

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 33 (1987)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE GALOIS GROUPS OF THE EXPONENTIAL TAYLOR POLYNOMIALS

Autor: Coleman, Robert F.

Kapitel: I. Review of the Newton Polygon

DOI: <https://doi.org/10.5169/seals-87891>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 02.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ON THE GALOIS GROUPS OF THE EXPONENTIAL TAYLOR POLYNOMIALS

by Robert F. COLEMAN

Let

$$f_n(x) = 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!}$$

denote the n^{th} Taylor polynomial of the Exponential Function.

In 1930, [S-2], Schur proved the following theorem about $f_n(x)$:

THEOREM (Schur). *The Galois group, G_n , of f_n is A_n , the alternating group on n letters, if 4 divides n and is S_n , the symmetric group on n letters, otherwise.*

In this note we shall give a proof different from Schur's. Our ulterior motive is to demonstrate the utility of Newton polygons.

We must:

- A. Show that f_n is irreducible.
- B. Show that G_n contains a p -cycle for any prime number p between $n/2$ and $n-2$.
- C. Calculate the discriminant, D_n , of f_n and determine when it is a square.

We need the following:

1. The main theorem about p -adic Newton polygons (see below).
2. Bertrand's Postulate [B], proven by Tschebyshev [T], which asserts that for each integer n , at least 8, there exists a prime number strictly between $n/2$ and $n-2$. (See also [H-W] Chapter 22.)
3. The theorem of Jordan which asserts that if G is a transitive subgroup of S_n which contains a p -cycle for some prime p strictly between $n/2$ and $n-2$ then G contains A_n . (See [J-1], Note C and [J-2], Theorem 1 or [Ha], Theorems 5.6.2 and 5.7.2.)
4. The fact that the Galois group of a polynomial of degree n is contained in A_n iff its discriminant is a square.

We shall use 1 for A and B . This will imply G_n is transitive and together with 2 and 3 will imply G_n contains A_n for $n \geq 8$. We shall use the differential equation satisfied by the exponential function and 2 again to perform C . Finally, we shall use 4 to complete the proof. (We shall also require liberal doses of Galois theory.)

I. REVIEW OF THE NEWTON POLYGON

Let


$$g(x) = a_0 + a_1x + \dots + a_kx^k$$

be a polynomial over \mathbf{Q}_p . Consider the points:

$$(i, \text{ord}(a_i)), \quad 0 \leq i \leq k,$$

in the Cartesian plane. The Newton polygon of g is defined to be the lower convex hull of these points.

Example. The Newton Polygon of $f_7(x)$ considered over \mathbf{Q}_2 , is



The main theorem about these polygons is:

THEOREM NP. Let $(x_0, y_0), (x_1, y_1), \dots, (x_p, y_p)$ denote the successive vertices of this polygon. Then over \mathbf{Q}_p , g factors as follows:

$$g(x) = g_1(x)g_2(x) \dots g_r(x)$$

where the degree of g_i is $x_i - x_{i-1}$ and all the roots of $g_i(x)$ in $\bar{\mathbf{Q}}_p$ have valuation $-\left(\frac{y_i - y_{i-1}}{x_i - x_{i-1}}\right)$.

We call the rational numbers, $\frac{y_i - y_{i-1}}{x_i - x_{i-1}}$, the slopes of g .

Example. The polynomial f_7 has three factors over \mathbf{Q}_2 , of degrees 4, 2 and 1, respectively, which have slopes $-3/4$, $-1/2$ and 0.

COROLLARY. Let d be a positive integer. Suppose that d divides the denominator of each slope (in lowest terms) of g . Then d divides the degree of each factor of g over \mathbf{Q}_p .

Proof. It suffices to show that d divides the degree of each irreducible factor of g . Let h be such a factor. Let $\alpha \in \bar{\mathbf{Q}}_p$ be a root of h . Since d divides the denominator of the valuation of α (by Theorem NP), it follows that d divides the index of ramification of the extension $\mathbf{Q}_p(\alpha)/\mathbf{Q}_p$ which divides the degree of the extension which equals the degree of h .

II. APPLICATION TO THE EXPONENTIAL TAYLOR POLYNOMIALS

Fix a prime number p .

LEMMA. Suppose k is a positive integer and

$$k = a_0 + a_1p + \dots + a_s p^s$$

where $0 \leq a_i < p$. Then

$$\text{ord}(k!) = \frac{k - (a_0 + a_1 + \dots + a_s)}{p - 1}.$$

This is easy and well known.

Now write

$$n = b_1 p^{n_1} + b_2 p^{n_2} + \dots + b_s p^{n_s}$$

where $n_1 > n_2 > \dots > n_s$ and $0 < b_i < p$. Let

$$x_i = b_1 p^{n_1} + \dots + b_i p^{n_i}.$$