

Zeitschrift: L'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: KNOTTING CODIMENSION 2 SUBMANIFOLDS LOCALLY
Kapitel: §3. Type 2 case
Autor: Masuda, Mikiya / Sakuma, Makoto
DOI: <https://doi.org/10.5169/seals-57360>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 09.11.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

§ 3. TYPE 2 CASE

In this section and the next section, we treat the case where a meridian of L^n in M^{n+2} is null homotopic in $M - L$. The following lemma follows from [Li, Lemma 1]. We shall give an alternative proof which is interesting by itself (the argument is also given in [Ms, Theorem 4.2]).

LEMMA 3.1. $I(S^n \times S^2, S^n \times \{*\}) = \mathcal{K}_n$ if $n \geq 3$.

Proof. Let (S^{n+2}, K) be an n -knot and consider $(S^n \times S^2, S^n \times \{*\}) \# (S^{n+2}, K)$. A subset $S^n \times \{*\} \cup K \cup \{x_0\} \times S^2$ ($x_0 \in S^n$) is exactly the wedge sum of S^n and S^2 . As easily observed the complement of an open regular neighborhood of the subset is contractible and hence diffeomorphic to D^{n+2} as $n + 2 \geq 5$. This means that one can express

$$(S^n \times S^2, S^n \times \{*\}) \# (S^{n+2}, K) = (S^n \times S^2, S^n \times \{*\}) \# \Sigma$$

where Σ is a homotopy $(n+2)$ -sphere and the connected sum at the right hand side is done away from the submanifold $S^n \times \{*\}$.

On the other hand the ambient manifold must be diffeomorphic to $S^n \times S^2$ because it is the connected sum of $S^n \times S^2$ with S^{n+2} . These mean that Σ belongs to the inertia group of $S^n \times S^2$. But the group is trivial ([Sc]), so Σ must be the standard sphere. This proves the lemma. Q.E.D.

We shall denote by $\langle m \rangle$ the class in $\pi_1(M - L)$ represented by a meridian of L in M .

LEMMA 3.2. Suppose M is spin, L is diffeomorphic to S^n , and $n \geq 3$. If $\langle m \rangle = 1$ for (M, L) , then $(M, L) = (S^n \times S^2, S^n \times \{*\}) \# M'$ with a closed oriented manifold M' of dimension $n + 2$.

Proof. Since $\langle m \rangle = 1$ and $\dim M \geq 5$, the meridian m bounds a 2-disk in $M - L$. Therefore $L \vee S^2$ is embedded in M . The normal bundle to L in M is trivial, because it is classified by the Euler class sitting in $H^2(L; \mathbf{Z})$ and $H^2(L; \mathbf{Z}) = 0$ as $L = S^n$ and $n \geq 3$. The normal bundle of the embedded S^2 is also trivial, because it is classified by the second Stiefel-Whitney class and it vanishes as M is spin. Hence the closed regular neighborhood of $L \vee S^2$ in M is diffeomorphic to that of $S^n \vee S^2$ naturally embedded in $S^n \times S^2$. In particular its boundary is diffeomorphic to S^{n+1} . This implies the lemma. Q.E.D.

Remark 3.3. A similar argument works even if M is not spin. But this time two cases arise according as the normal bundle of the embedded S^2 is trivial or not. If it is trivial, then the same conclusion as above holds. If it is not trivial, we have

$$(M, L) = (S^n \tilde{\times} S^2, S^n) \# M'.$$

Here $S^n \tilde{\times} S^2$ denotes the total space of the sphere bundle associated with the nontrivial $(n+1)$ -dimensional vector bundle over S^2 (note that it is unique as $\pi_1(SO(n+1)) \simeq Z_2$ for $n \geq 2$) and the submanifold S^n denotes a fiber.

Combining Lemma 3.1 with 3.2, we obtain

THEOREM 3.4. *Suppose M is spin, L is diffeomorphic to S^n , and $n \geq 3$. Then if $\langle m \rangle = 1$ for (M, L) , then $I(M, L) = \mathcal{K}_n$.*

Remark 3.5. If the inertia group $I(S^n \tilde{\times} S^2)$ is trivial, then the same argument as the proof of Lemma 3.1 proves that $I(S^n \tilde{\times} S^2, S^n) = \mathcal{K}_n$ and hence one could drop the spin condition for M by Remark 3.3.

If $L \neq S^n$, then the above argument does not work. For a general L we construct an s -cobordism between pairs $(M, L) \# (S^{n+2}, K)$ and (M, L) and apply lemma 1.6. We denote the set of all null-cobordant n -knots by \mathcal{K}_n^0 . According to Kervaire [K] (cf. [KW, Chap. IV]) $\mathcal{K}_n = \mathcal{K}_n^0$ if n is even, but $\mathcal{K}_n \neq \mathcal{K}_n^0$ if n is odd.

PROPOSITION 3.6. *Suppose $\langle m \rangle = 1$ for (M^{n+2}, L^n) and $n \geq 3$. Then $I_0(M, L)$ contains \mathcal{K}_n^0 . In particular, if n is even ≥ 4 , then $I_0(M, L) = I(M, L) = \mathcal{K}_n$.*

Proof. Let (S^{n+2}, K) bound a disk pair (D^{n+3}, D) , where D is a $(n+1)$ -disk. The boundary connected sum $(M, L) \times I \natural (D^{n+3}, D)$ at the 1-level gives a cobordism between (M, L) and $(M, L) \# (S^{n+2}, K)$.

We shall check the conditions (1) and (2) in Lemma 1.6 for this cobordism. First, since D is diffeomorphic to D^{n+1} , $L \times I \natural D$ is diffeomorphic to $L \times I$; so (1) is satisfied. Hence $E(L \times I \natural D)$ gives a cobordism relative boundary between $E(L)$ and $E(L \# K)$. We note that

$$(3.7) \quad E(L \times I \natural D) = E(L \times I) \cup E(D)$$

where $E(L \times I)$ and $E(D)$ are pasted together along $D^{n+1} \times S^1$ embedded in their boundaries. The S^1 factor corresponds to meridians of $L \times I$ and D . Then the van Kampen's theorem says that

$$\begin{aligned} \pi_1(E(L \times I \natural D)) &\simeq \pi_1(E(L \times I)) \underset{\langle m \rangle}{*} \pi_1(E(D)) \\ &\simeq \pi_1(E(L \times I)) * (\pi_1(E(D)) / \langle m \rangle) \end{aligned}$$

where the latter isomorphism is because $\langle m \rangle = 1$ in $\pi_1(E(L \times I))$ by the assumption. Since $\pi_1(E(D)) / \langle m \rangle \simeq \pi_1(D^{n+3}) \simeq \{1\}$, we have

$$(3.8) \quad \pi_1(E(L \times I \natural D)) \simeq \pi_1(E(L \times I)) \simeq \pi_1(E(L)).$$

Here the inclusion map $i: E(L) = E(L) \times \{0\} \rightarrow E(L \times I \natural D)$ induces the isomorphism.

We shall observe that i is a simple homotopy equivalence. For that purpose we consider the lifting of i to the universal covers. Since the map $\pi_1(E(D)) \rightarrow \pi_1(E(L \times I \natural D))$ induced by the inclusion map is trivial as observed above, it follows from (3.7) that

$$(3.9) \quad \tilde{E}(L \times I \natural D) = \tilde{E}(L \times I) \cup E(D) \times \Pi$$

where $\Pi = \pi_1(E(L \times I \natural D)) = \pi_1(M - L)$ and $\tilde{E}(L \times I)$ and $E(D) \times \Pi$ are pasted together Π -equivariantly along $D^{n+1} \times S^1 \times \Pi$ embedded in their boundaries. This means that $\tilde{i}_*: H_q(\tilde{E}(L); \mathbf{Z}) \rightarrow H_q(\tilde{E}(L \times I \natural D); \mathbf{Z})$ is an isomorphism as $\mathbf{Z}[\Pi]$ -modules. Hence $i_*: \pi_q(E(L)) \rightarrow \pi_q(E(L \times I \natural D))$ is an isomorphism by Namioka's theorem (see [W11, §4]) and hence i is a homotopy equivalence.

The assumption $\langle m \rangle = 1$ together with (3.9) tells us that the Whitehead torsion $\tau(i) \in Wh(\Pi)$ of the map i comes from an element of $Wh(1)$ through the map: $Wh(1) \rightarrow Wh(\Pi)$ induced from the inclusion $1 \rightarrow \Pi$. However $Wh(1) = 0$ and hence $\tau(i) = 0$. This shows that $E(L \times I \natural D)$ is an s -cobordism relative boundary. The proposition then follows from Lemma 1.6. Q.E.D.

Proposition 3.6 gives a complete answer to the case where n is even ≥ 4 . It would be interesting to ask if the same conclusion still holds in the case $n = 2$.

In the next section we will improve Proposition 3.6 when n is odd ≥ 5 .

§ 4. AN IMPROVEMENT

Throughout this section we assume n is odd ≥ 5 . Let V^{n+1} be a Seifert surface of an n -knot K in S^{n+2} . The normal bundle to V in S^{n+2} is trivial. We give the stable normal bundle of S^{n+2} a canonical framing so that V can be viewed as a framed manifold.