

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE ZERO-IN-THE-SPECTRUM QUESTION
Autor: LOTT, John
Kapitel: 4.1 One Dimension
DOI: <https://doi.org/10.5169/seals-87882>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 02.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2. If M is an irreducible noncompact globally symmetric space G/K , with $G = \text{Isom}(M)$ and K a maximal compact subgroup, then one can say more about the bottom of the spectrum. If $\text{rk}(G) = \text{rk}(K)$ then $\text{Ker} \left(\Delta_{\frac{\dim(M)}{2}} \right)$ is infinite-dimensional and the spectrum of Δ is bounded away from zero otherwise. If $\text{rk}(G) > \text{rk}(K)$ then $\text{Ker}(\Delta) = 0$ and $0 \in \sigma(\Delta_p)$ if and only if

$$p \in \left[\frac{\dim(M)}{2} - \frac{\text{rk}(G) - \text{rk}(K)}{2}, \frac{\dim(M)}{2} + \frac{\text{rk}(G) - \text{rk}(K)}{2} \right]$$

[19, Section VIIB].

Finally, we state a result about uniformly contractible Riemannian manifolds.

DEFINITION 6 [15, p. 29]. *A metric space Z has finite asymptotic dimension if there is an integer n such that for any $r > 0$, there is a covering $Z = \bigcup_{i \in I} C_i$ of Z by subsets of uniformly bounded diameter so that no metric ball of radius r in Z intersects more than $n + 1$ elements of $\{C_i\}_{i \in I}$. The smallest such integer n is called the asymptotic dimension $\text{asdim}_+(Z)$ of Z .*

PROPOSITION 8 (Yu [33]). *If M is a uniformly contractible Riemannian manifold with finite asymptotic dimension then $0 \in \sigma(\Delta_p)$ for some p .*

The proof of Proposition 8 uses methods of coarse index theory [28].

4. VERY LOW DIMENSIONS

In this section we show that the answer to the zero-in-the-spectrum question is “yes” for one-dimensional simplicial complexes and two-dimensional Riemannian manifolds.

4.1 ONE DIMENSION

As a one-dimensional manifold is either S^1 or \mathbf{R} , zero is clearly in the spectrum.

A more interesting problem is to consider a connected one-dimensional simplicial complex K . Let V be the set of vertices of K and let E be the set of oriented edges of K . That is, an element e of E consists of an edge of K and an ordering (s_e, t_e) of ∂e . We let $-e$ denote the same edge with the

reverse ordering of ∂e . For $x \in V$, let m_x denote the number of unoriented edges of which x is a boundary. We assume that $m_x < \infty$ for all x . Put

$$C^0(K) = \{f : V \rightarrow \mathbf{C} \text{ such that } \sum_{x \in V} m_x |f(x)|^2 < \infty\},$$

$$(4.1) \quad C^1(K) = \{F : E \rightarrow \mathbf{C} \text{ such that } F(-e) = -F(e) \text{ and } \frac{1}{2} \sum_{e \in E} |F(e)|^2 < \infty\}.$$

Then $C^0(K)$ and $C^1(K)$ are Hilbert spaces. The weighting used to define $C^0(K)$ is natural in certain respects [8].

There is a bounded operator $d : C^0(K) \rightarrow C^1(K)$ given by $(df)(e) = f(t_e) - f(s_e)$. Define the Laplace-Beltrami operators by $\Delta_0 = d^*d$ and $\Delta_1 = dd^*$. An element of $\text{Ker}(\Delta_1)$ is an $F \in C^1(K)$ such that for each vertex x the total current flowing into x vanishes, i.e. $\sum_{e \in E: t_e=x} F(e) = 0$.

The next proposition is essentially due to Gromov [15, p. 236], who proved it in the case when $\{m_x\}_{x \in V}$ is bounded.

PROPOSITION 9. $0 \in \sigma(\Delta_0)$ or $0 \in \sigma(\Delta_1)$.

Proof. As the nonzero spectra of d^*d and dd^* are the same, for our purposes it suffices to consider $\sigma(\Delta_0)$ and $\text{Ker}(\Delta_1)$. We argue by contradiction. Suppose that $0 \notin \sigma(\Delta_0)$ and $\text{Ker}(\Delta_1) = 0$. First, K must be infinite, as otherwise $\text{Ker}(\Delta_0) \neq 0$. Second, K must be a tree, as if K had a loop then we could create a nonzero element of $\text{Ker}(\Delta_1)$ by letting a current of unit strength flow around the loop.

We now show that K has lots of branching. For $x, y \in V$, let $[x, y]$ be the geodesic arc from x to y and let (x, y) be its interior. Let $d(x, y)$ be the number of edges in $[x, y]$.

LEMMA 5. *There is a constant $L > 0$ such that if $d(x, y) > L$ then there is an infinite subtree of K which intersects (x, y) but does not contain x or y .*

Proof. If the lemma is not true then for all $N > 1$, there are vertices x and y such that $d(x, y) > N$ but there are no infinite subtrees as in the statement of the lemma. In other words, the connected component C of $K - \{x\} - \{y\}$ which contains (x, y) is finite. As K is a tree, x is only connected to the vertices in C by a single edge, and similarly for y (see Fig. 5). Define $f \in C^0(K)$ by

$$(4.2) \quad f(v) = \begin{cases} 1 & \text{if } v \in C, \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$(4.3) \quad \frac{\langle df, df \rangle}{\langle f, f \rangle} \leq \frac{2}{2(d(x, y) - 1)} \leq \frac{1}{N}.$$

As N can be taken arbitrarily large, this contradicts the assumption that $0 \notin \sigma(\Delta_0)$. \square



FIGURE 5

It follows that K contains a subtree K' which is topologically equivalent to an infinite triadic tree, with the distances between branchings at most L (see Fig. 6). We can create a nonzero square-integrable harmonic 1-cochain F' on K' by letting a unit current flow through it, as in Fig. 6. Let $F \in C^1(K)$ be the extension of F' by zero to K . If x is a vertex of K' then the total current flowing into x is still zero, as no new current comes in along the edges of $K - K'$. Thus $\text{Ker}(\Delta_1) \neq 0$, which is a contradiction. \square




FIGURE 6