

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 43 (1997)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: POLYGON SPACES AND GRASSMANNIANS
Autor: Hausmann, Jean-Claude / Knutson, Allen
Kurzfassung
DOI: <https://doi.org/10.5169/seals-63276>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 01.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

POLYGON SPACES AND GRASSMANNIANS

by Jean-Claude HAUSMANN and Allen KNUTSON *)

ABSTRACT. We study the moduli spaces of polygons in \mathbf{R}^2 and \mathbf{R}^3 , identifying them with subquotients of 2-Grassmannians using a symplectic version of the Gel'fand-MacPherson correspondence. We show that the bending flows defined by Kapovich-Millson arise as a reduction of the Gel'fand-Cetlin system on the Grassmannian, and with these determine the pentagon and hexagon spaces up to equivariant symplectomorphism. Other than invocation of Delzant's theorem, our proofs are purely polygon-theoretic in nature.

1. INTRODUCTION

Let ${}^m\tilde{\mathcal{P}}^k$ be the space of m -gons in \mathbf{R}^k up to translation and positive homotheties (precise definitions in §2). This space comes with several structures: an action of $O(k)$, an action of S_m permuting the edges, and a function $\ell : {}^m\tilde{\mathcal{P}}^k \rightarrow \mathbf{R}^m$ taking a polygon ρ to the lengths of its edges (once the perimeter of ρ is fixed). The quotients of ${}^m\tilde{\mathcal{P}}^k$ by SO_k (or O_k) are the moduli spaces ${}^m\mathcal{P}_+^k$ (respectively, ${}^m\mathcal{P}^k$). Fixing a reflection in $O(k)$ provides an involution on ${}^m\tilde{\mathcal{P}}^k$ and ${}^m\mathcal{P}_+^k$ whose fixed point sets are ${}^m\tilde{\mathcal{P}}^{k-1}$ and ${}^m\mathcal{P}^{k-1}$. The goal of this paper is to understand the topology of these various spaces and the geometric structures that they naturally carry when $k = 2$ or 3 . They are closely related to more familiar objects (Grassmannians, projective spaces, Hopf bundles, etc.) The spaces ${}^m\mathcal{P}^k(\alpha) := \ell^{-1}(\alpha)$ of polygons with given side-lengths $\alpha \in \mathbf{R}^m$ are of particular interest.

The great miracle occurs when $k = 3$, because \mathbf{R}^3 is isomorphic to the space $I\mathbf{H}$ of pure imaginary quaternions, and the 2-sphere in \mathbf{R}^3 is Kähler. The tools of symplectic geometry can then be used. Most prominent is a

*) Both authors thank the Fonds National Suisse de la Recherche Scientifique for its support.