

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 45 (1999)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** GENERALIZED FØLNER CONDITION AND THE NORMS OF RANDOM WALK OPERATORS ON GROUPS  
**Autor:** UK, Andrzej  
**Kapitel:** 5. Lower bounds  
**DOI:** <https://doi.org/10.5169/seals-64454>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

**Download PDF:** 15.03.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## 5. LOWER BOUNDS

Now we will consider generalized Følner sequences for functions  $f$  such that

$$Pf \geq \|P\|f.$$

This will enable us to obtain some lower bounds on the norm of random walk operators on graphs.

As in Section 3, let  $X$  be a connected, locally finite graph and let  $P$  be the simple random walk operator on  $X$ .

In this section we will prove the following lower bound on the norm  $\|P\|$ :

**THEOREM 8.** *Let  $X$  be a graph such that at each vertex there are at most  $k$  edges. Then*

$$\|P\| \geq \frac{2\sqrt{k-1}}{k}.$$

The norm of the random walk operator  $\|P\|$  is equal to  $\frac{2\sqrt{k-1}}{k}$  for the random walk on the tree which has  $k$  edges at each vertex. In [9] Kesten proved this lower bound in the case of Cayley graphs.

*Proof of Theorem 8.* Let us consider a graph  $X$  such that at each vertex there are at most  $k$  edges. We can suppose that  $k \geq 3$  because for  $k = 2$  we obtain subgraphs of  $\mathbf{Z}$  or finite graphs, and necessarily  $\|P\| = 1$ . As it is enough to prove the desired bound for any connected component of  $X$ , we can suppose that  $X$  is connected.

In order to show that  $\|P\|$  is large enough, we will construct a sequence of functions  $f_n \in l^2(X, N)$  such that

$$\limsup_{n \rightarrow +\infty} \frac{\|Pf_n\|_{l^2(X, N)}}{\|f_n\|_{l^2(X, N)}} \geq \frac{2\sqrt{k-1}}{k}.$$

Let us endow the set of vertices of  $X$  with a metric. The distance between two vertices is the smallest number of edges needed to connect them. Let us choose a vertex  $e$  in  $X$  and for a vertex  $v$  let us denote by  $|v|$  its distance from  $e$ .

Let  $f$  be the unique (up to translations and multiplications) radial eigenfunction of  $P$  on the homogeneous tree of degree  $k$ , corresponding to the eigenvalue  $\frac{2\sqrt{k-1}}{k}$ , which is the norm of  $P$  on this tree, i.e.

$$(15) \quad f(v) = g(|v|) = \left( \frac{k-2}{k} |v| + 1 \right) \left( \frac{1}{\sqrt{k-1}} \right)^{|v|}.$$

Using (15) we can define  $f$  on  $X$ . We then prove

LEMMA 6. *For any vertex  $v \in X$ ,*

$$Pf(v) \geq \frac{2\sqrt{k-1}}{k} f(v).$$

*Proof.* If  $v = e$  the result is clearly true. Let us consider then a vertex  $v \in X$  such that  $n = |v| \geq 1$ . Let the number of neighbors of  $v$  which are at a distance  $n-1$  or  $n$  from  $e$  be equal respectively to  $p$  and  $q$ . So the number of neighbors of  $v$  which are at a distance  $n+1$  is equal to  $N(v) - p - q$ . Hence

$$Pf(v) = \frac{1}{N(v)} (pg(n-1) + qg(n) + (N(v) - p - q)g(n+1)).$$

As  $p \geq 1$  and  $g$  is a decreasing function,

$$Pf(v) \geq \frac{1}{N(v)} (g(n-1) + (N(v) - 1)g(n+1)).$$

As  $N(v) \leq k$  and  $g(n-1) \geq g(n+1)$ ,

$$Pf(v) \geq \frac{1}{k} (g(n-1) + (k-1)g(n+1)) = \frac{2\sqrt{k-1}}{k} g(n). \quad \square$$

Let us denote by  $S_n$  and  $B_n$  the vertices which are respectively at a distance  $n$  and less than or equal to  $n$ .

LEMMA 7.

$$\frac{\sum_{v \in S_{n+1}} f^2(v) N(v)}{\sum_{v \in B_n} f^2(v) N(v)} \xrightarrow{n \rightarrow \infty} 0.$$

*Proof.* As  $1 \leq N(v) \leq k$  it is enough to show that

$$\frac{\sum_{v \in S_{n+1}} f^2(v)}{\sum_{v \in B_n} f^2(v)} \xrightarrow{n \rightarrow \infty} 0.$$

Let us denote

$$a_n = \sum_{v \in S_n} f^2(v) = |S_n| g^2(n).$$

As  $|S_{n+1}| \leq (k-1)|S_n|$  one has

$$(16) \quad a_{n+1} = |S_{n+1}|g^2(n+1) \leq (k-1)|S_n|g^2(n+1) = \left(1 + \frac{k-2}{(k-2)n+k}\right)^2 a_n.$$

We have to show that

$$(17) \quad \frac{\sum_{v \in S_{n+1}} f^2(v)}{\sum_{v \in B_n} f^2(v)} = \frac{a_{n+1}}{a_1 + \dots + a_n} \rightarrow_{n \rightarrow \infty} 0.$$

It is a standard exercise to show that (16) implies (17).  $\square$

Let  $f_n$  be the sequence of functions which are restrictions of  $f$  to the vertices that are at a distance not greater than  $n$ :

$$f_n = f|_{B_n}.$$

By Lemma 6 and Lemma 7 it follows that

$$\limsup_{n \rightarrow +\infty} \frac{\|Pf_n\|_{l^2(X,N)}}{\|f_n\|_{l^2(X,N)}} \geq \frac{2\sqrt{k-1}}{k},$$

which proves Theorem 8.  $\square$

Some examples of upper bounds on the norm of the simple random walk operator on graphs and their comparison with the lower bound from Theorem 8 can be found in [22].

## REFERENCES

- [1] AOMOTO, K. and Y. KATO. Green functions and spectra on free products of cyclic groups. *Ann. Inst. Fourier (Grenoble)* 38, 1 (1988), 59–85.
- [2] BOUGEROL, PH. and L. ÉLIE. Existence of positive harmonic functions on groups and on covering manifolds. *Ann. Inst. H. Poincaré Probab. Statist.* 31, No. 1 (1995), 59–80.
- [3] CARTWRIGHT, D. I. and P. M. SOARDI. Random walks on free products, quotients and amalgams. *Nagoya Math. J.* 102 (1986), 163–180.
- [4] FØLNER, E. On groups with full Banach mean value. *Math. Scand.* 3 (1955), 243–254.
- [5] GOODMAN, F. M., P. DE LA HARPE and V. F. R. JONES. Coxeter graphs and towers of algebras. *Math. Sci. Res. Inst. Publ.* 14. Springer-Verlag (1989).
- [6] GROMOV, M. Groups of polynomial growth and expanding maps. *Inst. Hautes Études Sci. Publ. Math.* 53 (1981), 53–78.
- [7] KAIMANOVICH, V. Dirichlet norms, capacities and generalized isoperimetric inequalities for Markov operators. *Potential Anal.*, 1 (1992), 61–82.