

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 45 (1999)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: COUNTING PATHS IN GRAPHS
Autor: Bartholdi, Laurent
Kapitel: 1. Introduction
DOI: <https://doi.org/10.5169/seals-64442>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 29.03.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

COUNTING PATHS IN GRAPHS

by Laurent BARTHOLDI

ABSTRACT. We give a simple combinatorial proof of a formula that extends a result by Grigorchuk [Gri78a, Gri78b] relating cogrowth and spectral radius of random walks. Our main result is an explicit equation determining the number of ‘bumps’ on paths in a graph: in a d -regular (not necessarily transitive) non-oriented graph let the series $G(t)$ count all paths between two fixed points weighted by their length t^{length} , and $F(u, t)$ count the same paths, weighted as $u^{\text{number of bumps}} t^{\text{length}}$. Then one has

$$\frac{F(1-u, t)}{1-u^2 t^2} = \frac{G\left(\frac{t}{1+u(d-u)t^2}\right)}{1+u(d-u)t^2}.$$

We then derive the circuit series of ‘free products’ and ‘direct products’ of graphs. We also obtain a generalized form of the Ihara-Selberg zeta function [Bas92, FZ98].

1. INTRODUCTION

Let $\Gamma = \mathbf{F}_S/N$ be a group generated by a finite set S , where \mathbf{F}_S denotes the free group on S . Let f_n be the number of elements of the normal subgroup N of \mathbf{F}_S whose minimal representation as words in $S \cup S^{-1}$ has length n ; let g_n be the number of (not necessarily reduced) words of length n in $S \cup S^{-1}$ that evaluate to 1 in Γ ; and let $d = |S \cup S^{-1}| = 2|S|$. The numbers

$$\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{f_n}, \quad \nu = \frac{1}{d} \limsup_{n \rightarrow \infty} \sqrt[n]{g_n}$$

are called the *cogrowth* and *spectral radius* of (Γ, S) . The Grigorchuk Formula [Gri78b] states that

$$(1.1) \quad \nu = \begin{cases} \frac{\sqrt{d-1}}{d} \left(\frac{\alpha}{\sqrt{d-1}} + \frac{\sqrt{d-1}}{\alpha} \right) & \text{if } \alpha > \sqrt{d-1}, \\ \frac{2\sqrt{d-1}}{d} & \text{else.} \end{cases}$$

We generalize this result to a somewhat more general setting: we replace the group Γ by a regular graph \mathcal{X} , i.e. a graph with the same number of edges at each vertex. Fix a vertex \star of \mathcal{X} ; let g_n be the number of circuits (closed sequences of edges) of length n at \star and let f_n be the number of circuits of length n at \star with no backtracking (no edge followed twice consecutively). Then the same equation holds between the growth rates of f_n and g_n .

To a group Γ with fixed generating set one associates its Cayley graph \mathcal{X} (see Subsection 3.1). \mathcal{X} is a d -regular graph with distinguished vertex $\star = 1$; paths starting at \star in \mathcal{X} are in one-to-one correspondence with words in $S \cup S^{-1}$, and paths starting at \star with no backtracking are in one-to-one correspondence with elements of \mathbf{F}_S . A circuit at \star in \mathcal{X} is then precisely a word evaluating to 1 in Γ , and a circuit without backtracking represents precisely one element of N . In this sense results on graphs generalize results on groups. The converse would not be true: there are even graphs with a vertex-transitive automorphism group that are not the Cayley graph of a group [Pas93].

Even more generally, we will show that, rather than counting circuits and proper circuits (those without backtracking) at a fixed vertex, we can count paths and proper paths between two fixed vertices and obtain the same formula relating their growth rates.

These relations between growth rates are consequences of a stronger result, expressed in terms of generating functions. Define the formal power series

$$F(t) = \sum_{n=0}^{\infty} f_n t^n, \quad G(t) = \sum_{n=0}^{\infty} g_n t^n.$$

Then assuming \mathcal{X} is d -regular we have

$$(1.2) \quad \frac{F(t)}{1-t^2} = \frac{G\left(\frac{t}{1+(d-1)t^2}\right)}{1+(d-1)t^2}.$$

This equation relates F and G , and so relates *a fortiori* their radii of convergence, which are $1/\alpha$ and $1/(d\nu)$. We re-obtain thus the Grigorchuk Formula.

Finally, rather than counting paths and proper paths between two fixed vertices, we can count, for each $m \geq 0$, the number of paths with m backtrackings, i.e. with m occurrences of an edge followed twice in a row. Letting $f_{m,n}$ be the number of paths of length n with m backtrackings, consider the two-variable formal power series

$$F(u, t) = \sum_{m,n=0}^{\infty} f_{m,n} u^m t^n.$$

Note that $F(0, t) = F(t)$ and $F(1, t) = G(t)$. The following equation now holds:

$$\frac{F(1 - u, t)}{1 - u^2 t^2} = \frac{G\left(\frac{t}{1 + u(d - u)t^2}\right)}{1 + u(d - u)t^2}.$$

Setting $u = 1$ in this equation reduces it to (1.2).

A generalization of the Grigorchuk Formula in a completely different direction can be attempted: consider again a finitely generated group Γ , and an exact sequence

$$1 \longrightarrow \Xi \longrightarrow \Pi \longrightarrow \Gamma \longrightarrow 1,$$

where this time Π is not necessarily free. Assume Π is generated as a monoid by a finite set S . Let again g_n be the number of words of length n in Π evaluating to 1 in Γ , and let f_n be the number of elements of Ξ whose minimal-length representation as a word in S has length n . Is there again a relation between the f_n and the g_n ? In Section 8 we derive such a relation when Π is the modular group $\mathbf{PSL}_2(\mathbf{Z})$.

Again there is a combinatorial counterpart; rather than considering graphs one considers a locally finite cellular complex \mathcal{K} such that all vertices have isomorphic neighbourhoods. As before, g_n counts the number of paths of length n in the 1-skeleton of \mathcal{K} between two fixed vertices; and f_n counts elements of the fundamental groupoid, i.e. homotopy classes of paths, between two fixed vertices whose minimal-length representation as a path in the 1-skeleton of \mathcal{K} has length n . We obtain a relation between these numbers when \mathcal{K} consists solely of triangles and arcs, with no two triangles nor two arcs meeting; these are precisely the complexes associated with quotients of the modular group.

The original motivation for our research was the study of cogrowth in group theory [Gri78a]; however, as it turned out, the more general problem in graph theory has applications to other domains of mathematics, like the Ihara-Selberg zeta function and its evaluation by Hyman Bass [Bas92].

2. MAIN RESULT

Let \mathcal{X} be a graph, that may have multiple edges and loops. We make the following typographical convention for the power series that will appear: a series in the formal variable t is written $G(t)$, or G for short, and $G(x)$ refers to the series G with x substituted for t . Functions are written on the right, with $(x)f$ or x^f denoting f evaluated at x .