

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: IDEAL SOLUTIONS OF THE TARRY-ESCOTT PROBLEM OF DEGREE FOUR AND A RELATED DIOPHANTINE SYSTEM
Autor: Choudhry, Ajai
Kurzfassung
DOI: <https://doi.org/10.5169/seals-64802>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 02.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

IDEAL SOLUTIONS OF THE TARRY-ESCOTT PROBLEM OF DEGREE FOUR AND A RELATED DIOPHANTINE SYSTEM

by Ajai CHOUDHRY

ABSTRACT. In this paper, the complete ideal symmetric solution in integers of the Tarry-Escott problem of degree four, that is, of the system of simultaneous equations $\sum_{i=1}^s a_i^r = \sum_{i=1}^s b_i^r$, $r = 1, 2, 3, 4$, has been obtained. In addition, a parametric ideal non-symmetric solution has also been obtained, and this non-symmetric solution has been used to obtain a parametric solution of the diophantine system $\sum_{i=1}^s a_i^r = \sum_{i=1}^s b_i^r$, $r = 1, 2, 3, 4$ and 6.

1. INTRODUCTION

The Tarry-Escott problem of degree k consists of finding two sets of integers a_1, a_2, \dots, a_s and b_1, b_2, \dots, b_s such that

$$(1) \quad \sum_{i=1}^s a_i^r = \sum_{i=1}^s b_i^r, \quad r = 1, 2, \dots, k.$$

There is a well-known theorem [6, p.614] due to Frolov according to which the relations (1) imply that

$$(2) \quad \sum_{i=1}^s (Ma_i + K)^r = \sum_{i=1}^s (Mb_i + K)^r, \quad r = 1, 2, \dots, k,$$

where M and K are arbitrary integers. That is, if $(a_1, a_2, \dots, a_s; b_1, b_2, \dots, b_s)$ is a solution of the system (1), then

$$(Ma_1 + K, \dots, Ma_s + K; Mb_1 + K, \dots, Mb_s + K)$$

is also a solution of (1). This theorem is easily established by using the binomial theorem. If one solution of the system (1) is obtained from another through