

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	46 (2000)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	THE SPECTRAL MAPPING THEOREM, NORMS ON RINGS, AND RESULTANTS
Autor:	Laksov, D. / Svensson, L. / Thorup, A.
Kapitel:	4. The proof
DOI:	https://doi.org/10.5169/seals-64805

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 02.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3. A REDUCTION

To prove the Spectral Mapping Theorem it suffices to verify that it holds for the polynomial ring $k[t_0, \dots, t_m]$ in variables t_0, \dots, t_m over k , and for the polynomial $F(x) = t_0 + t_1x + \dots + t_mx^m$. This is because any polynomial $G(x) = b_0 + b_1x + \dots + b_mx^m$ with coefficients in a ring R containing k as a subring is the image of $F(x)$ by the map $g: k[t_0, \dots, t_m][x] \rightarrow R[x]$ defined by $g(a) = a$ for $a \in k$, by $g(t_i) = b_i$ for $i = 0, \dots, m$, and by $g(x) = x$. If we can prove the equality $\det F(M) = \prod_{i=1}^n F(\lambda_i)$ in $k[t_0, \dots, t_m]$ we obtain that $\det G(M) = g(\det F(M)) = \prod_{i=1}^n g(F(\lambda_i)) = \prod_{i=1}^n G(\lambda_i)$ in R .

4. THE PROOF

Clearly (2.1) holds when F is a constant a where it simply states that $\det(aI_n) = a^n$. We shall prove (2.1) for polynomials F of degree $m > 0$ by induction on m .

We first note that if $F(x)$ has a root λ in R , so that $F(x) = (x - \lambda)G(x)$ in $R[x]$, then (2.1) holds for $F(x)$. Indeed, $G(x)$ is of degree $m - 1$ so it follows from the induction hypothesis that $\det G(M) = \prod_{i=1}^n G(\lambda_i)$. Since $F(M) = (M - \lambda I_n)G(M)$ we obtain:

$$\begin{aligned} \det F(M) &= \det(M - \lambda I_n) \det G(M) \\ &= \prod_{i=1}^n (\lambda_i - \lambda) \prod_{i=1}^n G(\lambda_i) = \prod_{i=1}^n (\lambda_i - \lambda) G(\lambda_i) = \prod_{i=1}^n F(\lambda_i). \end{aligned}$$

As we saw in Section 3 it suffices to prove the result for the ring $Q = k[t_0, \dots, t_m]$ and the polynomial $F(x) = t_0 + t_1x + \dots + t_mx^m$. Let x and y be independent variables over the ring Q . The polynomial $F(x) - F(y)$ in x with coefficients in $Q[y]$ has the root $x = y$. Hence, as we just observed, (2.1) holds for the polynomial $F(x) - F(y)$. We obtain the equation:

$$(4.1) \quad \det(F(M) - F(y)I_n) = \prod_{i=1}^n (F(\lambda_i) - F(y))$$

in $Q[y]$.

The equation (2.1) is a consequence of (4.1). To see this we observe that $F(y)$ in $Q[y]$ is transcendent over Q , that is the element $F(y)$ in $Q[y]$ does not satisfy a polynomial relation $a_0 + a_1F(y) + \dots + a_lF(y)^l = 0$ with coefficients a_i in Q and $a_l \neq 0$, because the coefficient $a_l t_m^l$ of the highest

power y^{ml} of y that appears in the relation is non-zero. It follows that we can define a homomorphism of rings $h: Q[F(y)] \rightarrow Q$ by $h(a) = a$ for $a \in Q$, and $h(F(y)) = 0$. We apply the map h to both sides of (4.1) and obtain the equality (2.1).

5. NORMS ON ALGEBRAS

The only properties of determinants that we used in the proof of the Spectral Mapping Theorem is that they are multiplicative, functorial and homogeneous. It is therefore natural to place the proof into the more general framework of norms on algebras. The advantage of this point of view is that we obtain a deeper understanding of the Spectral Mapping Theorem, and we obtain a natural connection with resultants of polynomials.

A *norm* N of *degree* n on a, not necessarily commutative, k -algebra A is a family of maps $N_R: R \otimes_k A \rightarrow R$, one for every commutative k -algebra R , that satisfies the conditions :

- (1) $N_R(a \otimes 1) = a^n$ for all elements a in R .
- (2) $N_R(uv) = N_R(u)N_R(v)$ for all elements u and v of $R \otimes_k A$.
- (3) For every homomorphism $\varphi: R \rightarrow S$ of commutative k -algebras we have $\varphi N_R = N_S(\varphi \otimes \text{id}_A)$.

A norm on an algebra may be described as a *multiplicative homogeneous polynomial law* (see Roby [R], or [B1], §9, Définition 3, p. 52).

For any map $B \rightarrow A$ of k -algebras the norm N on A *restricts* to a norm on B of degree n . Moreover, for every homomorphism of commutative rings $k \rightarrow k'$ the norm N on A *induces* a norm of degree n on the k' -algebra $k' \otimes_k A$.

Let N be a norm of degree n on a k -algebra A . Denote by $k[t]$ the k -algebra of polynomials in the variable t with coefficients in k . For every element α in A the polynomial in $k[t]$:

$$P_\alpha(t) = P_\alpha^N(t) = N_{k[t]}(t - \alpha)$$

is called the *characteristic polynomial* of α . The *trace* $\text{Tr}^N(\alpha)$ of α is the element in k such that $-\text{Tr}^N(\alpha)$ is the coefficient of t^{n-1} in $P_\alpha(t)$.

We note that $P_\alpha(0) = (-1)^n N_k(\alpha)$.