

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 47 (2001)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE POSITIVE CONE OF SPHERES AND SOME PRODUCTS OF SPHERES
Autor: MATTHEY, Michel / SUTER, Ulrich
Kapitel: 4. The positive cone of the spheres
DOI: <https://doi.org/10.5169/seals-65432>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 02.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Let x and a be suitable generators of $\tilde{K}(S^6)$ and of $H^6(S^6; \mathbf{Z})$ respectively, and define $\bar{x} := q^*(x)$ and $\bar{a} := q^*(a)$. For obvious dimensional reasons, the Chern classes $c_1(\bar{x})$ and $c_2(\bar{x})$ vanish. Moreover, one has $c_3(\bar{x}) = q^*(c_3(x)) = q^*(2a) = 0$ (see Proposition 2.4), hence $c\text{-dim}(\bar{x}) = 0$. On the other hand, we have $\gamma^1(\bar{x}) = \bar{x} \neq 0$, so $\gamma\text{-dim}(\bar{x}) \geq 1$; more precisely, $\gamma^2(\bar{x})$ is $q^*(-S(3, 2) \cdot x) = q^*(-3x) = \bar{x} \neq 0$ and $\gamma^3(\bar{x}) = q^*(2S(3, 3) \cdot x) = 0$, so $\gamma\text{-dim}(\bar{x}) = 2$. Consequently, M is a connected finite CW-complex with a strict inclusion

$$K_\gamma(M) \subsetneq K_c(M).$$

iii) Let $Z = Y \vee M$ be the wedge of the preceding two examples. It is a 7-dimensional finite connected CW-complex for which none of $K_\gamma(Z)$ and $K_c(Z)$ contains the other one. (The product $Y \times M$ would also do.)

To end the present section, we prove that the cones are semigroups and homotopy invariants.

PROPOSITION 3.5. *The positive cone, the γ -cone and the c -cone of a connected finite CW-complex X are sub-semigroups of $K(X)$ and homotopy invariants of X . Moreover, the positive cone is a sub- λ -semiring of $K(X)$.*

Proof. The homotopy invariance is obvious for the three cones. We have already mentioned in the preliminaries that the positive cone is a sub-semiring of $K(X)$. It is also clear that it is a sub- λ -semiring. The ‘‘exponentiality’’ of γ_t and of c (the total Chern class) immediately show that the γ -cone and the c -cone are sub-semigroups of $K(X)$. \square

We do not know if in general the γ -cone and the c -cone are sub- λ -semirings of $K(X)$.

4. THE POSITIVE CONE OF THE SPHERES

We now intend to compute the positive cone of the spheres. For odd-dimensional spheres, there is nothing to do since $\tilde{K}(S^{2n+1}) = 0$. Whereas for even-dimensional spheres, one has $\tilde{K}(S^{2n}) = \mathbf{Z} \cdot x \cong \mathbf{Z}$, so we only have to compute $g\text{-dim}(lx)$ for all integers l .

By Proposition 2.4, we have

$$c(lx) = c(x)^l = (1 + (-1)^{n-1}(n-1)! \cdot a)^l = 1 + (-1)^{n-1}l(n-1)! \cdot a,$$

where a is the orientation class of S^{2n} . Therefore, by Proposition 3.2, we deduce that, for $l \neq 0$,

$$n = \text{c-dim}(lx) \leq \text{g-dim}(lx) \leq \dim(S^{2n})/2 = n,$$


and this shows that $\text{c-dim}(lx) = \text{g-dim}(lx) = n$. The sphere S^{2n} being a torsion-free space, the following theorem follows from Proposition 3.3.

THEOREM 4.1. *Let x be a generator of $\tilde{K}(S^{2n}) \cong \mathbf{Z}$. Then, for $l \in \mathbf{Z}$,*

$$\text{g-dim}(lx) = \begin{cases} 0 & \text{if } l = 0 \\ n & \text{otherwise.} \end{cases}$$

Moreover, the positive cone, the c -cone and the γ -cone of S^{2n} coincide:

$$K_+(S^{2n}) = K_c(S^{2n}) = K_\gamma(S^{2n}) = \mathbf{N} \times 0 \cup \{(l, x) \mid l \geq n\} \subset \mathbf{Z} \times \tilde{K}(S^{2n}).$$



There is another, purely homotopic, proof of the theorem. It is based on Bott's celebrated results on the homotopy groups of $BU(n)$ and Serre's computation of the rational homotopy groups of spheres. Let us also present this proof. We have

$$[S^{2n}, BU(k)] = \pi_{2n}(BU(k)) \quad \text{and} \quad \tilde{K}(S^{2n}) = [S^{2n}, BU] = \pi_{2n}(BU).$$

Consider the long exact sequence of the fibration $BU(k) \xrightarrow{i_k} BU$:

$$\dots \rightarrow \pi_{2n}(U/U(k)) \rightarrow \pi_{2n}(BU(k)) \xrightarrow{(i_k)_*} \pi_{2n}(BU) \rightarrow \pi_{2n-1}(U/U(k)) \rightarrow \dots$$

The fibre $U/U(k)$ of i_k is $2k$ -connected (see for example [MiTo], p.216) and it follows that $(i_k)_*$ is an isomorphism for $n \leq k$. According to Bott [Bott2], we have $\pi_{2n}(BU) \cong \mathbf{Z}$. It is well-known that for $k < n$, the group $\pi_{2n}(BU(k))$ is finite. Let us however give a short proof of this result.

LEMMA 4.2. For $m \geq 2k + 1$, the group $\pi_m(BU(k))$ is finite.

Proof. We fix $m \geq 3$. The fibration $BU(k-1) \rightarrow BU(k)$, with fibre S^{2k-1} , yields the following long exact sequence in homotopy:

$$\dots \rightarrow \pi_m(S^{2k-1}) \rightarrow \pi_m(BU(k-1)) \rightarrow \pi_m(BU(k)) \rightarrow \pi_{m-1}(S^{2k-1}) \rightarrow \dots$$

By Serre [Serre], $\pi_j(S^{2k-1})$ is finite for $j \neq 2k-1$, and we can conclude by induction over k (with $k \geq 1$ and $2k+1 \leq m$), since when $k=1$, one has $\pi_m(BU(1)) = \pi_{m-1}(U(1)) = 0$ for $m \geq 3$. \square

From this, we now infer that the image of $(i_k)_*$ is zero for $k < n$. This implies that $\text{g-dim}(lx) = n$ when $l \neq 0$, and concludes the second proof.

REMARK 4.3.

i) Since we were motivated by Elliott's classification of unital C^* -algebras of type AF by means of their K -theory, their positive cone and the K -theory class [1] of the unit (see [Bla1]), it is important to single out the fact that the positive cone of S^{2n} and that of S^{2m} are non-isomorphic as monoids if n is different from m . (There is no need here to distinguish the K -theory class 1 of the trivial one-dimensional bundle.) Let us provide a short proof of this claim. For $n \geq 1$, let M_n denote the positive cone of S^{2n} (identified as above with a sub-monoid of \mathbf{Z}^2 , in order to designate its elements). The abelian monoid M_n has a minimal set A_n of generators, in other words a generating set (as a monoid) that is contained in any other generating set, namely

$$A_n = \{(0, 1)\} \cup \{(k, n) \mid k \in \mathbf{Z} \setminus \{0\}\}.$$

Now, consider the function $\sigma: A_n \rightarrow \{2, 3, \dots\}$ defined, for $x \in A_n$, by

$$\sigma(x) := \min \{l \geq 2 \mid lx \text{ decomposes as a sum of elements of } A_n \setminus \{x\}\}.$$

Clearly, such an l exists for any $x \in A_n$ and $\sigma(A_n) = \{2, 2n\}$. Since A_n and σ are isomorphism invariants of M_n , this proves our claim.

ii) For odd-dimensional spheres the positive cone is "trivial"; in other words, $K(S^{2n-1}) = \mathbf{Z}$ and $K_+(S^{2n-1}) = \mathbf{N}$.

5. FURTHER PROPERTIES OF THE CONES

We now investigate naturality properties and behaviour under products of the positive cone, the γ -cone and the c -cone.

The following result is obvious.