

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	48 (2002)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	THE NONAMENABILITY OF SCHREIER GRAPHS FOR INFINITE INDEX QUASICONVEX SUBGROUPS OF HYPERBOLIC GROUPS
Autor:	Kapovich, Ilya
Kapitel:	4. Quasiconvex subgroups of hyperbolic groups
DOI:	https://doi.org/10.5169/seals-66081

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 02.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

4. QUASICONVEX SUBGROUPS OF HYPERBOLIC GROUPS

Detailed background information on quasiconvex subgroups of hyperbolic groups can be found in [1, 4, 20, 31, 38, 34, 32, 51, 54, 68] and other sources.

CONVENTION 4.1. Suppose G is a finitely generated group with a fixed finite generating set A . Let $X = \Gamma(G, A)$ be the Cayley graph of G with respect to A . We will denote the word-metric corresponding to A on X by d_A . Also, for $g \in G$ we will denote $|g|_A := d_A(1, g)$. For a word w in the alphabet $A \cup A^{-1}$ we will denote by \bar{w} the element of G represented by w .

DEFINITION 4.2 (Quasiconvexity). For $\epsilon \geq 0$ a subset Z of a metric space (X, d) is ϵ -quasiconvex if, for any $z_1, z_2 \in Z$ and any geodesic $[z_1, z_2]$ in X , the segment $[z_1, z_2]$ is contained in the closed ϵ -neighborhood of Z . A subset $Z \subseteq X$ is quasiconvex if it is ϵ -quasiconvex for some $\epsilon \geq 0$.

If G is a finitely generated group and A is a finite generating set of G , a subgroup $H \leq G$ is *quasiconvex in G with respect to A* if $H \subseteq \Gamma(G, A)$ is a quasiconvex subset.

It turns out [20, 32, 4, 31] that for subgroups of word-hyperbolic groups quasiconvexity is independent of the choice of a finite generating set for the ambient group. Thus a subgroup H of a hyperbolic group G is termed quasiconvex if $H \subseteq \Gamma(G, A)$ is quasiconvex for some finite generating set A of G .

We summarize some well-known basic facts regarding quasiconvex subgroups and provide some sample references:

PROPOSITION 4.3. *Let G be a word-hyperbolic group with a finite generating set A . Let $X = \Gamma(G, A)$ be the Cayley graph of G with the word-metric d_A induced by A . Then :*

1. *If $H \leq G$ is a subgroup, then either H is virtually cyclic (in which case H is called elementary) or H contains a free subgroup F of rank two which is quasiconvex in G (in this case H is said to be nonelementary) [20, 32].*
2. *Every cyclic subgroup of G is quasiconvex in G [1, 20, 32].*
3. *If $H \leq G$ is quasiconvex then H is finitely presentable and word-hyperbolic [1, 20, 32].*

4. Suppose $H \leq G$ is generated by a finite set Q inducing the word-metric d_Q on H . Then H is quasiconvex in G if and only if there is a $C > 0$ such that for any $h_1, h_2 \in H$

$$d_Q(h_1, h_2) \leq Cd_A(h_1, h_2)$$

(see [20, 32, 4, 31]).

5. The set \mathcal{L} of all A -geodesic words is a regular language that provides a bi-automatic structure for G . Moreover, a subgroup $H \leq G$ is quasiconvex if and only if H is \mathcal{L} -rational, that is the set $\mathcal{L}_H = \{w \in \mathcal{L} \mid \overline{w} \in H\}$ is a regular language [31].
6. If $H_1, H_2 \leq G$ are quasiconvex, then $H_1 \cap H_2 \leq G$ is quasiconvex [68].
7. [51, 46] Let $C \leq B \leq G$ where B is quasiconvex in G (and hence B is hyperbolic) and C is quasiconvex in B . Then C is quasiconvex in G [51, 46].
8. Let $C \leq B \leq G$ where C is quasiconvex in G and where B is word-hyperbolic. Then C is quasiconvex in B [51, 46].
9. Suppose $H \leq G$ is an infinite quasiconvex subgroup. Then H has finite index in its commensurator $\text{Comm}_G(H)$ (see [51]), where $\text{Comm}_G(H) := \{g \in G \mid [H : g^{-1}Hg \cap H] < \infty \text{ and } [g^{-1}Hg : g^{-1}Hg \cap H] < \infty\}$.

Part 1 of the above proposition implies that a nonelementary subgroup of a hyperbolic group is nonamenable.

5. PROOF OF THE MAIN RESULT

Let G be a nonelementary word-hyperbolic group with a finite generating set A . Let $X = \Gamma(G, A)$ be the Cayley graph of G with the word metric d_A . Let $\delta \geq 1$ be an integer such that the space $(\Gamma(G, A), d_A)$ is δ -hyperbolic. Let $H \leq G$ be a quasiconvex subgroup of infinite index in G . These conventions, unless specified otherwise, will be fixed for the remainder of the paper.

We shall need the following useful fact:

LEMMA 5.1. *There exists an integer constant $K = K(G, H, A) > 0$ with the following properties.*

Assume $g \in G$ is shortest with respect to d_A in the coset class Hg . Then for any $h \in H$ we have $(g, h)_1 \leq K$ (and hence $(g, H)_1 \leq K$).