

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA
Autor: Etingof, Pavel / Strickland, Elisabetta
Kapitel: 2.3 The quantum Calogero-Moser System
DOI: <https://doi.org/10.5169/seals-66677>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 02.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The Calogero-Moser system has a generalization to arbitrary Coxeter groups. Namely, consider a finite group W generated by reflections acting on the space \mathfrak{h} , and keep the notation of the previous section. Fix a W -invariant nondegenerate scalar product $(-, -)$ on \mathfrak{h} . It determines a scalar product on \mathfrak{h}^* . Define the “energy function”

$$E(x, p) = \frac{(p, p)}{2} + \frac{1}{2} \sum_{s \in \Sigma} \frac{\gamma_s(\alpha_s, \alpha_s)}{\alpha_s(x)^2}, \quad x \in \mathfrak{h}, \quad p \in \mathfrak{h}^*$$

on $T^*\mathfrak{h} = \mathfrak{h} \times \mathfrak{h}^*$, where $\gamma: \Sigma \rightarrow \mathbf{C}$ is a W -invariant function. Notice that although α_s is defined up to a non zero constant, by homogeneity, E is independent of the choice of α_s . We will call the system defined by E the Calogero-Moser system for W .

If W is the symmetric group S_n , $\mathfrak{h} = \mathbf{C}^n$, then Σ is the set of transpositions $s_{i,j}$, $i < j$, and we can take $\alpha_s = e_i - e_j$. Then we clearly obtain the usual Calogero-Moser system.

Below we will see that the Calogero-Moser system for W is completely integrable.

2.3 THE QUANTUM CALOGERO-MOSER SYSTEM

Let us now discuss quantization of the Calogero-Moser system. We start by quantizing the energy E by formally making the substitution

$$p_j \Rightarrow -i\hbar \frac{\partial}{\partial x_j},$$

where \hbar is a parameter (Planck's constant). This yields the Schrödinger operator

$$\widehat{E} := -\frac{\hbar^2}{2} \Delta + \frac{1}{2} \sum_{s \in \Sigma} \frac{\gamma_s(\alpha_s, \alpha_s)}{\alpha_s^2},$$

where Δ denotes the Laplacian.

In particular, in the case of $W = S_n$ we have

$$\widehat{E} = -\frac{\hbar^2}{2} \Delta + \sum_{i < j} \frac{c}{(x_i - x_j)^2},$$

where $\Delta = \sum_i \frac{\partial^2}{\partial x_i^2}$. Setting $\beta_s = \frac{\gamma_s}{2\hbar^2}$, we will from now on consider the operator

$$H := -\frac{2}{\hbar^2} \widehat{E} = \Delta - \sum_{s \in \Sigma} \frac{\beta_s(\alpha_s, \alpha_s)}{\alpha_s^2(x)},$$

called the Calogero-Moser operator.

We want to study the stationary Schrödinger equation:

$$(3) \quad H\psi = \lambda\psi, \quad \lambda \in \mathbf{C}.$$

As in the classical case, it is difficult to say anything explicit about solutions of this equation for a general Schrödinger operator H , but for the Calogero-Moser operator the situation is much better.

DEFINITION 2.1. A *quantum integral* of H is a differential operator M such that

$$[M, H] = 0.$$

We are going to show that there are many quantum integrals of H , namely that there are n commuting algebraically independent quantum integrals M_1, \dots, M_n of H . By definition, this means that the quantum Calogero-Moser system is completely integrable.

Once we have found M_1, \dots, M_n , observe that for fixed constants μ_1, \dots, μ_n , the space of solutions of the system

$$\begin{cases} M_1\psi = \mu_1\psi \\ \dots\dots\dots \\ M_n\psi = \mu_n\psi \end{cases}$$

is clearly stable under H . We will see that this space is in fact finite dimensional. Therefore, the operators M_i allow one to reduce the problem of solving the partial differential equation $H\psi = \lambda\psi$ to that of solving a system of ordinary linear differential equations. This phenomenon is called quantum complete integrability.

2.4 THE ALGEBRA OF DIFFERENTIAL-REFLECTION OPERATORS .

We are now going to explain how to find quantum integrals for H , using the Dunkl-Cherednik method.

First let us fix some notation. Given a smooth affine variety X , we will denote by $\mathcal{D}(X)$ the ring of differential operators on X . We are going to consider the case in which X is the open set U in \mathfrak{h} which is the complement of the divisor of the equation $\delta(x) := \prod_{s \in \Sigma} \alpha_s(x)$. Clearly $\mathcal{D}(U) = \mathcal{D}(\mathfrak{h})[1/\delta(x)]$.

LEMMA 2.2. An element of $\mathcal{D}(U)$ is completely determined by its action on $\mathbf{C}[U]^W = \mathbf{C}[U/W]$.