

**Zeitschrift:** Helvetica Physica Acta  
**Band:** 23 (1950)  
**Heft:** [3]: Supplementum 3. Internationaler Kongress über Kernphysik und Quantenelektrodynamik  
  
**Artikel:** Filter methods in neutronspectroscopy  
**Autor:** Vries, Hl. de  
**DOI:** <https://doi.org/10.5169/seals-422269>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 16.03.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Filter methods in neutronspectroscopy

by **Hl. de Vries**, Groningen, Holland.

In the last ten years several direct methods of neutronspectroscopy have been developed. (Mechanical and electrical velocityselectors, crystalspectrometers.) In some cases, however, the older indirect methods of filtering are still of interest. Indeed, optical spectroscopy with coloured glasses would be impossible, but it should be mentioned that the absorption bands for slow neutrons are relatively much broader than spectral lines of free atoms and this enables a study of resonance levels without "neutron spectroscopes". Some recent measurements will be described of the width and the positions of resonance levels which were made in Groningen\*). To explain the essential features of the method used, we confine ourselves first to the measurement of the shape of the resonance level of Ag. By an Ag absorber a "hole" is made in the energyspectrum of a beam of neutrons. After that this beam is reflected by a scattering substance. The energy loss in the elastic collisions is

$$\Delta E = \frac{2E(1 - \cos \varphi)}{A}.$$

In this way the hole in the spectrum is shifted to a lower energy and if  $\Delta E$  is large enough the intensity of resonance neutrons as detected by a silverfoil may be just as large as if no hole had been made in the spectrum; this depends on the value of  $\Delta E$  relative to the levelwidth. With decreasing  $\Delta E$  (increasing  $A$ ) the activity of the Ag detector decreases and from its dependance on  $\Delta E$  the shape of the resonance level can be determined. It is essential for the method (which will be called the *displacement method*) that the values of  $\Delta E$ , given above, happened to be of the order of magnitude of the width of the resonance levels, if we confine ourselves to values of  $E$  below 100 eV. The same technique was also used in studies of the overlapping of levels. In that case the detector and the absorber which made the "hole" were different elements.

---

\*) See Physica **8**, 825 (1941); **10**, 281 (1943); **10**, 299 (1943); **10**, 312 (1943); **10**, 381 (1943); **11**, 345 (1943); **11**, 396 (1946); **11**, 481 (1946); Nature **159**, 569 (1947).

It can be said that at present the resolving power of this method is at least as good as the resolving power of velocity selectors.

Some results are summarised in the tables I, II and in Fig. 1.




Fig. 1.

Activity of Ag detector as a function of the energy-loss  $\Delta E$  by the reflecting collision with nuclei of various elements. ( $b$  is a parameter proportional to  $\Delta E$ , containing also  $T$ . The activity caused by scattered neutrons ( $f_\lambda(b)$ ) is divided by  $f_\lambda(\infty)$ , the activity which was found when no hole had been made in the spectrum in order to account for the different reflecting powers of the various reflectors.  $G$  = curve predicted by a Gaussian form of the resonance level.  $BW$  = predicted by the BREIT-WIGNER formula. The value of the level with  $T$  has already been adjusted to give the best fit.

**Table I.**

Resonance energies (eV).  
 Second row: our results (BORON method).  
 Third row: our results (overlapping).  
 Fourth row: american reports.

Element . . . .	Ag <sup>108</sup>	Au <sup>198</sup>	Ag <sup>110</sup> <sub>(a)</sub>	Sb	Ag <sup>110</sup> <sub>(b)</sub>	Cd <sup>117</sup>	Cu <sup>66</sup>	Cu <sup>64</sup>	Zn <sup>69</sup>	Al <sup>28</sup>
Reson. energy .	23	4.6	5.6		90	110	100	140	500	60
Energy difference		0.31	0.7							
Time of flight .	16	4.8	5.1	5.8						

**Table II.**Results of various measurements of the level width  $\Gamma$ . $\sigma_0$  = absorption coefficient for resonance neutrons.

(Time of flight method from american reports.)

Method	Direct experimental result	Results for	
		Ag <sup>110</sup> (5.1 eV)	Au <sup>198</sup> (4.8 eV)
1. Displacement method	$\Gamma$	0.18	
2. Time of flight . . . .	$\sigma_0 \Gamma^2$	0.18	0.15 <sup>5</sup>
3. One level formula (BREIT-WIGNER)	$\sigma_0 \Gamma^2$		0.16
4. Activability . . . . .	$\sigma_0 \Gamma$		0.16 <sup>5</sup>
5. Overlapping . . . . .	$\Gamma, \frac{K_{12}}{K_{11}}, \frac{K_{21}}{K_{22}}$	0.18	0.17