

**Zeitschrift:** Helvetica Physica Acta

**Band:** 41 (1968)

**Heft:** 6-7

**Artikel:** Effective masses and curvature of the Fermi surface or energy bands

**Autor:** Jan, J.-P.

**DOI:** <https://doi.org/10.5169/seals-113954>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 16.03.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

tion des parois pouvant être aléatoires dans une large mesure. Ceci pourrait permettre un jour une application à l'étude de corps organiques non cristallins, formés par juxtaposition de chaînes de polymères.

Toutefois, les méthodes relativement nouvelles de canalisation et de blocage ont été appliquées jusqu'à présent surtout à certains métaux (Au, Cu, W, Fe) et à quelques semi-conducteurs (Ge, Si, GaAs, GaSb) et il semble bien qu'une extension aux autres substances inorganiques intéressant actuellement le physicien du solide promette de livrer des renseignements nouveaux et précieux sur la constitution de la matière.

### Bibliographie

- [1] J. C. PFISTER, *J. Phys.* **27**, 253–256 (1966).
- [2] A. F. TULINOV, *Soviet Physics-Uspekhi* **8**, 864–872 (1966).
- [3] W. BRANDT, *Sci. American* **278**, 91–98 (March 1968).
- [4] S. DATZ, C. ERGINSOY, G. LEIBFRIED, H. O. LUTZ, *Ann. Rev. Nucl. Sci.* **17**, 129–188 (1967).
- [5] G. DEARLANEY, M. A. WILKINS, *J. Sci. Instr.* **44**, 880–881 (1967).
- [6] H. O. LUTZ, S. DATZ, C. D. MOAK, T. S. NOGGLE, *Phys. Rev. Lett.* **17**, 285–287 (1966).
- [7] H. MATZKE, J. A. DAVIES, *J. appl. Phys.* **38**, 805–808 (1967).
- [8] V. S. KULIKAUSKAS, M. M. MALOV, A. F. TULINOV, *Zh. Eksper. Teor. Fiz.* **53**, 487–489 (1967).
- [9] J. U. ANDERSEN, J. A. DAVIES, K. O. NIELSEN, S. L. ANDERSEN, *Nucl. Instr. Meth.* **38**, 210–215 (1965).
- [10] M. W. THOMPSON, *Phys. Rev. Lett.* **13**, 756–760 (1964).
- [11] E. BØGH, *Phys. Rev. Lett.* **19**, 61–63 (1967).

## Effective Masses and Curvature of the Fermi Surface or Energy Bands

by J.-P. Jan

Division of Pure Physics, National Research Council of Canada, Ottawa, Canada

(26. I. 68)

*Abstract.* The curvature of the energy function on one hand, and of the Fermi surface on the other hand, is examined in relation to the effective mass tensor.

### 1. Introduction

This paper is of a didactic nature. It proposes to examine the components of the effective mass tensor of electrons in a crystal lattice. They involve second derivatives of the energy function and are therefore somehow related to curvatures, but what curvatures are involved is not always clear. We shall examine both the Fermi surface, and the curves representing the energy in function of the magnitude of the wave-vector in a given direction (energy bands), and the relationships we shall establish will show that in both cases the effective mass tensor is involved in a rather complicated way.

## 2. Effective Mass and Curvature of the Fermi Surface

If  $E(\mathbf{k})$  is the energy function, and  $E_F$  the Fermi energy, the equation of the Fermi surface is

$$E(\mathbf{k}) = E_F. \quad (1)$$

This is an implicit equation which does not lend itself easily to a local study of curvatures. Indeed, textbooks on differential geometry of surfaces prefer to adopt either an explicit form such as  $z = z(x, y)$  (in coordinates  $x, y, z$ ), or a parametric form  $x = x(u, v)$ ,  $y = y(u, v)$ ,  $z = z(u, v)$  involving two parameters  $u$  and  $v$ . The various curvatures can be easily derived in terms of these last equations. However, the inverse effective mass energy tensor

$$m_{ij}^{-1} = \frac{1}{\hbar^2} \frac{\partial^2 E}{\partial k_i \partial k_j} \quad (2)$$

is defined in terms of the implicit Equation (1), and its relation to the curvatures is not immediately obvious.

It should be noted also that the principal axes of the tensor (2) bear no simple relation to the local geometry of the Fermi surface. This is quite evident, for instance, in the case of the ellipsoidal surfaces

$$E(\mathbf{k}) = \frac{\hbar^2 k_1^2}{2 m_{11}} + \frac{\hbar^2 k_2^2}{2 m_{22}} + \frac{\hbar^2 k_3^2}{2 m_{33}} \quad (3)$$

for which the principal axes are at any point parallel to the coordinate axes.

A local study based on Equation (1) would make use of an expansion of the energy function to second order terms around a given point  $\mathbf{k}^0$ :

$$E(\mathbf{k}) = E(\mathbf{k}^0) + (k_i - k_i^0) \hbar v_i + \frac{1}{2} (k_i - k_i^0) (k_j - k_j^0) \hbar^2 m_{ij}^{-1} \quad (4)$$

where we have introduced the electron velocity

$$\mathbf{v} = \frac{1}{\hbar} \nabla_{\mathbf{k}} E \quad (5)$$

and adopted the standard summation convention over repeated indexes. When  $E(\mathbf{k})$  has the form (4), (1) represents a quadric (ellipsoid, hyperboloid or paraboloid). We now change our coordinate system to a new set  $(k'_1, k'_2, k'_3)$ , with the origin at point  $\mathbf{k}^0$ , the  $k'_3$  axis perpendicular to the Fermi surface in the direction of increasing energy, and the axes  $k'_1$  and  $k'_2$  chosen in the tangent plane to make  $(m_{12}^{-1})' = (m_{21}^{-1})' = 0$  (cf. PIPPARD [1]), and (4) becomes a new function

$$\begin{aligned} E'(\mathbf{k}') &= E'(0) + k'_3 \hbar v + \frac{1}{2} k_1'^2 \hbar^2 (m_{11}^{-1})' + \frac{1}{2} k_2'^2 \hbar^2 (m_{22}^{-1})' + \frac{1}{2} k_3'^2 \hbar^2 (m_{33}^{-1})' \\ &+ k'_1 k'_3 \hbar^2 (m_{13}^{-1})' + k'_2 k'_3 \hbar^2 (m_{23}^{-1})'. \end{aligned} \quad (6)$$

The principal radii of curvature of the Fermi surface at  $\mathbf{k}^0$  are therefore those of the paraboloid

$$k'_3 = - \frac{\hbar}{2 v m'_{11}} k_1'^2 - \frac{\hbar}{2 v m'_{22}} k_2'^2 \quad (7)$$

where  $m'_{11} = 1/(m_{11}^{-1})'$  and  $m'_{22} = 1/(m_{22}^{-1})'$ , namely

$$Q_1 = v m'_{11} / \hbar, \quad Q_2 = v m'_{22} / \hbar. \quad (8)$$

A sign convention is implicit in (8). The choice of  $k'_3$  makes  $v = v'_3$  positive. A positive  $m'_{11}$  or  $m'_{22}$  corresponds to a positive radius and to the center of curvature «inside» the Fermi surface, meaning that, if we move from  $\mathbf{k}^0$  towards the center of curvature, the energy decreases. This corresponds to a convex curve on the Fermi surface. The opposite holds for negative masses;  $m'_{11}$  and  $m'_{22}$  may of course have opposite signs, for instance at the necks of the Fermi surface of the noble metals.

The simple Equations (8) require the change of axes described above, and the components of the original inverse effective mass tensor (2) are not simply related to any curvature of the Fermi surface. It should also be noted that the components of the transformed tensor, which do not appear in (7) and (8), have nothing to do with the curvature of the Fermi surface; as PIPPARD [1] points out, the essential quantities are  $\rho_1$ ,  $\rho_2$  and  $v$ .

The preceding remarks apply not only to the Fermi surface, but to any energy surface.

### 3. Effective Mass and Curvature of the Energy Bands

In the one-dimensional case, the effective mass is

$$\frac{1}{m} = \frac{1}{\hbar^2} \frac{d^2 E}{dk^2}. \quad (9)$$

By energy band, we mean here the curve obtained by plotting  $E$  vs.  $k$ . Its curvature is given by

$$\frac{1}{\rho} = \frac{E''}{(1 + E'^2)^{3/2}} = \frac{\hbar^2/m}{(1 + \hbar^2 v^2)^{3/2}} \quad (10)$$

where  $E' = \hbar v$  and  $E'' = \hbar^2/m$  are the first and second derivatives of  $E(k)$  respectively, and  $\rho$  the radius of curvature. This curvature is seen to involve  $m$  directly, but also  $v$ . For instance, in a parabolic band,  $m$  is independent of  $k$ , but the curvature is not.

We generalize now these elementary remarks to three dimensions. Here, we consider how  $E$  changes in  $\mathbf{k}$ -space along a straight line whose direction is defined by a unit vector  $\mathbf{n}$ , such that  $\mathbf{k} = \mathbf{k}^0 + \mathbf{n} k$ . Such straight lines join points of high symmetry in the Brillouin zone in the popular energy band diagrams. It is easily shown that the curvature of  $E(k)$  is

$$\frac{1}{\rho} = \frac{\hbar^2/m_{av}}{(1 + \hbar^2 v^2 \cos^2 \alpha)} \quad (11)$$

where the average effective mass in the direction  $\mathbf{n}$  is defined by

$$\frac{1}{m_{av}} = n_i n_j m_{ij}^{-1} \quad (12)$$

and  $\alpha$  is the angle between  $\mathbf{v}$  and  $\mathbf{n}$ . It is seen that the effective mass tensor, through the average (12), comes directly into the curvature of the energy bands. For instance, in the [100] direction, only the  $m_{11}^{-1}$  component is involved; several components come into play in more complicated crystallographic directions.

The author is grateful to Dr. W. B. PEARSON for reading the manuscript.

### Bibliography

- [1] A. B. PIPPARD, *The Dynamics of Conduction Electrons*, in: *Low Temperature Physics* (Les Houches 1961); C. DE WITT et al. ed., Gordon and Breach, (New York 1962), or Blackie & Son Limited (London and Glasgow 1965).