

Zeitschrift: Helvetica Physica Acta
Band: 69 (1996)
Heft: 3

Artikel: Quasi-Einstein metrics and their renormalizability properties
Autor: Chave, Thierry / Valent, Galliano
DOI: <https://doi.org/10.5169/seals-116953>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 17.03.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Quasi-Einstein metrics and their renormalizability properties

By Thierry CHAVE, Galliano VALENT

Laboratoire de Physique Théorique et des Hautes Energies,
Unité associée au CNRS URA 280, Université Paris 7,
2 Place Jussieu, 75251 Paris Cedex 05

Abstract. We construct a family of kählerian quasi-Einstein metrics with an isometry group $U(n)$ acting linearly on the holomorphic coordinates. Suitable restrictions on the parameters give rise to complete non-compact as well as compact metrics whose geometrical structure is studied in detail. And we discuss The two loop renormalizability properties of the bosonic σ -models.

1 Introduction

Quasi-Einstein metrics are defined by the constraints

$$Ric_{\mu\nu} = \lambda g_{\mu\nu} + \frac{1}{2}(D_{\mu}v_{\nu} + D_{\nu}v_{\mu}). \quad (1.1)$$

In section 2, using Kähler geometry and an isometry group $U(n)$, we reduce (1.1) to a non-linear differential equation. Since this equation cannot be integrated to get the Kähler potential we switch to non-holomorphic coordinates which are useful to display explicitly the distance.

In section 3 we give two classes of complete non-compact metrics and we describe the compact metrics of the family.

In section 4 we discuss the two-loop renormalizability of the bosonic σ -models built on kählerian quasi-Einstein metrics. Our main result is that in the chosen isometry class the requirement of two-loop renormalizability selects uniquely CP^n , which is not only Einstein but symmetric!

2 Quasi-Einstein Kähler metrics

2.1 Isometry group $U(n)$

The idea to obtain explicit quasi-Einstein metrics is to use holomorphic coordinates $\{z_i, i = 1, \dots, n\}$ in $\mathbb{C}^n \setminus \{0\}$ and to choose for isometry group a $U(n)$ for which $\{z_i\}$ transform according to its fundamental representation. This implies that the Kähler potential K depends solely on $s = \bar{z} \cdot z = \sum_{i=1}^n \bar{z}_i z_i$. It follows that the metric is

$$\frac{1}{2} g = g_{i\bar{j}} dz^i d\bar{z}^j = A d\bar{z} \cdot dz + A' |\bar{z} \cdot dz|^2, \quad A(s) = \frac{dK(s)}{ds}.$$

The Ricci tensor reads

$$Ric_{i\bar{j}} = -\partial_{i\bar{j}}^2 \ln D, \quad D = \det(g_{i\bar{j}}) = A^{n-1}(sA)'.$$

The restrictions put on the metric by the quasi-Einstein requirement (1.1) imply ([1])

$$v_i = c \partial_i(sA), \quad A^{n-1}(sA)' = e^{-\lambda K - (Re c)sA}. \tag{2.1}$$

where c is an a priori complex integration constant.

2.2 The coordinates choice

In view of the complexity of (2.1), there is little hope to get the explicit Kähler potential $K(s)$ and the distance if we insist on using holomorphic coordinates. We define

$$z_i = \sqrt{s} \xi_i \quad i = 1, \dots, n$$

and

$$\xi = \frac{e^{i\tau}}{1 + \bar{u} \cdot u} (1, u), \quad \tau \in [0, 2\pi], \quad \{u_i, i = 1, \dots, n - 1\}.$$

This gives for the final form of our distance

$$\frac{1}{2} g = \frac{(dt)^2}{4\varrho} + \varrho (d\tau + \theta)^2 + t \frac{h}{2} (CP^{n-1})$$

with

$$\varrho = s \frac{dt}{ds}, \quad t = s K', \quad \theta = \frac{1}{1 + \bar{u} \cdot u} \frac{\bar{u} \cdot du - u \cdot d\bar{u}}{2i}.$$

The differential equation (2.1) becomes

$$\frac{d}{dt} (e^{ct} t^{n-1} \varrho(t)) = e^{ct} (n t^{n-1} - \lambda t^n). \tag{2.2}$$

The vector v is given by the holomorphic 1-form $v = c dt$ for some real constant c .

3 Complete and compact metrics

Integrating (2.2) gives

$$t^{n-1} \varrho = \frac{e^{-ct} \int_0^{ct} e^u \left(n u^{n-1} - \frac{\lambda}{c} u^n \right) du}{c^n} - B e^{-ct}, \quad n = 1, 2, \dots \tag{3.1}$$

3.1 Complete non-compact metrics

The first one appears for $B = 0, c > 0, \lambda < 0$.

At $t = 0$, setting $t = r^2$ the distance becomes

$$\frac{1}{2} g \approx dr^2 + r^2 \left[(d\tau + \theta)^2 + \frac{h}{2} (CP^{n-1}) \right] = dr^2 + r^2 d\bar{\xi} \cdot d\xi$$

which shows that $r = 0$ is a “nut”, [4].

For $t \rightarrow \infty$, setting $t = \frac{|\lambda|}{c} r^2$ the distance becomes

$$\frac{1}{2} g \approx dr^2 + r^2 \left\{ \left(\frac{|\lambda|}{c} \right)^2 |\bar{u} \cdot du|^2 + \frac{|\lambda|}{c} (d\bar{u} \cdot du - |\bar{u} \cdot du|^2) \right\}$$

showing that infinity is asymptotically flat (albeit not euclidean).

For $\lambda = 0$ the infinity is taubian, [2, p. 252].

The second one appears for $B > 0, c > 0, \lambda \leq 0$. But now t_n define by $\varrho(t_n) = 0$ is a “bolt” of twist $k = n + 1, n + 2, \dots$, [4].

3.2 Compact metrics

Let us suppose $c < 0, \lambda > |c|$ and $n \geq 2$. From (3.1) we can deduce there does exist a finite interval $[t_n^{(1)}, t_n^{(2)}]$ on which $\varrho(t)$ is positive. The constraints for $t = t_n^{(1)}$ and $t = t_n^{(2)}$ to be bolts of twist k are [5]

$$\begin{cases} \varrho(t_n^{(1)}) = 0, & \varrho'(t_n^{(1)}) = k, \\ \varrho(t_n^{(2)}) = 0, & \varrho'(t_n^{(2)}) = -k. \end{cases} \quad k = 1, 2, 3, \dots$$

which reduce, using the differential equation, $\varrho' + \left(c + \frac{n-1}{t} \right) \varrho = n - \lambda t$, to the transcendental equation

$$\int_{-k}^{+k} e^{-\xi u} u (n + u)^{n-1} du = 0, \quad \xi = \frac{|c|}{t}, \quad k = 1, 2, \dots, n - 1.$$

And this equation has a unique solution $\xi \in]0, 1[$.

4 Bosonic σ -models at two loops

For a given bosonic σ -model with classical action

$$\frac{1}{g^2} \int d^2x \frac{1}{2} g_{ij}(\varphi) \partial_\mu \varphi^i \partial_\mu \varphi^j$$

the one and two-loop on-shell divergences follow from Friedan [3] and are given by

$$\left(\frac{\hbar}{4\pi}\right) \frac{1}{\epsilon} \int d^2x Ric_{ij} \partial_\mu \varphi^i \partial_\mu \varphi^j + \left(\frac{\hbar}{4\pi}\right)^2 \frac{g^2}{\epsilon} \int d^2x \frac{1}{2} R_{is,tu} R_j^{s,tu} \partial_\mu \varphi^i \partial_\mu \varphi^j, \quad \epsilon = d - 2.$$

Two-loop renormalizability requires

$$X_{ij} = R_{is,tu} R_j^{s,tu} = 0 = D_i w_j + D_j w_i \quad \Rightarrow \quad w_i = c_2 \partial_i (sA)$$

and

$$X_{i\bar{j}} = R_{i\bar{k},l\bar{n}} R_{\bar{j}}^{\bar{k},l\bar{n}} = \lambda_2 g_{i\bar{j}} + D_i w_{\bar{j}} + D_{\bar{j}} w_i = \partial_{i\bar{j}}^2 (\lambda_2 K + (c_2 + \bar{c}_2) sA).$$

For the divergence $X_{i\bar{j}}$ to be absorbable it is therefore necessary that $X_{i\bar{j}}$ be Kähler. In the class of metrics considered in this article, the large isometry group enables one to compute the curvature tensor and

$$X_{i\bar{j}} = \mu(s) \delta_{ij} + \nu(s) \bar{z}_i z_j.$$

The Kähler condition requires therefore $\nu = \mu'$ which is integrated to $K(s) = a \ln(s + b)$. Among all kählerian metrics of dimension $n \geq 2$, with an isometry group $U(n)$ acting linearly on the holomorphic coordinates, only CP^n is one-loop and two-loop renormalizable.

References

- [1] G. Bonneau and F. Delduc, *Int. J. Mod. Phys. A*, **1** (1986) 997.
- [2] T. Eguchi, P. B. Gilkey and A. J. Hanson, *Phys. Report.* **66** (1980) 214.
- [3] D. H. Friedan, *Ann. Phys.* **163** (1985) 318.
- [4] G.W. Gibbons and S. W. Hawking, *Commun. Math. Phys.* **66** (1979) 291.
- [5] N. Koiso, in "Recent topics in differential and analytic geometry", *Advanced studies in pure Math.* **18-I** (1990) 327.