

Zeitschrift: Mitteilungen aus dem Gebiete der Lebensmitteluntersuchung und Hygiene = Travaux de chimie alimentaire et d'hygiène
Herausgeber: Bundesamt für Gesundheit
Band: 74 (1983)
Heft: 4

Artikel: Chlorobenzene residues in market milk and meat
Autor: Jan, J.
DOI: <https://doi.org/10.5169/seals-983017>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 15.03.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Chlorobenzene Residues in Market Milk and Meat

J. Jan

Public Health Institute of Slovenia, Ljubljana

Introduction

Chlorobenzenes (CB) represent a group of chlorinated hydrocarbons and are ubiquitous environmental pollutants (1, 2). Contamination of milk and meat with CB could appear owing to their high production, wide use, lipophilic character and slow degradation in the environment. CB are in minor use as pesticides, while more are used as solvents, heat transfer fluids, flame retardants and in the electro-industry. In the environment they could also be derived from pesticides (e. g. HCH) by metabolic processes (3), and probably during incineration of organic matter and chloro compounds (4). They have been found in water, fish, seeds and man (2, 5, 6, 7).

This investigation reports the data on CB (di-, tri-, tetra-, penta-, and hexa-) residues in milk and meat; of these only information on hexa-CB (8) in milk and meat, and on 1,4-di-CB in pork meat (9) is available. It is important to collect data on CB levels, since they have comparable toxicity and properties to organochloro pesticides and polychlorinated biphenyls (10, 11). The technique of destructive clean-up as described for organochloro compounds (12), was also applied for CB. The presence of CB in milk and meat was quantified by high resolution glass capillary GL chromatography.

Experimental

Samples

The 9 raw milk samples were received as an average daily sample from the city milk distribution in Ljubljana (Yugoslavia) in September—October 1981. The 3 fresh beef meat samples (*M. long. dorsi*) were purchased on the market.

Reagents

1,2-dichlorobenzene (1,2-DCB); 1,3-dichlorobenzene (1,3-DCB); 1,4-dichlorobenzene (1,4-DCB); 1,2,3-trichlorobenzene (1,2,3-TCB); 1,2,4-trichlorobenzene (1,2,4-TCB); 1,3,5-trichlorobenzene (1,3,5-TCB); 1,2,3,4-tetrachlorobenzene (1,2,3,4-TeCB); 1,2,3,5-tetrachlorobenzene (1,2,3,5-TeCB); 1,2,4,5-tetrachloroben-

zene (1,2,4,5-TeCB); pentachlorobenzene (PeCB); hexachlorobenzene (HCB) were supplied by Fluka.

To lower the blank value, glassware was rigorously cleaned with methanol and heated at 300 °C for 1 h. All solvents were glass distilled. Florisil, 60–100 mesh, (Fluka) was Soxhlet extracted with benzene for 3 h before activation for 3 h at 550 °C and maintained at 130 °C prior to use. The reagent blank was measured and corrected for in sample analyses. Recoveries of CB added to beef meat were over 80 %.

Instrumentation

A Varian model 3700 gas chromatograph; electron capture detector (^{63}Ni); glass capillary columns (a) i. d. 0.32 mm, length 20 m coated with SE-30 and (b) i. d. 0.30 mm, length 20 m coated with OV-1. Injector temperature 210 °C, detector temperature 240 °C. Temperature programme: initial column oven 60 °C held for 40 s (during the purge activation time) then the temperature raised to 65 °C in 20 s; 65–170 °C temperature programme 5 °C/min 170 °C isothermal; splitless injection, purge activation time 40 s. Sample 1–3 μl . Solvent n-hexane. Nitrogen flow: column 0.9, injector purge 120, and make-up 25 ml/min, respectively.

Procedures

Meat

To 20 g of finely chopped fresh beef meat and 15 ml of n-hexane were added 20 ml of 1:1 (v/v) H_2SO_4 -water in an erlenmeyer flask and shaken. After half an hour 30 ml of conc. sulfuric acid was added and digested overnight at room temperature. The reaction mixture was put in a separatory funnel, the acid phase discarded and the hexane extract cleaned-up with 20 ml conc. sulfuric acid. The hexane extract was passed through a florisil column 10 cm in height with an i. d. of 1 cm having 0.5 cm of anhydrous Na_2SO_4 on the top, and eluted with 25 ml of 6% diethyl ether in hexane. The eluate was concentrated to 2 ml at 50 °C in a flow of nitrogen, and chromatographed. The chromatogram is shown in figure 1.

Milk

20 g of milk and 10 ml of n-hexane were cooled below 0 °C and 20 ml of conc. sulfuric acid was slowly added with stirring. After 1 h when the mixture reached room temperature, a fresh portion of 10 ml of conc. sulfuric acid was added. After standing overnight, the hexane layer was separated and cleaned with 10 ml of conc. sulfuric acid and through the florisil column, and the eluate evaporated and analysed by GL chromatography.

An additional clean-up could be done with hydrolysis by a modified procedure of *Young and Burke* (13): to the hexane eluate evaporated to 1 ml, 1 ml of 2% ethanolic KOH was added, this was then sealed in a glass vial and hydrolysed for 1 h at 95 °C. After cooling, water was added to the reaction mixture, extracted with hexane and cleaned-up through conc. sulfuric acid and florisil as above, and




Fig. 1. Chromatogram of the beef meat extract on a SE-30 glass capillary column (for abbreviations, see text)

chromatographed. For quantitative assessment, the increased trichlorobenzene levels produced from hydrolysis of HCH isomers should be taken into consideration.

Results and discussion

In table 1 residue levels of CB and HCH (hexachlorocyclohexane) pesticides in milk and meat are presented. The level in both samples is similar. The total CB (dichloro to hexachloro) in milk falls in the range of total organochloro pesticides (sum of HCH, DDT and cyclodiene groups) as well as polychlorinated biphenyls; the corresponding values are approximately 0.4 (from table 1), 0.1 (8, 14), and 0.3 (from table 2) mg/kg recalculated and expressed on a fat basis, respectively. It is possible to speculate that the contamination of milk and meat with CB could originate through water (5, 20) and cattle feedstuffs prepared on a fish (7) and seed (6) basis, since CB have been found in these sources.

At present there are no available legal tolerances for CB in food except for HCB. WHO (10) and FDA (11) have proposed some tentative acceptable daily intakes (ADI) and recommended residue limits for organochloro contaminants. In table 2 a comparison of these limits with the average content in milk and the toxicity of CB is presented.

Table 1. Mean levels of chlorobenzenes in cow's milk and beef meat (ng/g «as in» basis)

| Compound Sample | DCB | | | TCB | | | TeCB | | PeCB | HCH | Total HCH ^a |
|-------------------------|-----------------|------|------|--------|--------|--------|--------------------------------|----------|------|------|------------------------|
| | 1,3- | 1,4- | 1,2- | 1,3,5- | 1,2,4- | 1,2,3- | 1,2,3,5- and/or 1,2,4,5- | 1,2,3,4- | | | |
| Cow's milk ^b | nd ^d | 5.3 | 2.6 | 0.9 | 0.7 | 1.5 | 1.10 | 0.20 | 0.05 | 0.35 | 1.31 |
| Beef meat ^c | nd | 5.0 | 1.0 | 1.2 | 1.0 | 1.8 | 1.00 | 0.02 | 0.05 | 1.10 | 2.40 |

The fat content of the milk was 3.2% and of the meat 2.1%. The GLC analytical technique was not sensitive enough (electron capture detector sensitivity of lower chlorobenzenes is low) to determine DCB and TCB to more than two digits

^a the sum of alpha-, beta- and gamma-hexachlorocyclohexane isomers

^b arithmetic means of 9 samples analysed

^c arithmetic means of 3 samples analysed

^d not detected

Table 2. Toxicological data, tolerances and residues of some organochloro contaminants in milk

| Compound | LD ₅₀ (mg/kg body weight, rat, orally) (15) | ADI (mg/kg body weight) (10) | Residue limit (10) and common content ^a in milk (mg/kg on fat basis) | |
|---------------------------|--|------------------------------|---|-------------------------|
| | | | | |
| Gamma-HCH (lindane) | 125 | 0.0125 | 0.1 | 0.02–0.03 (8, 14) |
| HCB | 3 500 | 0.0006 | 0.3 | 0.01–0.03 (8, 14) |
| 1,4-DCB | 500 | 0.014–0.0014 | no data | 0.16 ^b |
| Polychlorinated biphenyls | 4 000 (16) | 0.02 (11) | 1.5 (17) | 0.2–0.4 (8, 14, 18, 19) |

^a in some European countries

^b calculated from table 1

Results of this contribution illustrate that the CB levels in milk and meat are not so high as to cause concern; however, as the toxic effects of CB are comparable to those of the other organochloro contaminants, it would be advisable in food contamination monitoring to include, besides HCB, CB isomers with less than six chloro atoms on the benzene nucleus.

Acknowledgement

The Slovenian Science Foundation supported this work financially.

Summary

Chlorobenzene levels were determined in cow's milk and beef meat samples. Concentrations determined by capillary GL chromatography for some chlorobenzene isomers were from 0.05 to 5 ng/g, original weight basis. The total amount of chlorobenzenes (sum of dichloro to hexachloro), of approximately 10 ng/g, original basis, is found to be in the range of other organochloro contaminants — e. g. organochloro pesticides and polychlorinated biphenyls.

Zusammenfassung

In Kuhmilch- und Rindfleischproben wurden mittels Kapillargaschromatographie die Konzentrationen einzelner Chlorbenzole (Dichlor- bis Hexachlorbenzol) bestimmt. Sie lagen im Bereich von 0,05 bis 5 ng/g (Frischgewicht). Der Totalgehalt an Chlorbenzolen (Summe) betrug ungefähr 10 ng/g und war somit mit den Gehalten anderer chlorierter Kohlenwasserstoffe (Organochlorpestizide, polychlorierte Biphenyle) vergleichbar.

Résumé

Les résidus de chlorobenzènes ont été déterminés dans des échantillons de lait de vache et dans la viande de boeuf. Les concentrations, déterminées par chromatographie gaz liquide capillaire pour quelques isomères de chlorobenzène, varient de 0,05 à 5 ng/g de poids original. La quantité totale des chlorobenzènes (la somme des dichloro- à hexachlorobenzènes) est d'environ 10 ng/g du poids original, elle se trouve dans le domaine des autres contaminants organochlorés — par exemple des pesticides organo-chlorés et des biphenyles polychlorés.

Literature

1. *Fishbein, L.*: Potential halogenated industrial carcinogenic and mutagenic chemicals. IV. Halogenated aryl derivatives. *Sci. Total Environ.* **11**, 259–278 (1979).
2. *Morita, M.*: Chlorinated benzenes in the environment. *Ecotoxicol. Environ. Safety* **1**, 1–6 (1977).
3. *Engst, R., Macholz, R. M. and Kujawa, M.*: Recent state of lindane metabolism. Part II. *Residue Rev.* **72**, 71–95 (1979).
4. *Choudhry, G. G. and Hutzinger, O.*: Mechanistic aspects of the thermal formation of halogenated organic compounds including polychlorinated dibenzo-p-dioxins. Part V: Hypotheses and summary. *Toxicol. Environ. Chem.* **5**, 295–309 (1982).
5. *Oliver, B. G. and Nicol, K. D.*: Chlorobenzenes in sediments, water, and selected fish from lakes Superior, Huron, Erie, and Ontario. *Environ. Sci. Technol.* **16**, 532–536 (1982).
6. *Jan, J.*: Chlorinated benzene residues in some seeds. *Chemosphere* **9**, 165–167 (1980).

7. *Jan, J. and Malneršič, S.*: Chlorinated benzene residues in fish in Slovenia (Yugoslavia). *Bull. Environ. Contam. Toxicol.* **24**, 824–827 (1980).
8. *Heeschen, W.*: Rückstände von Arzneimitteln, Pestiziden und Umweltchemikalien in der Milch – ein Überblick zur gegenwärtigen Situation. *Tierärztl. Umschau* **37**, 538–545 (1982).
9. *Watson, A. and Patterson, R. L. S.*: Tainting of pork meat by 1,4-dichlorobenzene. *J. Sci. Food Agric.* **33**, 103–105 (1982).
10. *FAO/WHO*: Pesticide residues in food. WHO technical Report Series No. 574. WHO, Geneva 1975.
11. *US Department of Health, Education and Welfare, Food and Drug Administration*: Polychlorinated biphenyls (PCB's). *Fed. Reg.* **38**, 18096–18103 (1973).
12. *Veierov, D. and Aharonson, N.*: Economic method for analysis of fluid milk for organochlorine residues at the 10 ppb level. *J. Assoc. Off. Anal. Chem.* **63**, 532–535 (1980).
13. *Young, S. J. V. and Burke, J. A.*: Micro-scale alkali treatment for use in pesticide residue confirmation and sample cleanup. *Bull. Environ. Contam. Toxicol.* **7**, 160–167 (1972).
14. *Venant, A. et Richou-Bac, L.*: Contamination des produits laitiers français par les résidus de composés organo-chlorés. *Lait* **61**, 619–633 (1981).
15. *Christensen, H. E., Fairchild, E. J., Carroll, B. S. and Lewis, R. J.* (Editors): NIOSH registry of toxic effects of chemical substances. HEW Publ., Washington D.C. 1976.
16. *WHO*: Polychlorinated biphenyls and terphenyls. *Environ. Health Crit. Vol. 2*, WHO, Geneva 1976.
17. *US Department of Health, Education and Welfare, Food and Drug Administration*: Polychlorinated biphenyls (PCB's); reduction of tolerances. *Fed. Reg.* **44**, 38330 (1979).
18. *Van Renterghem, R. and Devlaminck, L.*: Polychlorinated biphenyl compounds in milk and dairy products. *Z. Lebensm.-Unters. Forsch.* **170**, 346–348 (1980).
19. *Sümmerrmann, W., Rohleder, H. und Korte, F.*: Polychlorierte Biphenyle (PCB) in Lebensmitteln. *Z. Lebensm.-Unters. Forsch.* **166**, 137–144 (1978).
20. *Grob, K. and Grob, G.*: Organic substances in potable water and its precursor, Part II, Applications in the area of Zürich. *J. Chromatogr.* **90**, 303–313 (1974).

Dr. J. Jan
 Public Health Institute of Slovenia
 Trubarjeva 2
 61000 Ljubljana
 Yugoslavia