

**Zeitschrift:** Mitteilungen der Schweizerischen Entomologischen Gesellschaft =  
Bulletin de la Société Entomologique Suisse = Journal of the Swiss  
Entomological Society

**Herausgeber:** Schweizerische Entomologische Gesellschaft

**Band:** 73 (2000)

**Heft:** 3-4

**Artikel:** Response of Orthoptera assemblage composition to land-use in the  
southern Alps of Italy

**Autor:** Guido, Marco / Chemini, Claudio

**DOI:** <https://doi.org/10.5169/seals-402790>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 16.03.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Response of Orthoptera assemblage composition to land-use in the southern Alps of Italy

MARCO GUIDO<sup>1</sup> & CLAUDIO CHEMINI<sup>1</sup>

The effects of land-use systems and thermal heterogeneity on the composition of Orthoptera assemblages in secondary grasslands were investigated on a site in the Southern Alps of Italy. Orthoptera species were sampled on seven plots with different management regimes, and different aspects. Species were classified on the basis of their distribution, in an effort to identify groups of species with particular responses to human disturbance. Stepwise multiple-regression analysis revealed that heterogeneity in thermal conditions as expressed by irradiation played a key role in determining species distribution across the study site. A group of thermophilous species related to South-facing plots was distinguished from more generalist species that also colonize East-facing plots. It was also possible to distinguish species that appeared to receive benefits from abandonment from species occurring mainly on managed (mown) meadows. Differences in group densities among plots reflected differences in ground level temperature and vegetation structure.

Keywords: Orthoptera, land-use change, assemblage composition, temperature, vegetation structure, Italy, Alps.

### INTRODUCTION

In the last decades, changes in agricultural production methods have led to substantial changes in land-use systems (TURNER II & MEYER, 1994), including intensification in productive landscapes and abandonment of less productive mountain soils. Intensification and abandonment have led to the disappearance of many traditionally managed grassland areas throughout the entire Alpine region (SURBER *et al.*, 1973; PENZ, 1978; BÄTZING, 1990).

Land-use systems affect the vegetation structure and diversity (TAPPEINER & CERNUSCA, 1993; CERNUSCA *et al.*, 1992), and so the fauna (MORRIS, 1978; SOUTHWOOD *et al.*, 1979; ERHARDT & THOMAS, 1991; review in TSCHARNTKE & GREILER, 1995). In the mountains, an important role in determining the local occurrence of species is also played by climate heterogeneity resulting from topographical complexity.

Orthoptera are important primary consumers of alpine grasslands (BLUMER & DIEMER, 1996) and are effective bioindicators due to their differentiated reaction to habitat disturbances (KÖHLER, 1990). The choice of habitat by species depends mainly on microclimate conditions (VAN WINGERDEN *et al.*, 1991a; COXWELL & BOCK, 1995; review in INGRISCH & KÖHLER, 1998) and vegetation patterns (KEMP *et al.*, 1990; QUINN *et al.*, 1991; FIELDING & BRUSVEN, 1995). Therefore, land-use practices that imply a change in the structure of vegetation and microclimate must

<sup>1</sup> Centro di Ecologia Alpina, 38040 Viote del Monte Bondone, Trento, Italy  
guido@cealp.it, chemini@cealp.it

significantly affect Orthoptera assemblages (VAN WINGERDEN *et al.*, 1991b; FRICKE & VON NORDHEIM, 1992; THORENS, 1993; ANTOGNOLI & ZETTEL, 1996; review in INGRISCH & KÖHLER, 1998).

This study deals with the effects of climate heterogeneity and land-use systems on the composition of Orthoptera assemblages from secondary grasslands in the Southern Alps. Species distribution was analyzed in relation to selected environmental variables in order to characterize the different groups and to distinguish the impact of different land-use systems from the effect of thermal heterogeneity resulting from topographical aspect. This study also aims to integrate the results of a paper that assessed the effects of land-use changes on Orthoptera abundance and diversity (GUIDO *et al.*, 1998), and it is part of a larger research project (EC project ECOMONT) carried out along a N-S transect from the Austrian to the Italian Alps.

MATERIALS AND METHODS

*Study site*

The study site was located on the Viote plateau (46°01'20"N, 11°02'30"E) on Mount Bondone, at an altitude of 1550 m. Mount Bondone range is part of the Southern Alps and is located on the west side of the River Adige near the city of Trento.




Fig. 1 - Location of the plot study sites at Monte Bondone. The location of sampled areas within each plot is also given by rectangles including the name of the plot. Plots characterized by different land-use are indicated with different scales of grey. Abandoned meadows: dark grey; mown meadows: mid grey; pasture: light grey.

The mean annual temperature is 5.5 °C, with mean monthly temperatures ranging from -2.7 °C in January to 14.4 °C in July. Precipitation is abundant through the year (mean annual rainfall 1189 mm), with two peaks in June and October, and a minimum in January.

At present, the most common land-use system is extensive mowing, as meadows are cut once a year and lightly fertilized (30 kg N ha<sup>-1</sup>). The vegetation of these meadows is dominated by golden oats field (*Geranio-Trisetum flavescens*) and nard field (*Sieversio-Nardetum strictae*) (TASSER *et al.*, 1999). The beech forest represents the climax vegetation. Detailed information about the study site is given in CESCATTI *et al.* (1999).

### Sampling methods

Sampling of adult Orthoptera was carried out on seven plots from 1 July to 15 September 1997 (Tab. 1 and Fig. 1). Plots were selected in relation to aspect (southern or eastern) and types of land-use system. Pasture plots were available only on E-facing slopes. Within each plot, abundance of adult Orthoptera was assessed by sweep netting. Four replicate patches of 5 m<sup>2</sup> per plot were sampled. A modified sweep net with a plastic bag attached was used. Sweeping was repeated eight times on the same replicate patch, to make sure that all insects were caught. Each sweeping consisted of 10 strokes per m<sup>2</sup>. This number of strokes was found to allow of a complete coverage of the patch surface in all the different investigated types of vegetation. After each sweeping, the plastic bag was changed. Therefore, abundance

Tab. 1 - Description of the sampling plots; Al = altitude (m), S = slope (°), As = Aspect, I = Irradiation (nh). Vegetation types according to TASSER *et al.* (1999).

Plot	Description	Al	S	As	Vegetation type	Management system	I
<b>Aba1</b>	Abandoned grassland	1530	10	E	Sieversio-Nardetum strictae with <i>Calluna vulgaris</i>	none since 30 years	1991
<b>Mow1</b>	Hay meadow	1550	10	E	Sieversio-Nardetum strictae	mown once a year and fertilised (30kgN ha <sup>-1</sup> )	1991
<b>Mow2</b>	Hay meadow	1530	10	E	Geranio-Trisetum flavescens	mown once a year and fertilised (30kgN ha <sup>-1</sup> )	1967
<b>Past</b>	Pasture	1560	15	E	Sieversio-Nardetum strictae	grazing by cattle and horses (1.5 animals ha <sup>-1</sup> )	1926
<b>Aba2</b>	Abandoned grassland	1565	15	S	Geranio-Trisetum flavescens	none since 10 years	2339
<b>Mow3</b>	Hay meadow	1540	15	S	Sieversio-Nardetum strictae trifolietosum	mown once a year and fertilised (30kgN ha <sup>-1</sup> )	2337
<b>Mow4</b>	Hay meadow	1570	15	S	Arrhenatheretum elatioris	mown once a year and fertilised (30kgN ha <sup>-1</sup> )	2339

data were calculated per 5 m<sup>2</sup>. Sampling was only carried out under fair weather conditions. Quantitative samplings were repeated every 15 days (i.e. six samples). On managed meadows, mowing occurred between the 2nd and 3rd sampling.

Heterogeneity in thermal conditions among plots resulting from topographical aspect was expressed by irradiation. Irradiation was estimated for each plot according to BARTORELLI (1965), who proposed an algorithm for the calculation of the amount of solar radiation hitting the ground as a function of latitude, slope and aspect. Irradiation is measured in normal hours (nh), which give the lap of time (measured in hours) during which the sun should stay directly perpendicular to a place in order to give the same amount of radiation that it gives under normal conditions to the same place during a year.

Above-ground phytomass was estimated by clipping all growth of the current year on four patches per plot. The size of each harvested patch was 0.25 m<sup>2</sup>. All clipped material was divided into grasses, herbs (including dwarf shrubs), and dead standing plant matter (necromass). Samples were obtained every 15 days during the vegetative season from June to September (i.e. eight samples; TELLINI, pers. comm.).

Ground level temperatures were measured by thermocouples connected to a portable data acquisition system that provided temperature data for each hour (TAPPEINER *et al.*, 1999). We considered the mean daily temperature and maximum daily temperature of eight periods of seven days, with Orthoptera and phytomass sampling date as the central day of each period.

#### *Data analysis*

An analysis of variance (ANOVA) was used to test differences between plots and periods in measured environmental variables. As homogeneity of variances was not achieved even by data transformation, non-parametric Kruskal-Wallis ANOVA by ranks was applied to the data, and the effects of plot, aspect (S vs. E), and land-use were tested separately. To assess the effects of aspect on temperature and vegetation biomass, the pasture plot was not considered due to the absence of a S-facing replicate. Effects of land-use systems were analyzed considering E-facing and S-facing plots separately. To test differences between means, non-parametric multiple comparisons using rank sums from the Kruskal-Wallis test (ZAR, 1984) were performed.

Species occurrence in the different plots was analyzed by cluster analysis (CA) in order to obtain groups of species. For CA, abundance data were treated using 1-Pearson r as the distance index, and weighted pair-group method using arithmetic averages (WPGMA) as the clustering strategy. The density of each group of species was calculated by summing the densities of individual member species. To test the differences in abundance of CA groups, data were then submitted to a Kruskal-Wallis ANOVA followed by non-parametric multiple comparisons, using plot, aspect (not considering the pasture plot), and land-use (considering S- and E-facing plots separately) as independent variables. Densities of groups were also correlated to the environmental variables by forward stepwise regression analysis. Although sweep netting is not a particularly adequate method for catching Tetrigid species, abundance data of *Tetratetrix bipunctata kraussi* were included into the statistical analyses, since the distribution among plots of this species as obtained by sweep netting reflected the distribution as estimated in the field by qualitative observations. All statistical analyses were performed using the software package STATSOFT Statistica, version 5.




Fig. 2 - Distribution of mean (A) and maximum (B) ground level temperature for plots with different land-uses. The mean (+ SD) temperature values corresponding to the different land-use systems separated by aspect are given; Aba = abandoned meadow, Mow = mown meadow, Past = pasture. Results of Kruskal-Wallis test performed separately for E-facing and S-facing plots using land-use as independent variable are also given. Different letters imply significant differences in pairwise comparison among land-uses under similar aspect as revealed by non-parametric Tukey-type test.

## RESULTS

### *Environmental variables*

Irradiation (Tab. 1) and ground level temperature (both mean and maximum) were higher on the S-facing plots ( $H_{1, 192} = 59.13$ ,  $p < 0.01$ , and  $H_{1, 192} = 31.49$ ,  $p < 0.01$ , respectively). The mean and maximum temperatures (under similar irradiation) were significantly higher on managed plots (Fig. 2A). In particular, temperature maxima were clearly greater on mown meadows (Fig. 2B).

Above-ground phytomass was highly affected by thermal heterogeneity and by land-use (Fig. 3). Necromass was significantly higher on E-facing plots ( $H_{1, 190} = 4.43$ ,  $p < 0.05$ ), where necromass was higher on the pasture. The first result can be explained as due to the relatively lower irradiation and temperature, that led to a slower decomposition process of litter on E-facing plots. The second one can be explained by the fact that the investigated pasture plot was covered by a high percentage of perennial grasses, which accumulated dead matter into their tufts. On S-facing plots, necromass was significantly higher on abandoned plots (Fig. 3A). The grass biomass was higher on S-facing plots ( $H_{1, 192} = 14.30$ ,  $p < 0.01$ ). No significant differences in grass biomass were revealed between managed and abandoned plots (Fig. 3B). Herb biomass had no differences due to aspect ( $H_{1, 185} = 1.93$ ,  $p > 0.05$ ); under similar irradiation, herb biomass was higher on abandoned plots (Fig. 3C).

### *Orthoptera assemblages*

A total of 3520 individuals belonging to 15 species were collected on the seven study plots (Tab. 2). Cluster analysis (CA) separated the species by their local occurrence. A first separation into two clusters can be observed at a distance of about 1 (Fig. 4A). In the cluster (a) are species which were exclusively found (*Leptophyes*




Fig. 3 - Distribution of necromass (A), grass biomass (B), and herb/dwarf shrub biomass (C) for different land-uses. The mean (+ SD) dry weight values per m<sup>2</sup> corresponding to different land-use systems separated by aspect are given; Aba = abandoned meadow, Mow = mown meadow, Past = pasture. Results of Kruskal-Wallis tests performed separately for E-facing and S-facing plots using land-use as the independent variable are also given. Different letters imply significant differences in pairwise comparison among land-uses under similar aspect as revealed by non-parametric Tukey-type test.

*bosci*, *Gomphocerus rufus*, *Chorthippus dorsatus*) or with higher densities (*Tetigonia caudata*, *Arcyptera fusca*, *Euthystira brachyptera*, *Omocestus viridulus*, *Stenobothrus lineatus*, *Stauroderus scalaris*, *Glyptobothrus biguttulus*) on the S-facing areas (Aba2, Mow3, Mow4) (Tab. 2). This cluster was significantly and positively correlated to irradiation, herb biomass and mean temperature (Tab. 3). In the cluster (b), there are 5 species with a significant negative statistical relationship with

irradiation and necromass (Tab. 3). These two clusters seem to be separated on the basis of the distribution of species in relation to thermal heterogeneity, and thus on the thermal requirements of species. In particular, Orthoptera from the group (a) are linked to warmer sites, while the species grouped in cluster (b) do not show precise thermal requirements, being found with similar densities on both E- and S-facing plots (Tab. 4).

A further classification can be operated at a distance level of about 0.8 (Fig. 4A). Each of the two main clusters (a) and (b) splits into two branches.

Cluster (a) (S preferring species) splits into a first group (1), which includes species that reached significantly higher densities on S-facing mown meadows (Figs 4B, 5). This cluster groups two species which showed a clear preference for S-facing mown meadows (*G. biguttulus* and *S. lineatus*), and two other species which occurred with relatively higher densities on such plots, although not showing such a strong preference for a particular land-use (*S. scalaris* and *C. dorsatus*) (Tab. 2). The relatively less clear preference shown by *S. scalaris* might be due to its high mobility by flying across the plots. The group was positively related to mean temperature and irradiation, and negatively related to necromass (Tab. 3). The other group (2) is characterized by species that preferred abandoned meadows (Fig. 5). This cluster includes species with a clear preference for abandoned habitats (*E. brachyptera*, *L. bosci*, *G. rufus*) as well as species that occurred only with a

Tab. 2 - Total captures per plot (20 m<sup>2</sup>) of the species of Orthoptera found in the surveyed areas. Species are named according to FAILLA *et al.* (1995).

Species	Plot						
	Aba1	Mow1	Mow2	Past	Aba2	Mow3	Mow4
<b>Fam. Tettigoniidae</b>							
<i>Poecilimon ornatus</i> (Schmidt)	4	1	4	2	2	1	1
<i>Leptophyes bosci</i> Brunner					2		
<i>Tettigonia caudata</i> (Charpentier)			2		3	1	1
<i>Decticus verrucivorus</i> (Linnaeus)	34	42	103	2	27	22	25
<i>Pholidoptera griseoptera</i> (De Geer)	22	3	1		4		1
<b>Fam. Tetrigidae</b>							
<i>Tetratetrix bipunctata kraussi</i> (Saulcy)	3	4			2		4
<b>Fam. Acrididae</b>							
<i>Arcyptera fusca</i> (Pallas)	21	5	6		49	11	48
<i>Euthystira brachyptera</i> (Ocskay)	197	49	18	46	309	103	82
<i>Omocestus viridulus</i> (Linnaeus)	20	33	14	30	137	56	138
<i>Stenobothrus lineatus</i> (Panzer)		2				9	4
<i>Gomphocerus rufus</i> (Linnaeus)					4		
<i>Stauroderus scalaris</i> (Fischer Waldheim)	43	27	26	5	56	83	72
<i>Chorthippus d. dorsatus</i> (Zetterstedt)					3	7	3
<i>Chorthippus p. parallelus</i> (Zetterstedt)	248	195	220	139	131	196	286
<i>Glyptobothrus biguttulus</i> (Linnaeus)		1			4	35	26



Tab. 3 - Results of forward stepwise regression analysis performed on groups obtained by classification of Orthoptera species. Environmental variables with significant effect on each group distribution and their contribution to change in R-square are given.

Group	Resume of Regression	Independent variables entered ( $p < 0.05$ )	Beta	R-square change
Cluster a	$R^2 = 0.66$ , $F(4,163) = 77.97$ , $p = 0.000$	<b>Irradiation</b>	0.40	0.38
		<b>Herb &amp; shrub biomass</b>	0.35	0.13
		<b>Mean temperature</b>	0.81	0.09
		<b>Maximum temperature</b>	-0.56	0.06
				S = 0.66
Cluster 1	$R^2 = 0.38$ , $F(3,164) = 32.96$ , $p = 0.000$	<b>Mean temperature</b>	0.35	0.30
		<b>Irradiation</b>	0.24	0.05
		<b>Necromass</b>	-0.15	0.02
				S = 0.38
Cluster 2	$R^2 = 0.60$ , $F(4,163) = 61.60$ , $p = 0.000$	<b>Irradiation</b>	0.38	0.28
		<b>Herb &amp; shrub biomass</b>	0.40	0.21
		<b>Mean temperature</b>	0.73	0.05
		<b>Maximum temperature</b>	-0.58	0.06
				S = 0.60
Cluster b	$R^2 = 0.31$ , $F(4,163) = 18.04$ , $p = 0.000$	<b>Necromass</b>	-0.45	0.07
		<b>Irradiation</b>	-0.57	0.03
		<b>Mean temperature</b>	0.59	0.09
		<b>Herb &amp; shrub biomass</b>	0.41	0.11
				S = 0.31
Cluster 3	$R^2 = 0.28$ , $F(4,163) = 16.05$ , $p = 0.000$	<b>Mean temperature</b>	0.60	0.10
		<b>Irradiation</b>	-0.48	0.03
		<b>Necromass</b>	-0.39	0.05
		<b>Herb &amp; shrub biomass</b>	0.38	0.10
				S = 0.28
Cluster 4	$R^2 = 0.16$ , $F(4,163) = 7.57$ , $p < 0.001$	<b>Irradiation</b>	-0.50	0.04
		<b>Necromass</b>	-0.36	0.06
		<b>Herb &amp; shrub biomass</b>	0.26	0.02
		<b>Mean temperature</b>	0.26	0.03
				S = 0.16

slight preference for the S-facing abandoned meadow (*A. fusca*, *O. viridulus*, *T. caudata*). In particular, *A. fusca* and *O. viridulus* reached similar high density values on both Aba 2 and Mow 4 (Tab. 2), although the results of classification seemed to be mostly affected by low density values recorded on Mow 3. In addition, some of the sampled patches of Mow 4 were close to the abandoned area (less than 10 m, Fig. 1), thus a migration process from the nearby abandoned meadow might have been involved in determining the high densities of these two species recorded on Mow 4. Stepwise regression analysis showed a significant positive relationship between this group and irradiation, herb biomass, and mean temperature. Maximum

Tab. 4 - Mean densities and SD of the groups of orthopteran species obtained by classification among areas at a distance of 1. Mean densities are referred to 5 m<sup>2</sup>. Results of Kruskal-Wallis tests using plot as main effect are also given. Different letters imply significant differences in pairwise comparisons among plots as revealed by non-parametric Tukey-type tests.

Cluster	Plot						
	Aba1	Mow1	Mow2	Past	Aba2	Mow3	Mow4
<b>a</b>	Kruskall-Wallis test: H (6,168) = 103.10, p < 0.01						
Mean	11.71	5.00	2.79	3.37	23.58	12.71	15.58
S.D.	5.77	4.90	2.65	3.24	9.89	8.17	8.38
	b	c	c	c	a	ab	ab
<b>b</b>	Kruskall-Wallis test: H (6,168) = 30.55, p < 0.01						
Mean	13.08	10.17	13.67	5.96	7.04	9.12	13.37
S.D.	6.41	6.37	9.51	3.86	4.76	6.37	7.41
	a	ab	a	b	b	ab	a

temperature, higher on mown meadows, was negatively related to this group (Tab. 3).

Cluster (b) is also divided into two groups. These groups are composed of few species (2 and 3 respectively) and the autoecology and abundance of individual species play a substantial role. The first cluster (3) grouped two species, *Chorthippus parallelus* and *T. bipunctata*, and the local distribution of this group reflected the distribution of the dominant and eurytopic *C. parallelus*. It occurred with significantly different densities among plots, but without any significant relationship with thermal heterogeneity and land-use system (Figs 4B, 5; Tab. 3). The second branch (4) reached significantly higher densities on E-facing plots and was negatively correlated to irradiation (Fig. 4B; Tab. 3). The local occurrence of the group results from the distribution pattern of *Decticus verrucivorus*, and that of *Pholidoptera griseoptera* and *Poecilimon ornatus*. The first species reached a higher density on mown meadows, whereas *P. griseoptera* was more abundant on abandoned meadows and *P. ornatus* occurred with similar density on both managed and abandoned meadows (Tab. 2).

#### DISCUSSION

Both thermal heterogeneity and land-use systems affect the species composition and especially the local occurrence of Orthoptera on Mount Bondone.

The role of solar radiation and general thermal conditions on Orthoptera was underscored by SAMWAYS (1990) and COXWELL & BOCK (1995), and a relatively





Fig. 4 - Classification of orthopteran species. (A) WPGMA dendrogram of the 15 orthopteran species found across the seven sampling plots obtained by cluster analysis. (B) Distribution of the four species groups obtained by classification at a distance of 0.8 according to aspect. The mean (+ SD) density values per 5 m<sup>2</sup> for E-facing and S-facing plots are given. Kruskal-Wallis ANOVA revealed that groups (1) and (2) reached significantly higher densities on S-facing plots (respectively  $H_{1,144} = 36.48$ ,  $p < 0.01$ , and  $H_{1,144} = 37.25$ ,  $p < 0.01$ ). However, no significant differences in group (3) density due to aspect occurred ( $H_{1,144} = 0.80$ ,  $p > 0.05$ ), whereas group (4) was significantly more abundant on E-facing plots ( $H_{1,144} = 18.29$ ,  $p < 0.01$ ).




Fig. 5 - Distribution of the four species groups obtained by classification at a distance of 0.8 according to land-use. The mean (+ SD) density values per 5 m<sup>2</sup> corresponding to different land-use systems separated by aspect are given; Aba = abandoned meadow, Mow = mown meadow, Past = pasture. Results of Kruskal-Wallis tests performed separately for E-facing and S-facing plots using land-use as the independent variable are also given. Different letters imply significant differences in pairwise comparison among land-uses under similar aspect as revealed by non-parametric Tukey-type test.

small difference in mean temperatures can play a key role in the regulation of population density (REMMERT, 1985). A key factor determining Orthoptera distribution is the temperature during egg development, and particularly during post diapause egg development (PDD) (VAN WINGERDEN *et al.*, 1991a). Some species that reached

higher densities on S-facing plots are known to have relatively long PDD, as *G. biguttulus* and *S. lineatus* (VAN WINGERDEN *et al.*, 1991a). On the opposite side, *C. parallelus* is a short PDD eurytopic species that can be found on warm as well as cold places (OSCHMANN, 1973; VAN WINGERDEN *et al.*, 1991a). Thermoregulation requirements are also of importance for completing the nymphal development (CARRUTHERS *et al.*, 1992). *Omocestus viridulus* has a short PDD duration (VAN WINGERDEN *et al.*, 1991a), but its larval development is stopped by lower temperatures (WILLOTT, 1997; WILLOTT & HASSAL, 1998) and thus its higher densities are restricted to relatively warmer study grasslands (Tab. 2).

The land-use (under similar irradiation) affects the local occurrence of Orthoptera both directly, through the disturbance caused by haymaking and grazing, and indirectly by modifying microclimate and grassland vegetation structure (Figs 2-3). Stepwise multiple regression highlighted a predominant role played by microclimate (ground level temperature) in determining habitat selection by species from mown meadows (group 1 on Fig. 4.A plus *D. verrucivorus*), while species from abandoned areas (group 2) were chiefly correlated with herb and dwarf shrub biomass, suggesting a more direct relationship with vegetation structure.

Species chiefly related to mown meadows (Fig. 4A; Tab. 2) lay their eggs just above or beneath the soil surface and thus are favoured by the light and thermal climate of mown meadows. The temperature of oviposition sites is strongly influenced by radiation extinction by vegetation (STOUTJESDIJK & BARKMAN, 1992). On mown meadows light absorption is evenly distributed throughout all layers (CERNUSCA *et al.*, 1992), and there is a higher temperature in the egg environment (Fig. 2). Light absorption on the abandoned grasslands takes place chiefly in the upper layer (61-70%) because of the accumulation of dead matter and the lignified stems of dwarf shrubs (CERNUSCA *et al.*, 1992); this layer insulates the ground from extreme irradiation leading to significantly lower temperatures at the soil surface (Fig. 2). Therefore, egg development of species that lay their eggs into the soil, as *D. verrucivorus* (INGRISCH, 1979, HARZ, 1964), *G. biguttulus* (THORENS, 1989), and *S. lineatus* (WALOFF, 1950; LOHER, 1959), is hindered, with a consequent delay in hatching (VAN WINGERDEN *et al.*, 1991a; 1991b).

Species linked to abandoned grasslands lay eggs on the soil surface, e.g. *O. viridulus* (group 2) (WALOFF, 1950; LOHER, 1959), or on vegetation, e.g. *E. brachyptera* (group 2) (SÄNGER, 1977), and *P. griseoptera* (RAGGE, 1965) (upper branch of group 4). They are less damaged by soil insulation from radiation resulting in lower temperatures in the soil. On the other hand, they could clearly be damaged by land-use systems that cause the removal of vegetation and prevent the development of shrubs (such as mowing and grazing).

Not all species with similar egg laying preferences reacted in the same way to land-use, and *C. parallelus*, probably due to its short PDD duration, occurred with high density on differently managed plots, in accordance to the results obtained by VOISIN (1990) and FRICKE & VON NORDHEIM (1992). As observed by ANDERSON *et al.* (1979), CHAPPEL (1983), WHITMAN (1987), and WILLOTT (1997) the thermoregulatory ability during larval development contributes to habitat partitioning in Orthoptera.

Classification results partly agree with SÄNGER (1977), who considered *G. biguttulus*, *S. lineatus* and *C. dorsatus* as species linked with short grass meadows (e.g. mown meadows), and *E. brachyptera*, *P. griseoptera*, *L. bosci*, and *G. rufus* as species more linked with tall grass ecosystems, as abandoned meadows.

No species were linked exclusively with pasture, where the most generalist species occurred, together with a few more sensitive species as occasional guests. This is in accordance with the results of previous studies (BONAVITA *et al.*, 1999; GUIDO *et al.*, 1999), which suggested that the examined pasture plot is over-exploited in relation to the environmental sustainability of the site.

Mown meadows of Mount Bondone are interesting habitats for the local conservation of several species of Orthoptera. Other orthopteran species receive short-term benefit from abandonment. In fact, abandoned grassland areas are transitory habitats. In the long-term, the beech forest would re-colonise the grasslands, causing the almost complete disappearance of the recorded species (GUIDO *et al.*, 1998). The maintenance of the current extensive management system of meadows is of particular importance for the conservation of orthopteran assemblages from the Viote plateau. Unmanaged areas of high and dense vegetation should be left as shelter on the borders of meadows, as suggested by ERHARDT & THOMAS (1991), THORENS (1993), and LÖRTSCHER *et al.* (1994), contributing to a diverse habitat structure and satisfying the requirements of the species linked with abandoned grasslands.

#### ACKNOWLEDGEMENTS

The study is part of a special topic, «Plant-Animal Interactions», integrated into the EC project ECOMONT (Project N. ENV4-CT95-0179, co-ordinated by A. CERNUSCA). We would like to thank A. GALVAGNI for helpful assistance with species identification, U. TAPPEINER and S. THOMASER for providing temperature data and D. GIANELLE and G. TELLINI for helpful assistance with vegetation data. We are also grateful to A. BATTISTI, H. HAUFFE, R.M.J.C. KLEUKERS and to the anonymous referees for their helpful suggestions.

#### RÉSUMÉ

Réponse de la composition des groupements d'Orthoptères aux types d'utilisation des prairies dans le sud des Alpes italiennes – Les effets des systèmes de gestion agricole et de l'hétérogénéité des conditions thermiques sur la composition des groupes d'Orthoptères des prairies secondaires d'un site du sud des Alpes italiennes sont analysés. Les Orthoptères ont été échantillonnés dans sept parcelles gérées de façon différente et selon différentes expositions. Les espèces ont été classées sur la base de leur distribution, dans le but d'identifier des groupes présentant des réactions communes face aux perturbations humaines. Une analyse par régression multiple pas à pas a révélé que l'hétérogénéité des conditions thermiques, exprimée par l'insolation potentielle, joue un rôle clé dans la distribution des espèces. Un groupe d'espèces thermophiles lié aux parcelles exposées au sud a été séparé d'un groupe d'espèces plus généralistes qui colonise également les zones exposées à l'est. On distingue également les espèces qui bénéficient d'un abandon de celles liées principalement aux prairies encore gérées (fauchées). Les différences de densité des groupes selon les parcelles dépendent de la température au niveau du sol et de la structure de la végétation.

#### REFERENCES

- ANDERSON, R.V., TRACY, C.R., & ABRAMSKY, Z. 1979. Habitat selection in two species of short horned grasshoppers. The role of thermal and hydric stress. *Oecologia* 38: 359-374.
- ANTOGNOLI, C. & ZETTEL, J. 1996. Orthoptera communities of differently managed meadows in Ticino. *Mitt. Schweiz. Ent. Ges.* 69: 465-478.
- BARTORELLI, U. 1964. *L'assolazione*. Pubblicazioni dell'Acc. It. Sc. For., 65 pp.
- BÄTZING, W. 1990. *Der italienische Alpenraum*. CIPRA, Kleine Schriften, 7/90.
- BLUMER, P. & DIEMER, M. 1996. The occurrence and consequences of grasshopper herbivory in an alpine grassland, Swiss Central Alps. *Arct. Alp. Res.*, 28(4): 435-440.
- BONAVITA, P., GUIDO, M., & CHEMINI, C. 1999. Patterns of consumer diversity under different land-use practices along the Alpine transect. In: CERNUSCA, A., TAPPEINER, U., & BAYFIELD, N. (eds.), *Land-use Changes in European Mountain Ecosystems. ECOMONT - Concepts and Results*, pp. 256-260. Blackwell Wissenschafts-Verlag, Berlin.
- CARRUTHERS, R.I., LARKIN, T.S., & FIRSTENCHEL, H. 1992. Influence of thermal ecology on the mycosis of a rangeland grasshopper. *Ecology* 73: 190-204.

- CERNUSCA, A., TAPPEINER, U., AGOSTINI, A., BAHN, M., BEZZI, A., EGGER, R., KOFLER, R., NEWSELY, C., ORLANDI, D., PROCK, S., SCHATZ, H., & SCHATZ, I. 1992. Ecosystem research on mixed grassland/woodland ecosystems. First results of the EC-STEP-project INTEGRALP on Mt. Bondone. *Studi Trent. Sci. Natur., Acta Biol.* 67: 99-133.
- CESCATTI, A., CHEMINI, C., DE SIENA, C., GIANELLE, D., NICOLINI, G., & WOHLFAHRT, G. 1999. Monte Bondone composite landscape, Italy. In: CERNUSCA, A., TAPPEINER, U., & BAYFIELD, N. (eds.), *Land-use Changes in European Mountain Ecosystems. ECOMONT - Concepts and Results*, pp. 60-73. Blackwell Wissenschafts-Verlag, Berlin.
- CHAPPELL, M.A. 1983. Metabolism and thermoregulation in desert and montane grasshoppers. *Oecologia* 56: 126-131.
- COXWELL, C.C. & BOCK, C.E. 1995. Spatial variation in diurnal surface temperatures and the distribution and abundance of an alpine grasshopper. *Oecologia* 104: 433-439.
- ERHARDT, A. & THOMAS, J.A. 1991. Lepidoptera as indicators of change in the seminatural grassland of lowland and upland Europe. In: COLLINS, N.M. & THOMAS, J.A. (eds.) *The conservation of insects and their habitats*, pp 213-234, Academic Press, London.
- FAILLA, M.C., LA GRECA, M., LOMBARDO, F., MESSINA, A., SCALI, V., STEFANI, R., & VIGNA TAGLIANTI, A. 1994 - Blattaria, Mantodea, Isoptera, Orthoptera, Phasmatodea, Dermaptera, Embioptera. In: MINELLI, A., RUFFO, S., & LA POSTA, S. (eds.) *Checklist delle specie della fauna italiana*, 36. Calderini, Bologna.
- FIELDING, D.J. & BRUSVEN, M.A. 1995. Ecological correlates between rangeland grasshopper (Orthoptera: Acrididae) and plant communities of Southern Idaho. *Environ. Entomol.* 24 (6): 1432-1441.
- FRICKE, M. & VON NORDHEIM, H. 1992. Auswirkungen unterschiedlicher landwirtschaftlicher Bewirtschaftungsweisen des Grünlandes auf Heuschrecken (Orthoptera, Saltatoria) in der Oker-Aue (Niedersachsen) sowie Bewirtschaftungsempfehlungen aus Naturschutzsicht. *Braunschweiger Naturkundliche Schriften* 4: 59-89.
- GUIDO M., BATTISTI, A., & CHEMINI, C. 1998. Effects of land-use changes on Orthoptera assemblages from Monte Bondone (Southern Alps, Italy). *Redia* 81: 61-72.
- GUIDO M., BONAVITA, P., & CHEMINI, C. 1999. Effects of land-use changes on animal diversity and plant-animal interactions. In: CERNUSCA, A., TAPPEINER, U., & BAYFIELD, N. (eds.): *Land-use Changes in European Mountain Ecosystems. ECOMONT - Concepts and Results*, pp. 170-175. Blackwell Wissenschafts-Verlag, Berlin.
- HARZ, K. 1964. Die Eiablage der heimischen Laubheuschrecken. *Naturwiss. Ges. Bayreuth 1889-1964, Festschrift*: 67-70.
- INGRISCH, S. 1979. Untersuchungen zum Einfluß von Temperatur und Feuchtigkeit auf die Embryogenese einiger mitteleuropäischer Laubheuschrecken (Orthoptera: Tettigoniidae). *Zool. Beitr., N. F.* 25: 343-364.
- INGRISCH, S. & KÖHLER, G. 1998. *Die Heuschrecken Mitteleuropas*. Westarp Wissenschaften, Magdeburg, 460 pp.
- KEMP, W.P., HARVEY, S.J., & O'NEILL, K.M. 1990. Patterns of vegetation and grasshopper community composition. *Oecologia* 83: 299-308.
- KÖHLER, G. 1990. Biogeographisch-ökologische Hintergründe der Faunenveränderung bei Heuschrecken (Saltatoria). *Articulata*, 5: 3-23.
- LOHER, W. 1959. Das Verhalten einiger Feldheuschreckenarten unmittelbar nach der Eiablage. *Nachrbl. Bayer. Ent.* 8: 101-104, 108-110.
- LÖRTSCHER, M., HÄNGGI, H., & ANTOGNOLI, C. 1994. Zoological arguments for managing the abandoned grasslands on Monte San Giorgio, based on data of three invertebrate groups (Lepidoptera, Araneae, Saltatoria). *Mitt. Schweiz. Ent. Ges.* 67: 421-435.
- MORRIS, M.G. 1978. Grassland management and invertebrate animals - a selective review. *Sci. Proc. R. Dublin Soc. Ser. A*, 6: 247-57.
- OSCHMANN, M. 1973. Untersuchungen zur Biotopbindung der Orthopteren. *Faun. Abh. Mus. Tierk. Dresden* 4: 177-206.
- PENZ, H. 1978. *Die Almwirtschaft in Österreich*. Münchner Studien zur Sozial- und Wirtschaftsgeographie, Bd. 15. Verlag Michael Lassleben, Kallmünz/Regensburg.
- QUINN, M.A., KEPNER, R.L., WALGENBACH, D.D., NELSON FOSTER, R., BOHLS, R.A., POOLER, P.D., REUTER, K.C., & SWAIN, J.L. 1991. Habitat characteristics and grasshopper community dynamics on mixed-grass rangeland. *Can. Ent.* 123: 89-105.
- RAGGE, D.R. 1965. *Grasshoppers, Crickets and Cockroaches of the British Isles*. Warne, London, 299 pp.
- REMMERT, H. 1985. Crickets in sunshine. *Oecologia* 68: 29-33.
- SAMWAYS, M.J. 1990. Landforms and winter habitat refugia in the conservation of montane grasshoppers in southern Africa. *Cons. Biol.* 4: 375-382.

- SOUTHWOOD, T.R.E., BROWN, V.K., & READER, P.M. 1979. The relationships of plant and insect diversities in succession. *Biol. J. Linn. Soc.* 12: 327-48.
- STOUTJESDIJK, P. H. & BARKMAN, J. J. 1992. *Microclimate, Vegetation and Fauna*. Opulus Press AB, Knivsta Sweden, 216 pp.
- SÄNGER, K. 1977. Über die Beziehungen zwischen Heuschrecken (Orthoptera: Saltatoria) und der Raumstruktur ihrer Habitate. *Zool. Jahrb. Syst.* 104: 433-488.
- SURBER, E., AMIET, R., & KOBERT, H. 1973. Das Brachlandproblem in der Schweiz. *Ber. Eidg. Anst. Forst. Vers.* 12: 1-138.
- TAPPEINER, U. & CERNUSCA, A. 1993. Alpine meadows and pastures after abandonment. Results of the Austrian MaB-programme and the EC-STEP project INTEGRALP. *Pirineos*, 141-142: 97-118.
- TAPPEINER, U., CERNUSCA, A., SIEGWOLF, R.T.W., SAPINSKY, S., NEWSELY, C., GEISSBUEHLER, P., & STEFANICKI, G. 1999. Microclimate, energy budget, and CO<sub>2</sub> gas exchange of ecosystems. In: CERNUSCA, A., TAPPEINER, U., & BAYFIELD, N. (eds.): *Land-use Changes in European Mountain Ecosystems. ECOMONT - Concepts and Results*, pp. 140-145. Blackwell Wissenschafts-Verlag, Berlin.
- TASSER, E., PROCKS, S., & MULSER, J. 1999. The Impact of land-use on the vegetation in the mountain region. In: CERNUSCA, A., TAPPEINER, U., & BAYFIELD, N. (eds.): *Land-use Changes in European Mountain Ecosystems. ECOMONT - Concepts and Results*, pp. 235-247. Blackwell Wissenschafts-Verlag, Berlin.
- THORENS, P. 1989. Description comparée des oothèques et des oeufs de *Chorthippus mollis* (CHARP.) et de *Chorthippus biguttulus* (L.) (Orthoptera, Acrididae). *Mitt. Schweiz. Ent. Ges.* 62: 87-106.
- THORENS, P. 1993. Effects de la fauche sur une population du Criquet *Chorthippus mollis* (Charp.) (Orthoptera, Acrididae) dans une prairie du pied sud du Jura suisse. *Mitt. Schweiz. Ent. Ges.* 66: 173-182.
- TSCHARNTKE, T. & GREILER, H.J. 1995. Insect Communities, grasses, and grasslands. *Ann. Rev. Entomol.* 40: 535-558.
- TURNER II, B.L. & MEYER, W.B. 1994. Global land-use and land-cover change: an overview. In: MEYER, W.B. & TURNER II, B.L. (eds.), *Changes in land use and land cover: a global perspective*, pp. 3-10. Cambridge University Press, Cambridge.
- VOISIN, J.F. 1990. Observations sur les Orthoptères du Massif Central. 4. *Chorthippus parallelus* (Zetterstedt, 1821) (Orth. Acrididae). *Bull. Soc. Entomol. Fr.* 95: 89-95.
- WALOFF, N. 1950. The egg pods of British short-horned grasshoppers (Acrididae). *Proc. R. Entomol. Soc. Lond. Ser. A Gen. Entomol.* 25: 115-126.
- WHITMAN, D.W. 1986. Developmental thermal requirements for the grasshopper *Taenipoda eques* (Orthoptera, Acrididae). *Ann. Entomol. Soc. Am.* 79: 711-714.
- WILLOTT, S.J. 1997. Thermoregulation in four species of British grasshoppers (Orthoptera: Acrididae). *Funct. Ecol.* 11: 705-713.
- WILLOTT, S.J. & HASSALL, M. 1998. Life-history responses of British grasshoppers (Orthoptera: Acrididae) to temperature change. *Funct. Ecol.* 12: 232-241.
- WINGERDEN, W.K.R.E. VAN, MUSTERS, J.C.M., & MAAKSKAMP, F.I.M. 1991a. The influence of temperature on the duration of egg development in West European grasshoppers (Orthoptera: Acrididae). *Oecologia* 87: 417-423.
- WINGERDEN, W.K.R.E. VAN, MUSTERS, J.C.M., KLEIKERS, R.M.J.C., BONGERS, W., & VAN BIEZEN J.B., 1991b. The influence of cattle grazing intensity on grasshopper abundance (Orthoptera: Acrididae). *Proc. Exp. & Appl. Entomol. N.E.V.* 2: 28-34.
- ZAR, H.J. 1984. *Biostatistical analysis*. Second edition. Prentice-Hall International Editions, 718 pp.

(received September 10, 1999; accepted June 10, 2000)



