

Miscellanea : Sources of carotene and vitamin A in Lake Province, Tanganyika

Autor(en): **McLaren, D.S.**

Objektyp: **Article**

Zeitschrift: **Acta Tropica**

Band (Jahr): **18 (1961)**

Heft 1

PDF erstellt am: **13.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-310942>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Sources of Carotene and Vitamin A in Lake Province, Tanganyika.

By D. S. McLAREN,

East African Institute for Medical Research, Mwanza, Tanganyika.

Vitamin A deficiency is still one of the most prevalent and devastating nutritional diseases in certain parts of the world (KINNEY & FOLLIS, 1958). It is made all the more important by the facts that the eye is the part of the body most frequently affected, and that the most severe changes which are known as xerophthalmia and keratomalacia, frequently leading to blindness, occur usually in very young children.

Whilst the condition is most severe and widespread amongst the rice-eating populations of South-East Asia, it has become evident from surveys carried out in recent years that it also occurs in parts of Africa, especially where the red palm (*Eloesia guineensis*) does not grow. Such areas are the whole of North Africa, Northern Nigeria, Ruanda Urundi, and the drier parts of East Africa. MCKENZIE (1939) recognised the problem of night blindness in Central Tanganyika 20 years ago, and surveys carried out recently from this Institute in the same area have shown that the whole spectrum of vitamin A deficiency, from keratomalacia in young children to xerosis conjunctivae, Bitot's spots, and night blindness in older children and young adults, still constitutes an important problem. Plasma carotenoid and vitamin A values are generally low in Mwanza (McLAREN, 1959).

Experience gained in recent years in the field of feeding programmes in under-developed countries has shown the importance of developing locally available foodstuffs, and the present work has been undertaken with a view to combatting the vitamin A deficiency problem in the Lake and Central Provinces of Tanganyika.

1. The vitamin A content of the livers of Lake Victoria fish.

The fish were obtained within an hour or two of being caught, and in many instances were still breathing when brought to the laboratory. Each fish was weighed and measured from tip of head to commencement of tail. The whole liver was dissected out and weighed, and then weighed fractions were taken

TABLE 1.

Vitamin A content of livers of fish commonly found at the southern end of Lake Victoria.

Scientific name	Local name	No. of observations	Mean length (cm)	Mean body weight (g)	Mean weight liver (g)	Mean concentrat. vitamin A (i.u./g)	Standard Deviation
<i>Schilbe mystus</i>	nembe	7	14.0	54.3	0.49	1659	893
<i>Labeo victorianus</i>	ningu	5	18.8	110.0	1.18	1197	669
<i>Haplochromis</i> spp.	furu	8	10.4	55.0	0.85	2846	2141
<i>Tilapia esculenta</i>	ngege	6	17.5	147.5	1.70	1689	770
<i>Synodontis victorianus</i>	gogogo	6	11.5	48.0	1.30	4375	6999
<i>Alestes nurse</i>	soga	6	13.6	76.0	0.82	5446	3090
<i>Tilapia variabilis</i>	mbiru	6	18.0	260.0	2.38	2070	1134
<i>Mormyrus kannume</i>	mbete	4	22.2	300.0	4.38	3902	3205
<i>Bagrus docmac</i>	mbofu	7	27.2	1110.0	6.23	4363	1152
<i>Clarias</i> spp.	mumi	6	27.9	621.1	5.40	5686	3362
<i>Protopterus aethiopicus</i>	kambari mamba	1	42.0	2500.0	34.60	725	—

TABLE 2. Carotene content of green leaves.

Scientific name	Local name	No. of observations	State	Mean carotene fresh leaves (i.u./100 g)	Standard Deviation
<i>Manihot utilisima</i> Pohl.	muhogo	8	old: uncooked	12,800	368
<i>Manihot utilisima</i> Pohl.	muhogo	8	old: cooked 1 hr.	6,550	147
<i>Manihot utilisima</i> Pohl.	muhogo	8	young: uncooked	7,530	188
<i>Manihot utilisima</i> Pohl.	muhogo	8	young: cooked ½ hr.	7,010	206
<i>Cucumis sativa</i> L.	matango	8	uncooked	8,150	423
<i>Cucumis sativa</i> L.	matango	8	cooked 1 hour	4,540	182
<i>Cucurbita</i> spp.	nyamwanga	8	uncooked	10,050	463
<i>Cucurbita</i> spp.	nyamwanga	6	cooked 1 hour	9,050	512
<i>Vigna unguiculata</i> Walp.	kunde	8	uncooked	9,510	403
<i>Vigna unguiculata</i> Walp.	kunde	6	cooked 1 hour	5,010	91
<i>Amaranthus dubius</i> Mart.	mchicha (native)	8	uncooked	10,690	552
<i>Amaranthus dubius</i> Mart.	mchicha (native)	8	cooked ½ hour	8,000	290
<i>Amaranthus cruentus</i> L. var. <i>paniculatus</i> (L.) Thellg.	mchicha (foreign)	7	uncooked	6,170	345
<i>Amaranthus cruentus</i> L. var. <i>paniculatus</i> (L.) Thellg.	mchicha (foreign)	9	cooked ½ hour	5,080	259
<i>Phaseolus vulgaris</i> L.	maharage	6	uncooked	8,070	197
<i>Phaseolus vulgaris</i> L.	maharage	6	cooked 1 hour	6,270	152
<i>Ipomoea batatas</i> (Linn.) Lam.	matembele	6	uncooked	5,830	164
<i>Ipomoea batatas</i> (Linn.) Lam.	matembele	6	cooked ½ hour	3,970	156
<i>Gynandropsis gynandra</i> (L.) Briq.	mgagani	8	uncooked	7,550	293
<i>Gynandropsis gynandra</i> (L.) Briq.	mgagani	8	cooked 1 hour	8,090	576
<i>Gisekia pharnaceoides</i> L.	imbala	4	uncooked	4,600	83
<i>Gisekia pharnaceoides</i> L.	imbala	4	cooked ½ hour	2,800	111
<i>Corchorus trilocularis</i> L.	lunani	6	uncooked	5,000	238
<i>Corchorus trilocularis</i> L.	lunani	6	cooked ½ hour	4,150	91
<i>Sesamum angustifolium</i> Engl. forma	mlenda	2	uncooked	11,600	0
<i>Sesamum angustifolium</i> Engl. forma	mlenda	2	cooked ½ hour	5,500	56

for duplicate determination of the vitamin A content by the antimony trichloride method.

Table 1 shows the details of the 11 species of fish which are commonly found at the southern end of Lake Victoria. The extreme variability of liver concentration of vitamin A is shown by the very large standard deviations. Different tribes have different preferences for these fish. The Wasukuma and other tribes of Lake Province eat most of them, but do not like *Mormyrus*, *Synodontis*, or *Protopterus*.

2. *The carotene content of certain green leaves.*

The 12 kinds of green leaves examined are all readily available most of the year near the lake. Fresh samples of the leaves weighing 1 gram were used, and the carotene was separated on an alumina adsorption column, following the method described by BOOTH in MOORE (1957).

Table 2 gives details of the carotene content of the leaves both fresh and after cooking for various periods. The leaves were cooked in the laboratory with amounts of water and for lengths of time similar to those commonly used by the people of this area, although these will naturally vary considerably in practice. In all instances except one, there was from slight to considerable loss of carotene after cooking. In the case of *Gynandropsis gynandra* (L.) Briq. there appeared to be a very slight increase in the mean value after cooking, and it may be that carotene was made available by cooking for of all the leaves examined, this one requires the most cooking according to local experience.

Summary.

1. The vitamin A content of the livers of Lake Victoria fish, and the carotene content of certain green leaves have been determined.

2. These are shown to be excellent sources of vitamin A readily available to the people of Lake and Central Provinces of Tanganyika, where vitamin A deficiency is common.

Acknowledgements.

Mr. M. H. ATKINSON, Fisheries Officer, Lake Victoria Fisheries Service, very kindly supplied the fish, and I am indebted to Miss D. M. NAPPER of the East African Herbarium, Nairobi, for identification of the plants. Permission to publish was granted by the Director, East African Institute for Medical Research, and Mr. MAURICE KWENA assisted with the vitamin A and carotene determinations.

References.

- BOOTH, V. H. (1957). In *Vitamin A* by MOORE, T. p. 570.— Elsevier, Amsterdam.
- KINNEY, T. D. & FOLLIS, R. H. Jr. (1958). Nutritional Disease: Proceedings of the Conference on Beriberi, Endemic Goiter and Hypovitaminosis A. Princeton, June 1-5, 1958. — *Fed. Proc.* 17, Part 2, No. 2, p. 1-144.
- MCKENZIE, A. (1939). An examination of vitamin A deficiency among African natives by the visual dysadaptation test. — *Trans. roy. Soc. trop. Med. Hyg.* 32, 717.
- MCLAREN, D. S. (1959). Urinary excretion of vitamin A in pregnancy. — *Proc. Nutr. Soc.* 18, p. xxx-xxxii.
-