

Miscellanea : Lack of incorporation of tritiated thymidine in "Trypanosoma vivax in vitro"

Autor(en): **Isoun, Miriam J. / Isoun, T.T.**

Objektyp: **Article**

Zeitschrift: **Acta Tropica**

Band (Jahr): **31 (1974)**

Heft 4

PDF erstellt am: **29.06.2024**

Persistenter Link: <https://doi.org/10.5169/seals-311967>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Lack of Incorporation of Tritiated Thymidine in *Trypanosoma vivax in vitro*

MIRIAM J. ISOUN and T. T. ISOUN

In order to determine some of the characteristics of the cell cycle and *in vitro* metabolism of *T. vivax* pathogenic in ruminants, we designed labeling experiments with tritiated thymidine (^3H -thymidine) for preparation of autoradiographs and scintillation counting. Our methods were those routinely used by cytologists with mammalian cells, but included relatively large doses of thymidine (up to $15\ \mu\text{Ci/ml}$ medium) as described for other species of trypanosomes: *T. musculi* (VIENS & TARGETT, 1972), *T. brucei* (BALBER, 1971), *T. evansi* and *T. gambiense* (INOKI & TADASUKE, 1969), and *T. mega* (STEINERT & STEINERT, 1962). The results were disappointing as background was high, no specific label was apparent, and scintillation counts were low.

By using *T. brucei* passaged in rats, we were able to obtain large numbers of trypanosomes to perfect our experimental system. Under the conditions available to us, we found the most important factors in obtaining high specific label (Fig. 1)




Fig. 1. *T. brucei* labeled with $15\ \mu\text{Ci/ml}$ ^3H -thymidine in Hanks saline for 2 hours. Fixed slides were mounted with Kodak (Rochester, New York, U.S.A.) AR-10 stripping film for 4 weeks.




Fig. 2. *T. vivax* labeled with 25 $\mu\text{Ci/ml}$ ^3H -thymidine in Hanks saline for 8 hours. Fixed slides were dipped in 1% PPO-toluene before mounting with stripping film. There is no specific label; a few background grains are apparent (arrow).

were the use of large amounts of high specific activity ^3H -thymidine (Amersham, Buckinghamshire, England: 15 $\mu\text{Ci/ml}$, Sp. Act. 26 Ci/mmol) with correspondingly low amounts of thymidine in the suspending medium, and the control of humidity. We were able to control humidity during incubation by the extensive use of silica gel and improved air conditioning. Four weeks incubation of the mounted autoradiographs was adequate. The use of scintillation autoradiography (BALBER, 1971) was not necessary, and when used, increased the scatter of the grains. The high doses of tritium were not toxic to the cells, as survival (counts of organisms using haemocytometer) of the trypanosomes was similar in labeled and unlabeled cultures. Scintillation counts of solubilized (NCSTTM Solubilizer, Amersham-Searle, Illinois, U.S.A.) trypanosomes were adequate (Table 1, A–D) for studies of comparative metabolism with different culture conditions (ISOUN & ISOUN, 1974a).

By using the improvements described, we set up experiments with *T. vivax*: trypanosomes were incubated in Hanks saline with 5% homologous serum and 15–24 $\mu\text{Ci/ml}$ ^3H -thymidine for two to eight hours. In no autoradiographs was specific label incorporated (Fig. 2). Scintillation counts of solubilized *T. vivax* (Table 1 F and G) were insignificant as compared to controls (Table 1, E and H). The lack of incorporation of ^3H -thymidine by *T. vivax* is not indicative of sluggish




Fig. 3. Effect of foetal calf serum and homologous serum as compared to a saline control on protein synthesis in *T. vivax in vitro* at 25°. The incubation mixture contained 3 ml saline, 3 μ Ci ¹⁴C-amino acids (10 mCi/mmol), 0.72×10^6 trypanosomes, and 0.5 ml variable: A, homologous serum, B, foetal calf serum, and C, Hanks saline. Trypanosomes were prepared for scintillation counting as described in Table 1.

metabolism of dying organisms, as there was evidence of growth of trypanosomes over the first 24 hours *in vitro* (ISOUN & ISOUN, 1974b), and incorporation of ¹⁴C-labeled amino acids (Amersham Buckinghamshire, England) continued during this period *in vitro* (Fig. 3).

The failure to incorporate ³H-thymidine into the DNA of *T. vivax* has not been reported; indeed we are not aware of the inability of other Trypanosomatidae except *Crithidia fasciculata* (GUTTERIDGE & AL CHALABI, 1973) to incorporate exogenous thymidine. *Plasmodium berghei* and *P. vinckei* (WALSH & SHERMAN, 1968), psitticosis organisms (MOULDER, 1966) and *Euglena* (SAGAN, 1965) are reported to be unable to utilize exogenous thymidine. It is logical to deduce that these organisms are able to synthesize their own pyrimidines for DNA synthesis and thus cell division, and multiplication. However, the mechanism by which *T. vivax* is able to synthesize pyrimidines for DNA synthesis is not known. It is tempting to speculate that like the plasmodial group of organisms that synthesize their own pyrimidines, *T. vivax* is capable of fixing carbon dioxide to form carbamyl phosphate, and with aspartic acid, synthesize the pyrimidines. Two enzymes, thymidine kinase and thymidine synthetase, the formation of which depends on the derepression action of the appropriate cistron (PRESCOTT & STONE, 1969), may determine the availability of pyrimidines for DNA synthesis and cell division.

Table 1. Scintillation counts of *T. brucei* and *T. vivax*, isolated from whole blood by erythrocyte lysis (LEEFLANG et al., 1974) followed by DEAE cellulose elution (LANHAM & GODFREY, 1970), and incubated with 15 μ Ci/ml 3 H-thymidine. After incubation, 1 ml of trypanosome suspension was washed 2 \times in 10 ml Hanks saline + 5 % foetal calf serum, and resuspended in 0.5 ml saline. 0.2 ml of this suspension was placed in each of two scintillation vials to which 0.7 ml NCS solubilizer was added at room temperature overnight. 5 ml toluene fluor containing 0.4 % PPO (2,5-Diphenyl-oxazole) and 0.005 % POPOP [1,4-bis 2-(5 phenyloxzoly)-Benzene; phenyloxzolyphenyl-oxazolyphenyl] was added and the vials counted in a liquid scintillation counter (Packard, Model 3101) for 1 minute. The values given below represent averages of the duplicate vials.

Sample	Tryp. count	Species	Time labeled	Medium	Counts/min
A	<i>T. brucei</i>	3.0×10^6	4 hours	saline ¹ + 5 % serum ²	2583
B	<i>T. brucei</i>	3.0×10^6	6 hours	saline + 5 % serum	5281
C	<i>T. brucei</i>	3.0×10^6	4 hours	199 H ³ + 15 % serum	638
D	<i>T. brucei</i>	3.0×10^6	6 hours	199 H + 15 % serum	1117
E	<i>T. brucei</i>	3.0×10^6	0	saline + 5 % serum	26
F	<i>T. vivax</i>	1.0×10^6	2 hours	saline + 5 % serum	30
G	<i>T. vivax</i>	1.0×10^6	8 hours	saline + 5 % serum	32
H	None: background	0			14

¹ Hanks basic salt solution, GIBCO, Grand Island, New York.

² Foetal calf serum, GIBCO.

³ Medium 199 with HEPES buffer, GIBCO.

Additionally *T. vivax* like *C. fasciculata* (GUTTERIDGE & AL CHALABI, 1973) may have, in the presence of exogenous thymidine, high concentration (activity) of the enzyme thymidine phosphorylase (thymidine: orthophosphate deoxyribosyl-transferase EC 2.4.2.4) capable of converting tritiated thymidine to thymine – a substance not utilized by the organism. Compounds that are capable of inhibiting enzymes involved in the pathway of the bio-synthesis of pyrimidines in *T. vivax* may serve as chemotherapeutic agents in *T. vivax* infection in ruminants.

In the search for improved media for the *in vitro* cultivation of *T. vivax* it may be found that factors which enhance *de novo* synthesis of pyrimidines in these organisms may also promote growth and multiplication. Currently we are investigating these factors.

Acknowledgement

We wish to thank the Rockefeller Foundation and the University of Ibadan Senate for financial support of this work.

References

- BALBER, A. E. (1971). Pleomorphism and the physiology of *Trypanosoma brucei*. – Ph.D. thesis, The Rockefeller University, New York.

- INOKI, S. & TADASUKE, O. (1969). Studies on the DNA synthesis in *Trypanosoma gambiense* and *Trypanosoma evansi* by means of ³H-thymidine and autoradiography. – Jap. J. Parasit. 18, 217–225.
- GUTTERIDGE, W. E. & AL CHALABI, K. (1973). Thymidine phosphorylase activity in *Crithidia fasciculata*. – Progress in Protozoology. Abstracts of papers read at the fourth International Congress on Protozoology, Clermont – Ferrand.
- ISOUN, M. J. & ISOUN, T. T. (1974a). The effect of HEPES buffered media on the *in vitro* cultivation of *Trypanosoma vivax* and *Trypanosoma brucei*. – Z. Tropenmed. Parasit. 25, 283–287.
- ISOUN, T. T. & ISOUN, M. J. (1974b). *In vitro* cultivation of *Trypanosomes vivax* pathogenic to ruminants. – (In press.)
- LANHAM, S. M. & GODFREY, D. G. (1970). Isolation of salivarian trypanosomes from man and other mammals using DEAE cellulose. – Exp. Parasit. 28, 521–534.
- LEEFLANG, P., BLOTKAMP, C., GODFREY, D. G. & KILGOUR, V. (1974). A hypotonic lysis method for concentrating trypanosomes from infected blood. – Trans. roy. Soc. trop. Med. Hyg. (In press.)
- MOULDER, J. W. (1966). The relation of psitticosis group (*Chlamydiae*) to bacteria and viruses. – Ann. Rev. Microbiol. 20, 107–130.
- PRESCOTT, D. M. & STONE, G. E. (1969). Replication and function of the protozoan nucleus, pp. 119–146. In: Research in Protozoology, Vol. I. Ed. by Tze Tuan Chen. – Oxford: Pergamon Press.
- SAGAN, L. (1965). An unusual pattern of tritiated thymidine incorporation in *Euglena*. – J. Protozool. 12, 105–109.
- STEINERT, M. & STEINERT, G. (1962). La synthèse de l'acide désoxyribonucléique au cours du cycle de division de *Trypanosoma mega*. – J. Protozool. 9, 203–211.
- VIENS, P. & TARGETT, G. A. T. (1972). Autoradiography of blood forms of *Trypanosoma musculi*. – Can. J. Microbiol. 18, 533–534.
- WALSH, C. J. & SHERMAN, I. W. (1968). Purine and pyrimidine synthesis by the avian malaria parasite *Plasmodium lophurae*. – J. Protozool. 15, 763–770.