

Field and laboratory observations on parasitization rates of "Glossina" puparia by "Syntomosphyrum" species in Nigeria : short communication

Autor(en): **Onyiah, J.A. / Riordan, K.**

Objektyp: **Article**

Zeitschrift: **Acta Tropica**

Band (Jahr): **35 (1978)**

Heft 3

PDF erstellt am: **09.08.2024**

Persistenter Link: <https://doi.org/10.5169/seals-312393>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Department of Entomology, Nigerian Institute for Trypanosomiasis Research,
Kaduna, Nigeria

Field and laboratory observations on parasitization rates of *Glossina* puparia by *Syntomosphyrum* species in Nigeria

Short communication

J. A. ONYIAH, K. RIORDAN

In Nigeria, field studies on *Glossina* species have rarely revealed parasites which are effectively involved in the reduction of natural population levels. Suggestions have been made for introduction of such controlling agents from areas in and outside Africa (Laird, 1977). There is a worldwide outcry against the blanket use of persistent insecticide formulations for control of *Glossina* species in the field. The introduction of biocontrol methods thus demands attention.

Syntomosphyrum species (Order, Hymenoptera; family, Eulophidae) have been tested as biocontrol agents of tsetse populations in Malawi (Lamborn, 1925), Tanzania (Nash, 1933) and Nigeria (Lloyd et al., 1927). In the third area *Syntomosphyrum albiclavus* (Kerrich), bred from materials obtained from England, was introduced in Sherifuri (near Azare, Bauchi State) to control *Glossina morsitans submorsitans* (Newstead) and *G. tachinoides* (Westwood). No parasitized field puparium was recovered. *S. glossinae* (Waterston) is incapable of penetrating sandy soils far enough to reach the tsetse puparia (Nash, 1933).

S. glossinae was first recorded in West Africa by Nash (1947) as a parasite of field-collected *G. palpalis* (R.-D.) puparia in the N.I.T.R. laboratories in Kaduna, Nigeria. Nash (1955) also reported the parasitization of *Periplaneta americana* (L.) oothecae by the same species within the same insectary. This report was later refuted by Jordan (1956). From the scanty information available, Laird (1977) summarized the life history of *Syntomosphyrum* spp. Three-day-old females drill a hole in the puparial case and oviposit about 40–50 eggs into the subpuparial space. A wide variety of dipteran puparia including *Glossina* spp. can be used. Hatching occurs within 48 h and the 3rd instar larva is attained 24 h later. The 4th instar has a 5-day duration. The prepupal and pupal stages last 2 and 10 days respectively. After eclosion, adults bite a small hole in the host puparium, through which they escape. Copulation then takes place

Correspondence: Dr. J. A. Onyiah, Nigerian Institute for Trypanosomiasis Research, Private Mail Bag 2077, Kaduna, Kaduna State, Nigeria

Table 1. Parasitization-rate of field-collected *G. palpalis* puparia by *Syntomosphyrum* species in Nigeria

Year	Number of puparia examined	Number parasitized	%
1972	16,877	10	0.059
1973	21,890	11	0.057
1974	20,367	18	0.088
Totals	59,134	39	0.066

almost immediately. Females are capable of reproducing parthenogenetically, all offspring being males.

The low rates of parasitization (0.04%–2.4% of puparia examined) of *Glossina* spp. in nature in East Africa are given by Laird (1977). In Nigeria the incidence of parasitization recorded for *G. palpalis* puparia collected over a period of 3 years, is given in Table 1. The parasitized puparia were collected at Kaffin-Koro, Kwakuti, Kuje, Akerri, Guni and Bwari in Niger State of Nigeria. The overall rates of incidence varied little from locality to locality or from year to year, excepting Kuje (total rate for the 3 years, 0.133%) where the incidence was about twice the average.

At N.I.T.R., an attempt was made in January 1977 to raise laboratory colonies of *G. palpalis* and *G. tachinoides* in the insectary. 44,250 and 11,160 field-collected puparia of the two species respectively from the above collecting localities were used. Onyiah (1977) described the routine maintenance methods for adults and puparia. In May, a large number of *Syntomosphyrum* spp. was observed in the tsetse emergence cages. Cages were covered with fine-mesh netting, which ensured separation of the field- and laboratory-bred puparia within the insectary. All unemerged puparia were then examined for the tell-tale emergent holes. Those without holes were dissected for evidence of infection. The dissection indicated that parasitization could become established in well developed pharate flies in the puparium. The remains of tsetse flies (heads, probosces and legs) were seen in some cases. Once, 22 adults were observed crawling out from the emergence hole of a *Glossina* puparium. The summary of observations is given in Table 2. The parasitization rate amongst laboratory-bred puparia (of both *Glossina* spp.) was twice that amongst puparia from the field. Unlike the field-collected puparia (Table 1), parasitization was significantly higher amongst puparia kept in the insectary.

S. glossinae occurs naturally in Nigeria; the presence of *S. albiclavus* is not definitely confirmed. Reliable separation of the two species is only by a cross-breeding technique. No specimen of *Syntomosphyrum* has ever been taken during field studies in Nigeria. This indicates that in nature the hymenopterous parasite of various dipteran puparia occurs in very low densities. *Glossina*

Table 2. Rate of parasitization by *Syntomosphyrum* species of both field-collected and laboratory-bred *G. palpalis* kept in the Kaduna tsetse breeding laboratory

Field-collected puparia	Laboratory-bred puparia	Total No. unemerged	No. parasitized	No. found parasitized on dissection	Total parasitized	% of parasitization amongst unemerged puparia
<i>G. palpalis</i>						
44,450	–	7,524	666	19	685	9.1
–	5437	1,433	290	38	328	22.9
<i>G. tachinoides</i>						
11,160	–	2,095	785	–	785	37.5
–	413	209	145	–	145	69.4
Totals		11,261	1,886	57	1,943	17.3

larvae rarely pupariate on the soil surface. Therefore, few are exposed to infection since the hymenopterous parasites are unable to dig deeply into the sandy soil. This restricting factor is probably responsible for the very low parasitization rates amongst field-collected puparia. The barrier constitutes a major disadvantage in the use of the parasite in biocontrol of *Glossina* populations. Conversely, the parasite is highly effective in the destruction of *Glossina* puparia in an insectary; where puparia are exposed and readily available to ovipositing females. The higher infection rates amongst laboratory-bred puparia suggest that younger puparia are easier to infect than field-collected ones of older ages.

There is no evidence that exposed *Glossina* larvae, are infected before pupariation in the soil. This needs further investigation, as releases of the parasites can therefore be made in the dry season localized larviposition sites in the field. Laboratory experiences emphasize the care necessary to prevent the presence of the parasite in an insectary.

Acknowledgments. We are profoundly grateful to all field staff of the Entomology Department, who were energetically involved in field collection of puparia during 1972 to 1974, to Miss F. O. Apata, Mallam Abare Lawiye and others who were actively involved in the insectary aspect of this report and to the Director, N.I.T.R., for permission to publish.

- 1 Jordan A. M.: A note on the parasitization of the oothecae of *Periplaneta americana* (L) by the chalcid, *Syntomosphyrum glossinae* Wtstn – a correction. Bull. ent. Res. 47, 683 (1956).
- 2 Laird M. (ed.): Tsetse. The future for biological methods in integrated control. I.D.R.C. – 077e. (1977).
- 3 Lamborn W. A.: An attempt to control *Glossina morsitans* by means of *Syntomosphyrum glossinae* Waterston. Bull. ent. Res. 15, 303–309 (1925).
- 4 Lloyd L., Johnson W. B., Rawson P. H.: Experiments in the control of tsetsefly. Bull. ent. Res. 17, 423–455 (1927).

- 5 Nash T. A. M.: The ecology of *Glossina morsitans* Westw., and two possible methods for its destruction. Part. II. Bull. ent. Res. 24, 163–195 (1933).
- 6 Nash T. A. M.: A record of *Syntomosphyrum glossinae* from Nigeria. Bull. ent. Res. 38, 525 (1947).
- 7 Nash T. A. M.: A note on the parasitization of oothecae of *Periplaneta americana* (L.) by the chalcid, *Syntomosphyrum glossinae* Wtstn. Bull. ent. Res. 46, 111–112 (1955).
- 8 Onyiah J. A.: The Kaduna *Glossina palpalis* (R.-D.) colony, 1972–1976. 15th Meeting of ISCTRC, Banjul, Gambia (in press) (1977).

International Conference on Malaria and Babesiosis Mexico City, Mexico, April 30 through May 3, 1979

An International Conference on Malaria and Babesiosis will be held in Mexico City April 30 through May 3, 1979. Outstanding scientists from throughout the world will present current knowledge on the diseases, their treatment, transmission, immunology, immunopathology, immunization, morphologic and biochemical properties and the cultivation of the causative agents.

In conjunction with the conference, a workshop on malaria and babesiosis will be held on Monday Evening, April 30. Those wishing to make a presentation at the workshop, please submit an informative abstract for consideration before December 31, 1978.

Attendance at the Conference is open to the public, permitting interested individuals for the first time to obtain direct information from the research scientists about the recent exciting developments on these diseases.

The symposium is being jointly organized by the Instituto Nacional de Investigaciones Pecuarias of Mexico, The Ohio State University, The University of Illinois and The Rockefeller Foundation.

Those interested in receiving the program and registration materials or submitting an abstract for the workshop should send their full name and mailing address to the Conference Organizing Committee in care of Professor Miodrag Ristic, College of Veterinary Medicine, University of Illinois, Urbana, IL 61801.

World Health Organization Special Programme for Research and Training in Tropical Diseases

The search for scientists to search for the tools to control the tropical diseases

The search for new tools to control disease in the tropical countries requires scientists of many disciplines. Molecular and cell biologists, biochemists, immunologists, parasitologists and entomologists are among those whose contributions are needed.

The research areas covered by the Special Programme for Research and Training in Tropical Diseases are:

- malaria, schistosomiasis, filariasis, trypanosomiasis, leishmaniasis and leprosy;
- epidemiology, biomedical sciences, biological control of vectors, and socio-economic research.

Specific fields within these research areas are developed by the Programme's Scientific Working Groups. Scientists from any country are welcome to submit proposals for research grants within the specific fields. All enquiries should be addressed to:

The Special Programme for Research and
Training in Tropical Diseases
World Health Organization
1211 Geneva 27, Switzerland