

Zeitschrift: Acta Tropica

Band: 36 (1979)

Heft: 3

Artikel: The presence of basic proteins associated to the kinetoplast-DNA of "Trypanosoma cruzi" epimastigotes : short communication

Autor: Souza, W. de / Souto-Padrón, T. / Docampo, R.

DOI: <https://doi.org/10.5169/seals-312529>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 19.11.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Institutos de Biofísica y Microbiología, Universidade Federal do Rio de Janeiro, Brasil





The presence of basic proteins associated to the kinetoplast-DNA of *Trypanosoma cruzi* epimastigotes

Short communication

W. DE SOUZA, T. SOUTO-PADRÓN, R. DOCAMPO¹, LIANE CHAVES, W. LEON

Trypanosoma cruzi epimastigotes possess a single mitochondrion with a differentiated DNA containing region, which is termed the kinetoplast. Steinert (1965), based on cytochemical techniques has suggested that, like mitochondrial DNA, the kinetoplast DNA is not associated with basic proteins. The existence of a highly compact, well-structured association of molecules raises the question of stability. One wonders how the negatively charged DNA molecules can remain in such close proximity without any basic proteins to neutralize the charge (Simpson, 1972). Evidence of the existence of basic proteins in the kinetoplast has been presented by Souto-Padrón and De Souza (1978). However, the possibility that these basic proteins are not associated with DNA has not been definitely ruled out. One way of demonstrating whether these basic proteins are associated with K-DNA is to examine their alterations after inducing the loss of K-DNA with ethidium bromide (EB). *T. cruzi* epimastigotes (Y strain) were grown in Warren's liquid medium (Warren, 1960) for 4 days either in the presence or absence of 0.5 $\mu\text{g}/\text{ml}$ ethidium bromide. The concentration of EB used induced 50% diskinetoplasty of the parasite population as determined by light microscopical observation of Giemsa stained cells. The cells were harvested and washed twice by centrifugation in 0.15 M NaCl and prepared for electron microscopy (Souto-Padrón and De Souza, 1978). For cytochemistry, the ethanolic phosphotungstic acid (E-PTA) method to detect basic proteins was employed (Souto-Padrón and De Souza, 1978). As described previously (Delain and Riou, 1969) the initial effect of growth in a low concentration of EB on the ultrastructural appearance of K-DNA in *T. cruzi* culture forms is a disruption of the orderly double-layered arrangement of DNA fibrils. This occurred in several locations along the DNA band as seen in longitudinal section,

¹ Permanent address: Instituto de Química Biológica, Facultad de Medicina, Universidad de Buenos Aires, Buenos Aires, Argentina



producing several "arcs" of fibers. Then the DNA band either fragmented or retracted from the membrane, or formed a globular mass with a filamentous structure. In both cases the DNA mass remained attached to the membranes by fibers. The final phase of the process involved a further clumping of the DNA and finally an apparent decrease in the amount of DNA present.

T. cruzi epimastigotes stained with E-PTA displayed a homogenous electron-dense reaction in the nucleus, on some microtubules (basal body and peripheral doublets of the flagellum) and at the periphery of the kinetoplast DNA (Fig. 1). A similar clumping and retraction and eventual decrease in quantity of the material stained with E-PTA was observed in epimastigotes grown in EB and stained with E-PTA, thus indicating a close relationship between K-DNA and basic proteins (Figs. 2–4).

Acknowledgments. This work has been supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Financiadora de Estudos e Projetos (FINEP).

- 1 Delain E., Riou G.: Ultrastructure des altérations du DNA du kinétoplaste de *Trypanosoma cruzi* traité par le bromure d'éthidium. C.R. Acad. Sci. Paris 268, 1327–1330 (1969).
- 2 Simpson L.: The kinetoplast of the haemoflagellates. Int. Rev. Cytol. 32, 139–207 (1972).
- 3 Souto-Padrón T., De Souza W.: Ultrastructural localization of basic proteins in *Trypanosoma cruzi*. J. Histochem. Cytochem. 26, 349–358 (1978).
- 4 Steinert M.: L'absence d'histone dans le kinétonucleus des trypanosomes. Exp. Cell. Res. 39, 69–72 (1965).
- 5 Warren L.: Metabolism of *Schizotripanum cruzi* Chagas. I. Effect of culture age and substrate concentration on respiratory rate. J. Parasit. 40, 529–539 (1960).

Fig. 1. Control cell showing PTA reaction at the periphery (arrow) of the kinetoplast (k). $\times 45,000$.
Figs. 2–4. Ethidium bromide-treated cells showing reaction product in the basal body (bb), flagellar microtubules (f) and in the kinetoplast (k). In some cells both strongly and less intensely reacting areas can be seen (Figs. 2–3). In other cells the reaction product appears on electron-dense spherical bodies (Fig. 4, arrow). $\times 45,000$.

