

A new in vitro test for human serum resistance of "Trypanosoma (T.) brucei" : short communication

Autor(en): **Jenni, L. / Brun, R.**

Objektyp: **Article**

Zeitschrift: **Acta Tropica**

Band (Jahr): **39 (1982)**

Heft 3

PDF erstellt am: **14.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-312986>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Swiss Tropical Institute, Basel, Switzerland

A new in vitro test for human serum resistance of *Trypanosoma (T.) brucei*

Short communication

L. JENNI, R. BRUN

It has been known since Laveran (1902) that normal human serum has a trypanocidal effect on *Trypanosoma b. brucei*. After various attempts to identify the cytotoxic factor(s) (Aaronovitch and Terry 1972; Hawking et al., 1973; Hawking, 1979; Van Meirvenne et al., 1973), Rifkin (1978) found a high density lipoprotein in normal human blood and serum with trypanocidal activity.

The trypanocidal activity of normal human blood or serum has been used by Rickman and Robson (1970, 1972) who developed a simple indirect test to determine the subspecific identity of *Trypanosoma (T.) brucei*. This test, known as the blood-incubation infectivity test (BIIT) includes incubation of trypanosomes in human blood or serum in vitro at 37° C and subsequent inoculation of the suspension into susceptible laboratory animals. *T. b. rhodesiense* and *T. b. gambiense* are relatively resistant to the cytotoxic effect of the normal human blood or serum whereas *T. b. brucei* is neutralized.

Brun et al. (1981) described an in vitro culture system for bloodstream forms of different subspecies of the subgenus *Trypanosoma (T.) brucei* which enabled the continuous growth of *T. b. rhodesiense* and *T. b. gambiense* in the presence of normal human serum. In this system, stocks of *T. b. brucei* were rapidly lysed whereas they grew in inactivated rabbit serum. This in vitro culture system was used to test the human serum resistance of several stocks of *Trypanosoma (T.) brucei*.

The in vitro test system consists of a feeder layer of fibroblast-like cells derived from embryos of *Microtus montanus*, with HEPES-buffered Minimum Essential Medium, with Earle's Salts (EMEM), supplemented with 15% normal human serum in the test wells and 15% inactivated rabbit serum in the control wells (Brun et al., 1981).

Correspondence: Dr. L. Jenni, Swiss Tropical Institute, Socinstrasse 57, CH-4051 Basel, Switzerland

Table 1 shows the stocks and clones derived from primary isolates of *Trypanosoma (T.) brucei* which were tested. The history of these isolates, except for STIB 703 and STIB 710, is given elsewhere (Geigy et al., 1973a, 1975; Brun et al., 1981). STIB 703 was isolated from a patient in Southern Tanzania in 1982 and stabilized after one mouse passage. STIB 710 was isolated from a patient in Daloa (Ivory Coast) in 1982 by the direct transfer of blood parasites into culture.

The initiation and maintenance of the bloodstream form cultures was carried out according to Brun et al. (1981). Parasites were taken from the first raising parasitaemia of mice or *Microtus montanus* (TH-3/78 [031]) infected with trypanosomes from stabilates. Almost confluent feeder layers in 2-cm² wells were inoculated with 10⁵ trypanosomes/well. The test wells contained 15% normal human serum and the control wells 15% inactivated rabbit serum in EMEM. The tests were carried out at least 3 times and the cultures were maintained for at least 10 days with daily medium replacement.

Table 1 shows the test results of 12 different stocks and 4 corresponding clones. STIB 247, STIB 246, STIB 348 (clone of 246) and EATRO 1856 were consistently neutralized by the action of normal human serum. All the other stocks and clones showed continuous growth in the presence of human serum. Stabilate STIB 349 (clone of STIB 324) did not infect one volunteer (Geigy et al., 1975). However, the in vitro test was positive although human serum resistance was only present in a few parasites.

Initially, some stocks (e.g. STIB 229, 347, 350, 324 and STIB 349) contained only a minority of human serum resistant forms whereas most trypanosomes were lysed within a few hours in the presence of human serum. Especially the clones contained only very few resistant forms as indicated by the initial disappearance of the trypanosomes in the culture with human serum and the reappearance of a resistant population after 4–6 days.

The sensitivity of the test was analyzed by adding a single human serum resistant trypanosome of STIB 389 to 10⁵ trypanosomes of STIB 247 (well A). As controls 10⁵ trypanosomes of STIB 247 (well B) and a single trypanosome of STIB 389 (well C) were inoculated into culture wells. All three wells (A, B and C) contained 15% human serum. Well A showed a small growing parasite population after one week which resembled in its growth characteristic the clone growing in well C. The 10⁵ trypanosomes in well B were all lysed within a few hours and no trypanosomes could be detected after 2 weeks. These results show that the test is able to detect single human serum resistant parasites among high numbers of human serum sensitive parasites.

This new test allows the detailed analysis of an eventual fluctuation of human serum resistance in a stock or primary isolate during long term in vitro cultivation. The stability of human serum resistance of different primary isolates before and after tsetse fly transmission is currently being investigated.

Table 1. Human-serum resistance and sensitivity of *Trypanosoma (T.) brucei* stocks and clones

| Isolated from | Place, Country | Year | Stabilate No. | Continuous growth in vitro with | | Man-tested ¹ |
|---------------|---------------------|------|---------------|---------------------------------|---------------------|-------------------------|
| | | | | human serum | inact. rabbit serum | |
| Hartebeest | Serengeti, Tanzania | 1970 | EATRO 1856 | - | + | ND |
| Hartebeest | Serengeti, Tanzania | 1970 | EATRO 1873 | + | + | + |
| Hartebeest | Serengeti, Tanzania | 1971 | STIB 229 | + | + | ND |
| Hartebeest | Serengeti, Tanzania | 1971 | STIB 347* | + | + | ND |
| Hartebeest | Serengeti, Tanzania | 1971 | STIB 246 | - | + | ND |
| Hartebeest | Serengeti, Tanzania | 1971 | STIB 348* | - | + | - |
| Hartebeest | Serengeti, Tanzania | 1971 | STIB 247 | - | + | ND |
| Lion | Serengeti, Tanzania | 1971 | STIB 241 | + | + | + |
| Lion | Serengeti, Tanzania | 1971 | STIB 350* | + | + | + |
| Hyaena | Serengeti, Tanzania | 1971 | STIB 324 | + | + | + |
| Hyaena | Serengeti, Tanzania | 1971 | STIB 349* | + | + | + |
| Man | Homa Bay, Kenya | 1973 | STIB 365 | + | + | - |
| Man | Northern Tanzania | 1980 | STIB 389 | + | + | + |
| Man | Southern Tanzania | 1982 | STIB 703 | + | + | + |
| Man | Ivory Coast | 1978 | TH-3/78 (031) | + | + | + |
| Man | Ivory Coast | 1982 | STIB 710 | + | + | + |

* Clone of the above isolate; ¹ see Geigy et al. (1973b, 1975); ND = Not done

Acknowledgments

We are indebted to M. Schönenberger, C. Kunz and M. Kauffmann for their excellent technical assistance, to Dr. M. Mehltz, Hamburg, for kindly providing us with TH-3/78 (031) and Dr. P. Cattand, Daloa, Ivory Coast, for helping us to isolate STIB 710. These investigations were supported by a Swiss National Science Foundation Grant No. 3.681-0.80.

- Aaronovitch S., Terry R. J.: The trypanolytic factor in normal human serum. *Trans. roy. Soc. trop. Med. Hyg.* 66, 344 (1972).
- Brun R., Jenni L., Schönenberger M., Schell K.-F.: In vitro cultivation of bloodstream forms of *Trypanosoma brucei*, *T. rhodesiense*, and *T. gambiense*. *J. Protozool.* 28 (4), 470-479 (1981).
- Geigy R., Kauffmann M., Jenni L.: Wild mammals as reservoirs for Rhodesian sleeping sickness in the Serengeti 1970/71. *Trans. roy. Soc. trop. Med. Hyg.* 67, 284-286 (1973a).
- Geigy R., Kauffmann M., Mayende J. S. P., Mwambu P. M., Onyango R. J.: Isolation of *Trypanosoma (Trypanozoon) rhodesiense* from game and domestic animals in Musoma District, Tanzania. *Acta trop. (Basel)* 30, 49-56 (1973b).
- Geigy R., Jenni L., Kauffmann M., Onyango R. J., Weiss N.: Identification of *T. brucei*-subgroup strains isolated from game. *Acta trop. (Basel)* 32, 190-205 (1975).
- Hawking F.: The action of human serum upon *Trypanosoma brucei*. *Protozool. Abstr.* 3, 199-206 (1979).
- Hawking F., Ramsden D. B., Whytock S.: The trypanocidal action of human serum and of baboon plasma. *Trans. roy. Soc. trop. Med. Hyg.* 67, 501-516 (1973).
- Laveran A.: De l'action du serum humain sur le Trypanosome du Nagana (*T. brucei*). *C. R. Acad. Sci. (Paris)* 134, 735 (1902).
- Rifkin M. R.: Identification of the trypanocidal factor in normal human serum: high-density lipoprotein. *Proc. nat. Acad. Sci. (Wash.)* 75, 3450-3454 (1978).
- Rickman L. R., Robson J.: The blood incubation infectivity test - a simple test which may serve to distinguish *Trypanosoma brucei* from *T. rhodesiense*. *Bull. WHO* 42, 650-651 (1970).
- Rickman L. R., Robson J.: Some supplementary observations on the blood incubation infectivity test. *Bull. WHO* 46, 403-404 (1972).
- Van Meirvenne N., Janssens P. G., Magnus E.: Further studies on the lytic and neutralizing action of human serum on *T. brucei*. *Ann. Soc. belge Méd. trop.* 53, 49-56 (1973).