

Exergy changes in lakes around the world under pressure from global change

Autor(en): **Silow, Eugene A.**

Objektyp: **Article**

Zeitschrift: **Archives des sciences [2004-ff.]**

Band (Jahr): **65 (2012)**

Heft 1-2

PDF erstellt am: **10.08.2024**

Persistenter Link: <https://doi.org/10.5169/seals-738362>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Exergy changes in lakes around the world under pressure from global change

Eugene A. SILOW¹

Ms. received the 26th September 2012, accepted 17th December 2012

Abstract

Eco-Exergy, an indicator reflecting the state of an ecosystem and the degree of disturbance acting on it is described and a method of calculation is given. The possible changes in lake ecosystems due to Global Change according to the widely accepted "new paradigm" in limnology are listed and discussed. The expected changes of exergy caused by changes in ecosystems are the following: the total pelagic Eco-Exergy content in large lakes must have the tendency to increase, while structural Eco-Exergy of pelagic community of such lakes is to have a demonstrated trend to decrease. While in case of two large lakes (Geneva and Baikal), for which the long-term dynamics of exergy is preliminary analysed, these effects are not observed, the question needs further investigation.

Keywords: Large lakes, pelagic ecosystems, Eco-Exergy, structural Eco-Exergy, Global Change

Introduction


At present, a so called "new paradigm" (Livingstone 2008; Gerten 2008) of limnology, denies the individuality of lake ecosystems and relative constancy of limits, within which parameters of every lake ecosystem fluctuate. This new paradigm proclaims that lake ecosystems respond in concordance, being united by common hydrological cycles and climate, and the unidirectional trend caused by global changes in the environment (climate change and pollution). If this is so, comparable changes in the pelagic community will cause similar trends in such ecosystem state goal functions as exergy and structural exergy (see next paragraph). Taking into account that these goal functions are used for the assessment of the degree of ecosystem perturbation, it is necessary to know which changes of exergy can be caused by general tendencies for lakes according to the "new paradigm", now accepted by many limnologists

(Wilhelm and Adrian 2008; Thackeray et al. 2008; Jöhnk et al. 2008 and many others), if not by everybody among them, believing in Global Change.

Eco-Exergy

The term "exergy" denotes a measure of the quality of energy; as energy is used in any process, it loses quality and decreases in exergy. Basically, the exergy is a measure of the thermodynamic distance of a system from equilibrium with the surrounding environment, and therefore, it is both a quantitative and qualitative measure of the energy (mostly free energy in the context of ecological systems) incorporated into a system. It represents the maximum capacity of energy available to perform useful work as

Fig. 1. Exergy is calculated for the system relatively to the reference environment, Eco-Exergy relatively to the same system at the same temperature and pressure, but as an inorganic solution without life and even organic molecules. (after Jørgensen 2011).



¹ Institute of Biology, UNESCO Chair of Water Resources, Irkutsk State University, Russia; E-mail: eugenasilow@gmail.com

the system proceeds towards equilibrium, with irreversibly increasing its entropy at the expense of exergy (Ludovisi 2009). Taken by itself, the total exergy of an ecosystem is a measure of the difference in entropy between the equilibrium and the actual state (Svirezhev 2000). The exergy of a system in equilibrium with its environment is zero.

We may distinguish between technological exergy and Eco-Exergy: technological exergy uses the environment as reference state and is useful to find the first class energy (work) that a power plant can produce, Eco-Exergy uses as reference state the same ecosystem with the same temperature and pressure but at thermodynamic - chemical equilibrium (Fig. 1). Below, we use the terms exergy and Eco-Exergy as synonyms

The equation for exergy calculation was proposed by S.E. Jørgensen in 1977:

$$Ex = R \cdot T \cdot \sum_{i=0}^N [c_i \cdot \ln(c_i / c_{i,eq}) - (c_i - c_{i,eq})], \quad [J] \quad (1)$$

where Ex - exergy, J; R - gas constant, J Mol⁻¹*K⁻¹; T - temperature, K; c_i - concentration of component i, Mol; c_{i,eq} - concentration of the same component in the state of thermodynamic equilibrium with environment. Mol; n - number of components. This equation is transformed into the following working formula (look for details Silow et al. 2011a, b): total exergy of ecosystem, based on chemical energy of organic matter (biomass) and information, stored in living organisms (recalculating coefficient β), can be calculated as

$$Ex / RT = \sum_{i=1}^N c_i \cdot \beta_i \cdot [\text{g detritus equivalent m}^{-3}] \quad (2)$$

where Ex – exergy, J; R - gas constant, J Mol⁻¹ K⁻¹; T - temperature, K; c_i - biomass concentration, g·m⁻³; and β_i - recalculating coefficient, reflecting quantity of information, stored in organism. This exergy now is often called Eco-Exergy (sometimes - exergy index) to distinguish it from physical or technological exergy (Jørgensen 2011).

Another indicator of ecosystem state, based on Eco-Exergy, was proposed – structural or specific exergy (structural or specific Eco-Exergy). Structural exergy is the exergy related to total biomass (Silow 1998; Xu et al. 1999). Unlike total exergy it does not depend on biomass and it reflects the ability of an ecosystem to accept and utilize the flow of energy from external sources, serving simultaneously as an indicator of ecosystem development, its complexity and the level of evolutionary development of biological species composed in it.

$$Ex_{str} = \left(\sum_{i=1}^N c_i \cdot \beta_i \right) \cdot \left(\sum_{i=1}^N c_i \right)^{-1} \quad [J] \quad (3)$$

We can measure the following aspects of an ecosystem state with Eco-Exergy: 1) the distance from thermodynamic equilibrium, i.e. a general measure of total complexity of ecosystem; 2) structure (biomass and network size) and functions (available information) of ecosystem; 3) ability of ecosystem to survive (expressed via biomass and information of system). Structural exergy reflects: 1) efficiency of energy use by organisms; 2) relative information content of ecosystem and, consequently, the ability of ecosystem to regulate interactions between organisms or groups of organisms.

Corresponding β coefficients (recalculating coefficients, reflecting quantity of information, stored in organism and depending on the number of informative genes and number of cell types in the organism) were calculated for many systematic groups and are published regularly (Jørgensen et al. 2005; Jørgensen 2006; Jørgensen and Faith 2011; Jørgensen 2011). These coefficients reflect relative complexity of organisms (simpler organisms have lower β values). New β values are added every year (Table 1).

Changes in Lakes Caused by Global Warming

Let's observe the possible consequences of climate change for lake ecosystems basing on the modern limnology paradigm on structure and functioning of aquatic ecosystems (Hutchinson 1957, 1967; Wetzel 2001; Kalff 2002; Schwoerbel and Brendelberger 2005).

If the surface temperature of the air rises, so the temperature of the surface water increases too. Long-term increases of air temperature will cause an increase in water temperature of the surface layer of a lake. If persistent the temperature of the whole epilimnion will increase. There are three phenomena connected with these processes: (i) strengthening of water column stability, (ii) prolongation of stratification, and (iii) increase of epilimnion volume. These phenomena were observed in Scandinavian lakes (Pettersson et al. 2003; Jackson et al. 2007), Lake Zürich, Switzerland (Livingstone 2003), Lake Constance, Switzerland and Germany (Straile et al. 2003), other European lakes (George et al. 2000; Livingstone Dokulil 2001; Elliott et al. 2005; Blenckner et al. 2007), in Great African Lakes (O'Reilly et al. 2003), in Great Laurentian Lakes (King et al. 1997), and in Experimental Lake Area in Ontario, Canada (Schindler et al. 1996; Schindler 2001; Findlay et al. 2004), lake Washington (Arhonditsis et al. 2004), lakes of Wisconsin, USA (Magnuson et al. 1990), South America (Baiguin and Marinone 1995; Solo 2002), Antarctica (Quayle et al. 2002).

Climatic changes must be reflected by hydrochemical parameters also, as climate influences both processes

Table 1. Exergy/Biomass Conversion factors for different groups of organisms (after Silow & Mokry 2010).

Group	Exergy conversion factor, β	Group	Exergy conversion factor, β
Minimal cell	5.8	Brachiopoda	109
Bacteria	8.5–12	Seedless vascular plants	158
Archaea	13.8	Rotifera	163
Yeasts	18	Insecta	167–446
Alga	15–298	Chironomida	300
Cyanobacteria	15	Moss	174
Dynophyta	18	Crustaceans	230–300
Green microalgae	20	Cladocera	232
Diatoms	66	Copepoda	240
Macrophyta (alga)	67–298	Amphipoda	290
Rhodophyta	92	Mollusca	297–450
Protozoa	31–97	Bivalves	297
Amoeba	38	Gastropoda	312–450
Gastrotricha	97	Gymnosperm	314
Fungi	61	Macrophytes (Phanerogam)	356–520
Nemertina	76	Flowering plants	393–543
Worms	91–133	Fish	499–800
Cnidaria	91	Amphibia	688
Plathelminthes	120	Reptilia	833
Oligochaeta	130	Aves	980
Nematoda	133	Mammalia	2127
Sponges	98	Homo sapiens	2173

in the waterbody and watershed. So, we can expect a decrease of oxygen content and an increase of the nutrients concentration in the hypolimnion, due to strengthening and lengthening of lake stratification. Actually lowering of oxygen content is already observed in some lakes (Straile et al. 2003; Livingstone 2003; Jankowski et al. 2006; Peeters et al. 2007; Wilhelm and Adrian 2008). An increase in nutrients, in particularly phosphorus also has been observed in the hypolimnion of many lakes, due to internal processes (Pettersson et al. 2003; Wilhelm and Adrian, 2008), and an increase of nutrient loading from the watershed (Yoshioka et al. 2002; Prowse et al. 2006).

These serious changes in hydrology and hydrochemistry of water body must unavoidably be reflected in the biota. We can expect an intensive development of phytoplankton, changes in its composition, an increase of microbial activity, and consequently changes in zooplankton composition and fish population.

In deep oligo- and monomictic lakes the decrease of spring mixing causes lowering of silicates and phosphates concentration prior to phytoplankton development (Goldmann et al. 1989; Straile et al. 2003; Salmaso 2005). The consumption of nutrients by phytoplankton further decreases the concentration of dissolved nutrients. Consequently, earlier phytoplankton development due to warming (Adrian et al.

1999; Peeters et al. 2007) leads to earlier lowering of silicates, nitrates and phosphates (Anneville et al. 2002; Weyhenmeyer et al. 2007; Thackeray et al. 2008).

From the other hand, long and stable summer stratification causes the growth of cyanobacteria (Jöhnk et al. 2008) and, consequently / potentially (depending on the cyanobacterial taxa) nitrogen fixation. High temperatures can stimulate the growth of other algae (Elliott et al. 2005; Huber et al. 2008). So, significant increase of primary production in Arctic lakes is demonstrated for XXth century (Michelutti et al. 2005). The rise of temperature was shown experimentally (Rae and Vincent, 1998) to stimulate first the smallest algae (0,2-2,0 μm), than intermediate sized ones (2,0-20 μm), and then large ones (20-200 μm). The earlier ice-off and reinforced input of nutrients from the watershed favours the mass development of small forms of diatoms (Bradbury et al. 1994; Mackay 2007).

The growth of small-sized rapidly reproducing algae causes on the one hand, better feeding conditions for infusoria, flagellates, rotifers and cladocerans (filter feeding zooplankton), and on the another hand, worse feeding conditions for copepods and other catchers. The transfer of energy to higher trophic levels can be decreased by 40-65% (Moline et al., 2004). Both, direct influences of higher temperatures as well as changes in feeding conditions affect the shifts in zooplankton composition (Patalas 1990; Patalas and Salki, 1992). Warming causes a shift in the cladocerans: copepods ratio, as cladocerans react faster to an increase in food availability and to temperature (Straile et al. 2005). With an increase in temperature, the development of cladocerans influences greatly the whole ecosystem (Mayer 1997; Elser and Urabe 1999; Gillooly and Dodson 2000; Carpenter et al. 2001; Sommer and Sommer 2006). The whole structure of the ecosystem changes due to trophic chains restructuring (Winder and Schindler 2004, Yoshioka et al. 2002).

Expected and Observed Changes of Eco-Exergy in Lakes

Now we can transform the changes in lake ecosystems discussed above into changes of Eco-Exergy we may expect, using Table 1. We can predict the

increase of total Eco-Exergy, as the primary production is predicted to increase with warming, and total biomass will be increasing, in a way similar to eutrophication.

Structural Eco-Exergy, vice versa, must decline if there are changes as listed above. Actually, for phytoplankton the preferential development of cyanobacteria and other small sized fast reproducing forms algae is shown under global warming. They have lower conversion factors, than large-sized slowly growing phytoplankton species. It is the first driver of structural Eco-Exergy decrease. An increase of bacterial biomass is expected. This is a second driver of structural Eco-Exergy decrease. Changes in zooplankton composition can be the third driver of structural Eco-Exergy decrease, as the coefficient for cladocerans is lower than that for copepods. In the case of extreme events with hypolimnion isolation and anoxic conditions, fish kills will be followed by dramatic decrease of structural Eco-Exergy.

Nevertheless, if we take lakes with calculated long-term dynamics of exergy and eco-exergy for their pelagic ecosystems (Silow et al. 2011c), e.g. lake Geneva (1974-2005) and lake Baikal (1950-1999), we see that Eco-Exergy in these lakes does not (yet?) behave in complete accordance with our predictions. The total Eco-Exergy in lake Baikal tends to increase, while in Geneva lake it remains approximately the same. Structural Eco-Exergy in Geneva lake is decreasing, while in lake Baikal it has no trend, neither a decrease nor an increase. Long-term dynamics

of the lake Geneva is more determined by the program of lake restoration, including re-oligotrophication and pollution prevention, which caused a more intensive grazing of fishes on zooplankton and consequent phytoplankton development, resulted in a structural Eco-Exergy decrease. As for lake Baikal, though some trends in its ecosystem can be explained by climate changes, this giant lake remains too conservative and too inert to follow general tendencies rapidly.

Conclusion

So, if the hypothesis proposed in the “new paradigm” in limnology is true, the total pelagic Eco-Exergy content in large lakes must have a tendency to increase, while structural Eco-Exergy of pelagic community of such lakes would tend to decrease. Yet, in the case of two big lakes (Geneva and Baikal), for which the long-term dynamics of exergy are preliminary analysed, these effects are not observed, hence the question needs further investigation.

Acknowledgment

Author is pleased to acknowledge the Federal Targeted Program “Scientific and Pedagogical Staff for Innovative Russia” for 2009-2013, that supported this research with contract 14.B37.21.1252.

Bibliographie

- **ADRIAN R, WALZ N, HINTZE T, HOEG S, RUSCHE R.** 1999. Effects of ice duration on the plankton succession during spring in a shallow polymictic lake. *Freshw. Biol.*, 1999, 41, 621-623.
- **ANNEVILLE O, SOUSSI S, IBAÑEZ F, GINOT V, DRUART J-C, ANGELI N.** 2002. Temporal mapping of phytoplankton assemblages in Lake Geneva: annual and interannual changes in their patterns of succession. *Limnol. Oceanogr.*, 47, 1355-1366.
- **ARHONDITSIS GB, BRETT MT, DE GASPERI CL, SCHINDLER DE.** 2004. Effects of climatic variability on the thermal properties of Lake Washington. *Limnol. Oceanogr.*, 49(1), 256-270.
- **BAIGÚN C, MARINONE MC.** 1995. Cold-temperate lakes of South America: do they fit northern hemisphere models? *Arch. Hydrobiol.*, 135 (1), 23-51.
- **BLENCKNER T, ADRIAN R, LIVINGSTONE DM, JENNINGS E, WEYHENMEYER GA, GEORGE DG, JANKOWSKI T, JARVINEN M, AONGHUSA CN, NOGES T, STRAILE D, TEUBNER K.** 2007. Large-scale climatic signatures in lakes across Europe: a meta-analysis. *Glob. Change Biol.*, 2007, 13, 1314-1326.
- **BRADBURY JP, BEZRUKOVA YV, CHERNYAIEVA GP, COLMAN SM, KHURSEVICH G, KING JW, LIKOSHWAY YV.** 1994. A synthesis of post-glacial diatom records from Lake Baikal. *Journal of Paleolimnology*, 10, 213-252.
- **CARPENTER SR, COLE JJ, HODGSON JR et al.** 2001. Trophic cascades, nutrients, and lake productivity: whole-lake experiments. *Ecological Monographs*, 71, 163-186.
- **ELLIOTT JA, THACKERAY SJ, HUNTINGFORD C et al.** 2005. Combining a regional climate model with a phytoplankton community model to predict future changes in phytoplankton in lakes. *Freshw. Biol.*, 50, 1404-1411.
- **ELSER JJ, URABE J.** 1999. The stoichiometry of consumer-driven nutrient recycling: theory, observations, and consequences. *Ecology*, 80, 735-751.

- **FINDLAY DL, KASIAN SEM, STANTON MP, BEATY K, LYNG M.** 2001. Climatic influences on algal populations of boreal forest lakes in the Experimental Lakes Area. *Limnol. Oceanogr.*, 2001, 46(7). 1784-1793.
- **GEORGE DG, TALLING JF, RIGG E.** 2000. Factors influencing the temporal coherence of five lakes in the English Lake District. *Freshw. Biol.*, 43, 449-461.
- **GERTEN D.** 2008. Climatic change, aquatic science, multiple shifts in paradigms. *Internat. Rev. Hydrobiol.*, 93, 397-403.
- **GILLOOLY JF, DODSON SI.** 2000. Latitudinal patterns in the size distribution and seasonal dynamics of new world, freshwater cladocerans. *Limnol. Oceanogr.*, 45, 22-30.
- **GOLDMAN CR, JASSBY A, POWELL T.** 1989. Interannual fluctuations in primary production: Meteorological forcing at two subalpine lakes. *Limnol. Oceanogr.*, 34, 310-323.
- **HUBER V, ADRIAN R, GERTEN D.** 2008. Phytoplankton response to climate warming modified by trophic state. *Limnol. Oceanogr.*, 53(1), 1-13.
- **HUTCHINSON GE.** 1957. *A Treatise on Limnology*. John Wiley & Sons, New York. Vol. 1. Geography, Physics and Chemistry.
- **HUTCHINSON GE.** 1967. *A Treatise on Limnology*. John Wiley & Sons, New York. Vol. 2. Introduction to Lake Biology and Limnoplankton.
- **JACKSON LJ, LAURIDSEN TL, SONDERGAARD M, JEPPESEN E.** 2007. A comparison of shallow Danish and Canadian lakes and implications of climate change. *Freshw. Biol.*, 52, 1782-1792.
- **JANKOWSKI T, LIVINGSTONE DM, BUHRER H, FORSTER R, NIEDERHAUSER P.** 2006. Consequences of the 2003 European heat wave for lake temperature profiles, thermal stability, and hypolimnetic oxygen depletion: Implications for a warmer world. *Limnol. Oceanogr.*, 51, 815-819.
- **JÖHNK KD, HUISMAN J, SHARPLES J, SOMMEIJER B, VISSER PM, STROOM JM.** 2008. Summer heatwaves promote blooms of harmful cyanobacteria. *Glob. Change Biol.*, 14, 495-512.
- **JØRGENSEN SE.** 2006. Application of holistic thermodynamic indicators. *Ecological Indicators*, 2006, Vol. 6, pp. 24-29.
- **JØRGENSEN SE.** 2011. *An Introduction to Ecosystem Theory*, Taylor & Francis Group/CRC Press, Boca Raton.
- **JØRGENSEN SE, LADEGAARD N, DEBELJAK M, MARQUES JC.** 2005. Calculations of exergy for organisms. *Ecological Modelling*, Vol. 185, pp. 165-175
- **JØRGENSEN SE, FATH BD.** 2011. *Fundamentals of Ecological Modelling: Applications in Environmental Management and Research*, Elsevier, 4th ed., Amsterdam, The Netherlands.
- **KALFF J.** 2002. *Limnology: Inland Water Ecosystems*. Prentice Hall, New Jersey.
- **KING JR, SHUTER BJ, ZIMMERMAN AP.** 1997. The response of the thermal stratification of South Bay (Lake Huron) to climatic variability. *Can. J. Fish. Aquat. Sci.*, 54, 1873-1882.
- **LIVINGSTONE DM.** 2003. Impact of secular climate change on the thermal structure of a large temperate central European lake. *Climatic Change*, 57, 205-225.
- **LIVINGSTONE DM.** 2008. A change of climate provokes a change of paradigm: taking leave of two tacit assumptions about physical lake forcing. *Internat. rev. Hydrobiol.*, 93, 404-414.
- **LIVINGSTONE DM, DOKULIL MT.** 2001. Eighty years of spatially coherent Austrian lake surface temperatures and their relationship to regional air temperature and the North Atlantic Oscillation. *Limnol. Oceanogr.*, 46, 1058-1066.
- **LUDOVISI A.** 2009. Exergy vs information in ecological successions: Interpreting community changes by a classical thermodynamic approach. *Ecological Modelling*. 220, 1566-1577.
- **MACKAY AW.** 2007. The paleoclimatology of Lake Baikal: A diatom synthesis and perspectives. *Earth-Science Reviews*, 82, 181-215.
- **MAGNUSON JJ, BENSON BJ, KRATZ TK.** 1990. Temporal coherence in the limnology of a suite of lakes in Wisconsin, U.S.A. *Freshw. Biol.*, 23, 145-159.
- **MAYER CM.** 1997. The relationship between prey selectivity and growth and survival in a larval fish. *Can. Journ. Fish. Aquat. Sci.*, 54, 1504-1512.
- **MICHELUTTI N, WOLFE AP, VINEBROOKE RD, RIVARD B, BRINER JP.** 2005. Recent productivity increases in Arctic lakes. *Geophysical Research Letters*, 32, L19715, doi:10.1029/2005GL023693.
- **MOLINE MA, CLAUSTRE H, FRAZER TK et al.** 2004. Alteration of the food web along the Antarctic Peninsula in response to a regional warming trend. *Glob. Change Biol.*, 10, 1973-1980.
- **O'REILLY CM, ALIN SR, PLISNIER PD, COHEN AS, MCKEE BA.** 2003. Climate change decreases aquatic ecosystem productivity of Lake Tanganyika, Africa. *Nature*, 424, 766-768.
- **PATALAS K, SALKI A.** 1992. Crustacean Plankton in lake Winnipeg: variation in space and time as a function of lake morphology, geology, and climate. *Can. Journ. Fish. Aquat. Sci.*, 49, 1035-1059.
- **PATALAS K.** 1990. Patterns in zooplankton distribution and their causes in North American great lakes. *In: Tilzer M.M., Serruya C. (eds.) Large Lakes: Ecological Structure and Function*. Springer-Verlag, Berlin, pp. 440-458.
- **PEETERS F, STRAILE D, LORKE A, LIVINGSTONE DM.** 2007. Earlier onset of the spring phytoplankton bloom in lakes of the temperate zone in a warmer climate. *Glob. Change Biol.*, 13, 1898-1909.
- **PETTERSSON K, GRUST K, WEYHENMEYER G, BLENCKNER T.** 2003. Seasonality of chlorophyll and nutrients in Lake Erken - effects of weather conditions. *Hydrobiologia*, 506, 75-81.
- **PRAWSE TD, WRONA FJ, REIST JD, GIBSON JJ, HOBBIJE JE, LEVESQUE LMJ, VINCENT WF.** 2006. Climate change effects on hydroecology of Arctic freshwater ecosystem. *Ambio*, 35, 347-358.
- **RAE R, VINCENT WF.** 1998. Phytoplankton production in subarctic lake and river ecosystems: development of a photosynthesis-temperature-irradiance model. *J. Plankton Research*, 20, 1293-1312.

- QUAYLE WC, PECK LS, PEAT H, ELLIS-EVANS JC, HARRIGAN PR. 2002. Extreme responses to climate change. *Science*, 295, 645.
- SALMASO N. 2005. Effects of climatic fluctuations and vertical mixing on the interannual trophic variability of Lake Garda, Italy. *Limnol. Oceanogr.*, 50, 553-565.
- SCHINDLER DW, CURTIS PJ, PARKER BR, STANTON MP. 1996. Consequences of climate warming and lake acidification for UV-B penetration in North American boreal lakes. *Nature*, 379, 705-708.
- SCHINDLER DW. 2001. The cumulative effects of climate warming and other human stresses on Canadian freshwaters in the new millennium. *Can. J. Fish. Aquat. Sci.*, 58, 18-29.
- SCHWOERBEL J, BRENDENBERGER H. 2005. Einführung in die Limnologie. 9 Auflage. Elsevier, München.
- SILOW EA. 1998. The changes of ecosystem goal functions in stressed aquatic communities. *Journal of Lake Science*, 10, 421-435.
- SILOW EA, MOKRY AV. 2010. Exergy as a Tool for Ecosystem Health Assessment. *Entropy*, 12, 902-925.
- SILOW EA, MOKRY AV, JØRGENSEN SE. 2011a. Some Applications of Thermodynamics for Ecological Systems. In: J.C. Moreno-Pirajan (ed.). *Thermodynamics - Interaction Studies - Solids, Liquids and Gases*, InTech, Vienna, pp. 319-342.
- SILOW EA, MOKRY AV, JØRGENSEN SE. 2011b. Eco-Exergy use for ecosystem health assessment. *International Journey of Exergy*, Paris Ouest University, Paris, 22 p. Available from http://leme.u-paris10.fr/exergy/files/17_06_11/SilowA.pdf
- SILOW EA, ANNEVILLE O, MONTUELLE B, MOKRY AV, XU F-L. 2011c. Case studies of Eco-Exergy use for ecosystem health assessment. *International Journey of Exergy*, Paris Ouest University, Paris, 18 p. Available from http://leme.u-paris10.fr/exergy/files/17_06_11/SilowB.pdf
- SOMMER U, SOMMER F. 2006. Cladocerans versus copepods: the cause of contrasting top-down controls on freshwater and marine phytoplankton. *Oecologia*, 147, 183-194.
- SOTO D. 2002. Oligotrophic patterns in southern Chilean lakes: The relevance of nutrients and mixing depth. *Revista Chilena de Historia Natural*, 75 (2), 377-393.
- STRAILE D, JOEHNK K, ROSSKNECHT H. 2003. Complex effects of winter warming on the physico-chemical characteristics of a deep lake. *Limnol. Oceanogr.*, 48, 1432-1438.
- STRAILE D. 2005. Food webs in lakes – seasonal dynamics and the impact of climate variability. In: Belgrano A., Scharler U., Dunne J., Ulanowicz R.E. (eds.) *Aquatic food webs: an ecosystem approach*, Oxford University Press, Oxford, pp. 41-50.
- SVIREZHEV YM. 2000. Thermodynamics and ecology. *Ecological Modelling*, 132, 11-22.
- THACKERAY SJ, JONES ID, MABERLY SC. 2008. Long-term change in the phenology of spring phytoplankton: species-specific responses to nutrient enrichment and climatic change. *Journal of Ecology*, 96, 523-535.
- WEYHENMEYER GA, JEPPESEN E, ADRIAN R, ARVOLA L, BLECKNER T, JANKOWSKI T, JENNINGS E, NOGES P, NOGES T, STRAILE D. 2007. Nitrate-depleted conditions on the increase in shallow northern European lakes. *Limnol. Oceanogr.*, 52, 1346-1353.
- WETZEL RG. 2001. *Limnology: Lake and River Ecosystems*. 3d ed. Academic Press, London / Sydney / Tokyo.
- WILHELM S, ADRIAN R. 2008. Impact of summer warming on the thermal characteristics of a polymictic lake and consequences for oxygen, nutrients and phytoplankton. *Freshw. Biol.*, 53, 226-237.
- WINDER M, SCHINDLER DE. 2004. Climate change uncouples trophic interactions in an aquatic ecosystems. *Ecology*, 85, 2100-2106.
- XU FL, JØRGENSEN SE, TAO S. 1999. Ecological indicators for assessing freshwater ecosystem health. *Ecological Modelling*, 116, 77-106.
- YOSHIOKA T, UEDA S, KHODZHER TV, BASHENKHAeva N, KOROVYAKOVA IV, SOROKOVIKOVA LM, GORBUNOVA I. 2002. Distribution of dissolved organic carbon in Lake Baikal and its watershed. *Limnology*, 3, 159-168.