

Behaviour, morphology and molecular characterization of a *Monophadnus* sawfly species (Hymenoptera: Tenthredinidae) feeding on *Helleborus* spp. in Western Switzerland

Autor(en): **Malacari, Lucas / Pétremand, Gaël / Rochefort, Sophie**

Objektyp: **Article**

Zeitschrift: **Archives des sciences [2004-ff.]**

Band (Jahr): **69 (2017)**

Heft 2

PDF erstellt am: **13.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-825731>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Behaviour, morphology and molecular characterization of a *Monophadnus* sawfly species (Hymenoptera: Tenthredinidae) feeding on *Helleborus* spp. in Western Switzerland

Lucas MALACARI¹, Gaël PÉTREMAND¹, Sophie ROCHEFORT¹, Bastien COCHARD¹, Gautier CALMIN² et François LEFORT^{1*}

Ms. received the 14th February 2017, accepted the 9th June 2017

Abstract

Recently, an undescribed sawfly species (Hymenoptera : Symphyta) has been causing serious damage to the *Helleborus* collections of the Conservatory and Botanical Garden of Geneva, destroying up to 80% of some species populations. These sawflies were very aggressive towards *Helleborus* and were shown to be able to feed on 14 *Helleborus* species and subspecies. This study aimed to identify this oligophagous sawfly species, which was morphologically identified as belonging to the genus *Monophadnus* (Hymenoptera : Tenthredinidae). The morphological description was supplemented with molecular identification using several genes, which showed this species to be genetically different from *M. monticola*, often considered as a synonym of *M. latus*, and of any species for which molecular data are available. Therefore, it is likely that this sawfly species could be *Monophadnus latus* sensu Lacourt 1999, which could be confirmed in a next future, at the occasion of a revision of the genus *Monophadnus* supported by genetic identifications.

Keywords : Molecular identification, Symphyta, *Monophadnus*, Switzerland, botanical garden, hellebores, Geneva, horticulture

Résumé

Comportement, morphologie et caractérisation moléculaire d'une espèce symphyte de *Monophadnus* (Hymenoptera: Tenthredinidae) se nourrissant sur diverses espèces d'*Helleborus* en Suisse Occidentale. – Depuis quelques années, une espèce non décrite de mouche à scie (Hymenoptera : Symphyta) cause d'importants dégâts aux collections d'hellébore des Conservatoire et Jardin botaniques de Genève, détruisant jusqu'à 80% des plantes de certaines espèces. Ces mouches à scie, très voraces, ont été observées se nourrissant sur 14 espèces et sous-espèces d'hellébore. Cette étude avait pour objectif d'identifier cette espèce de mouche à scie laquelle a été morphologiquement identifiée comme appartenant au genre *Monophadnus* (Hymenoptera : Tenthredinidae). La description morphologique a été complétée par une identification moléculaire à plusieurs gènes, qui montre que cette espèce est génétiquement différente de

¹ Research Institute Land Nature and Environment, hepia, HES-SO University of Applied Sciences and Arts Western Switzerland, Route de Presinge 150, 1254 Jussy, Geneva, Switzerland

² Faculty of Engineering and Architecture, HES-SO University of Applied Sciences and Arts Western Switzerland, Rue de la Jeunesse 1, 2800 Delémont, Switzerland

* Corresponding author: francois.lefort@hesge.ch

M. monticola, souvent considérée comme un synonyme de *M. latus*, et de toutes les espèces pour lesquelles une description moléculaire est disponible. Il est probable que cette espèce soit *M. latus* sensu Lacourt 1999, ce qui pourrait être confirmé dans le futur par une révision du genre *Monophadnus* complétée par des identifications génétiques.

Mots-clés : Identification moléculaire, Symphytes, *Monophadnus*, Suisse, Jardin botanique, hellébères, Genève, horticulture

Introduction

More than 8000 Symphyta (Hymenoptera) species distributed worldwide have been described, including important pest insects, which are major defoliators in the Northern hemisphere, in natural areas as well as in horticultural and agricultural production sites (Boevé 2004; Taeger et al. 2010). Symphyta include more than 1000 species, for which the phylogeny has been recently refined (Prous et al. 2014; Welz & Vilhelmsen 2014; Vilhelmsen 2015), and new sawflies species are continuously being discovered and described (Haris 2015; Niu et al. 2015; Looney et al. 2016).

In the literature, several authors have reported *Monophadnus* spp. larvae as feeding on *Helleborus* spp., especially on *H. niger* L., *H. foetidus* L. and *H. viridis* L. (Fenili 1965; Jahn & Schedl 1992; Lacourt 1994). The *Helleborus* collections of the Conservatory and Botanical Garden of Geneva (CBG) have been particularly infested by sawflies for the last five years, with damage reaching 80% of individuals for some *Helleborus* species. For the purpose of developing biological control trials against this pest, it appeared important to first try to secure its identity. Additionally, the *Helleborus* collection found at the CBG provided an opportunity to observe this sawfly on 14 *Helleborus* species and subspecies.

Materials and Methods

Survey of Plant Species Hosting Sawflies in the CBG Collections

Observations were carried out in Spring 2015 in all *Helleborus* collections and isolated plants of the CBG in order to evaluate the range of species attacked by

this sawfly. Specimens, both adults and larvae, were collected on *Helleborus* plants at the CBG and deposited at the Museum of Natural History of Geneva under voucher numbers MHNG ENTO 9805 to 9808.

Morphological Identification

The morphological description of the adult was initially achieved by using the determination keys for Hymenoptera Tenthredoidea by Berland (1947).

Molecular Identification

Samples of 3-4 larvae at the fourth stage were ground in liquid nitrogen with a mortar and pestle. Larvae were sampled and submitted to DNA extraction. Total nucleic acids were extracted from the resulting powder using the DNeasy Blood & Tissue kit (Qiagen AG, Switzerland), according to the protocol «Purification of total DNA from insects». Genomic DNA quality was checked and assayed with a Nanodrop ND-1000 spectrophotometer (WITEC AG, Switzerland). PCR amplifications were carried out for the 18S and 28SrDNA nuclear genes and the cytochrome oxidase subunit 1 mitochondrial gene (COI) with specific primers, whose sequences and sources are given in Table 1. PCR reactions were carried out in 50 µl final volume containing 50 ng DNA, using 1.25 u of MyTaq™ HS DNA Polymerase (Bioline, UK) in 1x MyTaq™ Reaction Buffer. PCR programs used for the amplification of the three genes are given in Table 2. Touch-down programs were used for 28S and COI genes. PCR products were checked for quality by gel electrophoresis, then purified with the Wizard® SV Gel and PCR Clean-up system (Promega AG, Switzerland) and submitted to Sanger sequencing. Result-

Table 1. Primers used for the phylogenetic study.

Genes	Primer	Sens	Sequence (5' - 3')	Source
18S	2880	F	CTG GTT GAT CCT GCC AGT AG	Von Dohlen & Moran, 1995
	Br	R	CCG CGG CTG CTG GCA CCA GA	Von Dohlen & Moran, 1995
28S	S3660	F	GAG AGT TMA ASA GTA CGT GAA AC	Downton & Austin, 1998
	A335	R	TCG GAR GGA ACC AGC TAC TA	Whiting et al., 1997
COI	LepF1	F	ATT CAA CCA ATC ATA AAG ATA TTG G	Hebert et al., 2004
	LepR1	R	TAA ACT TCT GGA TGT CCA AAA AAT CA	Hebert et al., 2004



Fig. 1. Damage caused by *Monophadnus* sawfly larvae on *Helleborus foetidus*.

ing sequences were edited with FinchTV and registered in NCBI GenBank (Benson et al. 2013), where they were compared to deposited sequences with the Basic Local Alignment Search Tool (BLAST) algorithm (Altschul et al. 1990).

Outgroup sequences for the same genes were recovered from GenBank. Phylogenetic inference was based on a multigene dataset (28S and COI). Sequence alignment was constructed using Bioedit 7.1.9 (Hall 1999). Phylogeny was generated using maximum likelihood. The best-fitting nucleotide substitution model was selected via the Akaike information criterion (AIC), as implemented in the PhyML-SMS online execution program (<http://www.atgc-montpellier.fr/phyml-sms/>). The Gen-

eral Time Reversible model, with gamma-distributed among-site rate heterogeneity and invariant sites (GTR+G+I), was recovered as most appropriate. ML analysis was inferred using PhyML online execution platform (Guindon et al. 2010), applying model suggested by PhyML-SMS. Node support was quantified with 1000 rapid bootstrap replicates. The tree was visualised using Fig-Tree 1.4.2 (Rambaut 2009).

Results

Survey of Host Plant Species in CBG collection

The unknown sawfly species was observed feeding on all 14 *Helleborus* species and subspecies found at the CBG: *H. argutifolius* Viv., *H. cyclophyllus* Boiss., *H. dumetorum* ssp. *atrorubens* (Waldst. & Kit.) Merxm. & Podlech, *H. foetidus*, *H. lividus* Aiton, *H. multifidus* ssp. *bocconeii* (Tenore) Mathew, *H. niger*, *H. odoratus* Waldst. & Kit. ex Willd., *H. orientalis* Lam., *H. orientalis* ssp. *abchasicus* (A. Braun) B. Mathew, *H. purpurascens* Waldst. & Kit., *H. torquatus* Archer-Hind, *H. viridis* and *H. viridis* ssp. *occidentalis* (Reut.) Schiffn.. The damages were most important on *H. foetidus* (Fig. 1), while *H. niger* supported the least significant damages.

Morphological description

The morphological description was carried out with the descriptors of the determination key established by Berland (1947). Following Berland (1947), two characters enable this *Monophadnus* species to be identified as *Monophadnus longicornis* Har-

Table 2. PCR programs used to amplify the 3 genes used for the phylogenetic study.

Gene		Initial denaturation	Denaturation	30-35 cycles		
				Hybridation	Amplification	Final extension
18S	Temperature [°C]	94	94	55	72	72
	Time	4 min	1 min	50 s	45 s	7 min
28S	Temperature [°C]	95	95	49 - 61	72	72
	Time	4 min	1 min	50 s	50 s	7 min
COI	Temperature [°C]	95	95	48 - 60	72	72
	Time	5 min	1 min	1 min	1 min	10 min

Fig. 1837: 1) simple tarsal claws and 2) third segment of antennae only one and a half length of fourth segment (Fig. 2). Other morphological characters of adults of this *Monophadnus* species are: body length 6.5-8.2 mm; antennae of nine articles and shorter than the abdomen; legs almost entirely black, but upper half of tibiae yellow-brown; dark brown to black veins and stigma. Our observations on the CBG sawfly are congruent with the morphological descriptions of *M. longicornis* given by Berland (1947), Fenili (1965) and Jahn and Schedl (1992). The general aspects of the adult and larva are shown in Figure 3. Given that the key of Berland (1947) is obsolete for identification purposes and that the *Monophadnus* genus includes many undescribed or wrongly named species (Lacourt 1999), our morphological and behavioural observations allow us to identify the CBG specimens as an undescribed species of the genus *Monophadnus*, close to *Monophadnus latus sensu* Lacourt 1999, labelled *Monophadnus* sp. MHNG ENTO 9805.

Molecular Identification

The DNA sequences obtained have been registered in NCBI GenBank under accessions numbers KU302717-KU302718 for the 18S rDNA gene, KU302719-

KU302720 for the 28S rDNA gene, and KU302721-KU302722 for the COI gene. Genbank searches showed that genetic information on *Monophadnus* was very scarce. There are, for instance, no 18S rDNA sequences available, except the ones produced in this study for *Monophadnus* sp. MHNG ENTO 9805, and very few sequences registered for both 28S rDNA and COI genes. Sequence comparison allowed to confirm that the sequences for *Monophadnus* sp. MHNG ENTO 9805 were the first ones registered for this undescribed species. A phylogram built with the concatenated genes 28S and COI shows that *Monophadnus* sp. MHNG ENTO 9805 was very different from *M. monticola* Hartig 1837 (Fig. 4). A second phylogram, built only with COI sequences, clearly showed that these two species differ from each other and are also different from other *Monophadnus* species for which COI sequences are available (Fig. 5). A specimen labeled *Monophadnus* sp. A GM-2013 is the closest genotype to *Monophadnus* sp. MHNG ENTO 9805. Cited by Boevé et al. (2013), this specimen from an unidentified species was found on *H. foetidus* in Delémont, Northern Switzerland. The *Monophadnus* sp. B GM-2013 (Boevé et al. 2013) was sampled in Meride (Ticino), South East Switzerland, as was the *Monophadnus* sp. DEIGISHym11547 sampled on *H. viridis*. Though close to *Monophadnus* sp. MHNG ENTO 9805, at 99% similarity, these




Fig.2. Morphological criteria used for identification; third antennal segment only one and a half length of fourth segment (2A), simple tarsal claws (2B).




Fig.3. General aspects of the adult (3A) and the larva (3B).




Fig. 4. ML tree topology derived from multigene analysis (28S and COI). Bootstrap support values are given at nodes. Only bootstrap values > 850 are shown. Blanks indicate values that were below the corresponding thresholds. Bar = 0.008 substitution positions.

three specimens, are nevertheless distinct genotypes. These first registered DNA sequences for three genes add to the morphological description of *Monophadnus* sp. MHNG ENTO 9805 and will help to place it in the phylogeny of the genus in the future.

Discussion and Conclusion

This study identified a *Monophadnus* species as the species causing damage to the *Helleborus* collections of the CBG. However, morphology as well as genetic data do not allow us to name this species, which could likely be *M. latus sensu* Lacourt 1999. Although the species described here was easily differentiated from the four other European species *Monophadnus alpicola* Benson 1954, *M. monticola*, *M. pallescens*

Gmelin 1790, and *M. spinolae* Klug 1816, it has not been possible to name this species because of historical confusions, using three different names *Monophadnus latus*, *M. longicornis* and *M. monticola*, with no name being correct. The key of Berland does not allow *Monophadnus latus sensu* Lacourt 1999, to be distinguished from *M. monticola* and does not include *M. alpicola*. *M. monticola* is known to feed on *Ranunculus* spp. and *M. alpicola* on *Pulsatilla* spp., and for this reason, these specimens have not been considered in this study focusing on *Helleborus* spp. There is no morphological characters that would help distinguish whether it is *M. latus sensu* Lacourt 1999, or another undescribed species, and it is not clear, from the literature, if the aforesaid synonyms correspond to the same species. There is however a consensus that the *Monophadnus* species feeding




Fig. 5. ML tree topology for the COI gene. Bootstrap support values are given at nodes. Only bootstrap values > 850 are shown. Blanks indicate values that were below the corresponding thresholds. Bar = 0.05 substitution positions.

on *Helleborus* spp. could be named *M. latus sensu* Lacourt 1999. The species present in CBG, feeds on a large range of *Helleborus* spp. and seems to be genetically different from all referenced specimens. Concerning the behavior of *M. latus sensu* Lacourt 1999, females lay their eggs in the mesophyll of the leaves and petals of hellebores and are reported to feed only on this genus: they have been found feeding only on *H. viridis*, *H. foetidus* and *H. niger* (Fenili 1965; Jahn & Schedl 1992; Lacourt 1994). Concerning the host plants, Fenili (1965) stated that *M. longicornis* attacked *H. viridis* and *H. foetidus*. If Lacourt (1994) considered that *M. longicornis* could

be restricted to *H. niger*, while *M. monticola* auct. *gallica* would be a specialist on *H. viridis* and *H. foetidus*, he proposed later that there was only one species, *Monophadnus latus sensu* Lacourt 1999 feeding on *Helleborus* spp. (Lacourt 1999, 2003). Jahn and Schedl (1992) reported that *M. longicornis* might also attack these three *Helleborus* species. Thus *M. latus sensu* Lacourt 1999 was reported to be limited to *H. niger*, *H. viridis* and *H. foetidus*, while our observations found that *Monophadnus* sp. MHNG ENTO 9805 is able to forage on 14 *Helleborus* species and sub-species, with *H. foetidus* being the most attacked among these species.

Concerning its distribution, *M. latus sensu* Lacourt 1999, is restricted to Europe and Taeger et al. (2006) have given the best inventory available at this point, completed with recent reports. The presence of *M. latus sensu* Lacourt 1999 has been recorded in Albania, Austria, Bosnia and Herzegovina, Bulgaria, Croatia, Czech Republic, Germany, Hungary, Italy and Sardinia, Romania, Spain, Switzerland and Ukraine, and previously recorded as *M. longicornis* in Austria, Italy, Romania and Ukraine. Since then it has also been reported in South West France (Savina & Chevin 2012; Chevin & Savina 2013) and North West France. Concerning Switzerland, *M. latus sensu* Lacourt 1999 was not, even recently, considered to be present in Geneva (Boillat 2010). There was only one report by Peter (2006), who mentioned a first report for South Eastern Switzerland in Ticino in 1987 on *H. niger* and another report by Pschorn-Walscher and Altenhofer (2000),

in Northern Switzerland, who also used the name *M. longicornis*. The DNA sequences provided here are the first registered for this *Monophadnus* species, identified as potentially being *M. latus sensu* Lacourt 1999. Whether the specimens from CBG correspond to this species or another undescribed species remains unknown. For this reason, a revision of the genus *Monophadnus*, supported by genetic identification would be required in order to precisely name this species.

Acknowledgements

We are grateful to Nicolas Freyre, Pascale Steinmann and the horticulturists of the domain of Rocailles, from the Conservatory and Botanical Garden of Geneva for their help in this study.

References

- ALTSCHUL SF, GISH W, MILLER W, MYERS EW, LIPMAN DJ. 1990. Basic local alignment search tool. *Journal of Molecular Biology*, 215: 403-410.
- BENSON DA, CAVANAUGH M, CLARK K, KARSCH-MIZRACHI I, LIPMAN DJ, OSTELL J, SAYERS EW. 2013. GenBank. *Nucleic Acids Research*, 41: 36-42.
- BERLAND L. 1947. Hyménoptères Tenthredoïdes [Hymenoptera Tenthredinids]. *Faune*, 47: 1-496.
- BOEVÉ JL. 2004. Sawflies (Hymenoptera: Tenthredinidae). In: Capinera JL (ed.), *Encyclopedia of Entomology*, Volume 3, Kluwer Academic Publishers; Dordrecht, pp. 1949-1954.
- BOEVÉ JL, BLANK SM, MEIJER G, NYMAN T. 2013. Invertebrate and avian predators as drivers of chemical defensive strategies in tenthredinid sawflies. *BMC Evolutionary Biology*, 13(1): 198.
- BOILLAT H. 2010. Les Hyménoptères Symphytes dans le canton de Genève, hier et aujourd'hui (Hymenoptera, Symphyta) [Hymenoptera Symphyta in the township of Geneva, yesterday and today]. *Miscellanea Faunistica Helvetica*, 7: 1-125.
- CHEVIN H, SAVINA H. 2013. Contribution à l'inventaire des Hyménoptères Symphytes du département de la Haute-Garonne [Contribution to the inventory of Hymenoptera Symphyta of the Haute-Garonne department]. *Bulletin de la Société Entomologique de France*, 118(3): 379-390.
- DOWTON M, AUSTIN AD. 1998. Phylogenetic relationships among the microgastroid wasps (Hymenoptera: Braconidae): combined analysis of 16S and 28S rDNA genes and morphological data. *Molecular Phylogenetics Evolution*, 10(3): 354-366.
- FENILI GA. 1965. Contributo allo studio morfologico-etologico del *Monophadnus longicornis* Hartig (Hymenoptera, Tenthredinidae) [Contribution to the morphological and ethological study of *Monophadnus longicornis* Hartig (Hymenoptera, Tenthredinidae)]. *Redia*, 49: 255-309.
- GUINDON S, DUFAYARD JF, LEFORT V, ANISIMOVA M, HORDIJK W, GASCUEL O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology*, 59(3): 307-21.
- HALL TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, 41: 95-98.
- HARIS A. 2015. Two new sawfly species from India (Hymenoptera: Tenthredinidae). *Natura Somogyiensis*, 26: 73-78.
- HEBERT PDN, PENTON FH, BURNS JM, JANZEN DH, HALLWACHS W. 2004. Ten species in one: DNA barcoding reveals cryptic diversity in the neotropical skipper butterfly *Astraptes fulgerator*. *Proceedings of the National Academy of Sciences of the United States of America*, 101: 14812-14817.
- JAHN E, SCHEDL W. 1992. Beobachtungen zum Auftreten einer Schneerosenblattwespe, *Monophadnus longicornis* (Hartig, 1837), in Südkärnten (Hymenoptera: Tenthredinidae) [Observations on the occurrence of snow rose sawfly, *Monophadnus longicornis* (Hartig, 1837), in Southeast Carinthia (Hymenoptera: Tenthredinidae)]. *Carinthia II*, 182(102): 453-459.

- **LACOURT J.** 1994. *Monophadnus longicornis* Hartig et *Monophadnus monticola* Hartig, deux espèces de Blennocampinae inféodées au genre *Helleborus* (Tourn.) L. (Hymenoptera, Tenthredinidae) [*Monophadnus longicornis* Hartig and *Monophadnus monticola* Hartig, two species of *Blennocampinae* subservient to the genus *Helleborus* (Tourn.) L. (Hymenoptera, Tenthredinidae)]. *L'Entomologiste*, 50(6): 335-336.
- **LACOURT J.** 1999. Répertoire des Tenthredinidae ouest paléarctiques (Hymenoptera, Symphyta) [Repertory of Tenthredinidae west-paleoartic (Hymenoptera, Symphyta)]. *Mémoires de la Société entomologique de France*, 3: 432 p.
- **LACOURT J.** 2003. Réflexions sur la classification des Blennocampinae, avec description d'un nouveau genre et d'une nouvelle espèce du sud de la France et de Corse (Hymenoptera, Tenthredinidae) [Reflexions on the classification of Blennocampinae, with the description of a new genus and a new species from South France and Corsica (Hymenoptera, Tenthredinidae)]. *Bulletin de la Société entomologique de France*, 108(5): 495-529.
- **LOONEY C, SMITH DR, COLLMAN SJ, LANGOR DW, PETERSON MA.** 2016. Sawflies (Hymenoptera, Symphyta) newly recorded from Washington State. *Journal of Hymenoptera Research*, 49: 129-159. doi: 10.3897/JHR.49.7104
- **NIU G, LIU T, CHU B, WEI M.** 2015. Four new species and a key to species of *Siobla* (Hymenoptera: Tenthredinidae) from Tibet, China. *Entomotaxonomia*, 37(4): 303-313 doi: 10.11680/entomotax.2015043
- **PETER B.** 2006. Erstnachweise von 18 Pflanzenwespenarten (Hymenoptera: Symphyta) für die Schweiz [First records of 18 sawflies species (Hymenoptera: Symphyta) for Switzerland]. In: Blank SM, Schmidt S, Taeger A (eds), *Recent sawfly research: synthesis and prospects*, : Antiquariat Goecke & Evers, Keltern, Germany, pp. 349-352.
- **PROUS M, BLANK S, GOULET H, HEIBO E, LISTON A, MALM T, NYMAN T, SCHMIDT S, SMITH D, VÅRDAL H,** et al. 2014. The genera of Nematinae (Hymenoptera, Tenthredinidae). *Journal of Hymenoptera Research*, 40: 1-69. doi: 10.3897/JHR.40.7442
- **PSCHORN-WALCHER H, ALTENHOFER E.** 2000. Langjährige Larvenaufsammlungen und Zuchten von Pflanzenwespen (Hymenoptera, Symphyta) in Mitteleuropa [Old larvae collections and breeding of sawflies (Hymenoptera, Symphyta) in Central Europe]. *Linzer biologische Beiträge*, 32(1): 273-327.
- **RAMBAUT A.** 2009. FigTree version 1.4.2. Computer program distributed by the author. Available from: <http://tree.bio.ed.ac.uk/software/figtree>. Accessed 8 March 2015.
- **SAVINA H, CHEVIN H.** 2012. Contribution à l'inventaire des Hyménoptères Symphytes du département de l'Ariège et complément sur les plantes-hôtes des espèces du genre *Abia* inféodées aux Dipsacaceae [Contribution to the inventory of Hymenoptera Symphytes of the Ariège department and supplementary on host-plant species of the genus *Abia* subservient to Dipsacaceae]. *Bulletin de la Société Entomologique de France*, 117(1): 53-70.
- **TAEGER A, BLANK S.M, LISTON AD.** 2006. European Sawflies (Hymenoptera: Symphyta) - A Species Checklist for the Countries. In: Blank SM, Schmidt S, Taeger A (eds.) *Recent Sawfly Research: Synthesis and Prospects*. Antiquariat Goecke & Evers, Keltern, Germany, pp. 399-504.
- **TAEGER A, BLANK S M; LISTON AD.** 2010. World Catalog of Symphyta (Hymenoptera). *Zootaxa*, 2580: 1-1064.
- **VILHELSEN L.** 2015. Morphological phylogenetics of the Tenthredinidae (Insecta: Hymenoptera). *Invertebrate Systematics*, 29(2): 164-190. <http://dx.doi.org/10.1071/IS14056>
- **VON DOHLEN D, MORAN NA.** 1995. Molecular phylogeny of the Homoptera: a paraphyletic taxon. *Journal of Molecular Evolution*, 41(2): 211-223.
- **WELZ C-E, VILHELSEN L.** 2014. The saws of sawflies: exploring the morphology of the ovipositor in Tenthredinoidea (Insecta: Hymenoptera), with emphasis on Nematinae. *Journal of Natural History*, 48(3-4): 133-183. <http://dx.doi.org/10.1080/00222933.2013.791941>
- **WHITING MF, CARPENTER JC, WHEELER QD, WHEELER WC.** 1997. The Strepsiptera problem: phylogeny of the holometabolous insect orders inferred from 18S and 28S ribosomal DNA sequences and morphology. *Systematic Biology*, 46: 1-68.