

# Studies on the metabolic interaction of triiodobenzoic acid during the transport of externally applied indole acetic acid

Autor(en): **Vardar, Y.**

Objektyp: **Article**

Zeitschrift: **Berichte der Schweizerischen Botanischen Gesellschaft = Bulletin de la Société Botanique Suisse**

Band (Jahr): **80 (1970)**

PDF erstellt am: **03.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-56315>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# Studies on the Metabolic Interaction of Triiodobenzoic Acid During the Transport of Externally Applied Indole Acetic Acid

By *Y. Vardar*

(Department of Botany, Ege University, Izmir, Turkey)

Manuscript received March 26, 1970

According to Kuse (1953) 2,3,5-triiodobenzoic acid (TIBA) has a blocking effect on the transport of indole acetic acid (IAA). From earlier experiments (Vardar, 1959) we concluded that TIBA causes a metabolic change of IAA. Several workers (Goldsmith and Thimann, 1962; Christie and Leopold, 1965; Winter, 1967) observed a considerable accumulation of IAA in TIBA treated plant tissues and attributed it to the immobilization effect of TIBA on IAA transport. In the present paper the effect of TIBA during transportation of IAA-C<sup>14</sup> applied externally to young seedlings is studied.

We used *Helianthus annuus* hypocotyls grown and treated with IAA-C<sup>14</sup> according to the procedures in our previous work (Vardar, 1959). The radioactive auxin was provided by the Waltham Tracer Laboratory; its specific activity was 1.1 mc/mM and the concentration 0.11 mg/ml; the amount of auxin applied was 2.2 mg and the activity applied 100 × 13 CpM. The extractions obtained from hypocotyl sections according to Nitsch and Nitsch (1955) were subjected to paper and thin layer chromatography and the radioactive counting of these was done with a Nuclear Chicago M-5 Gas Flow Counter. Paper chromatograms were run on Whatman No. 2 paper in isopropanol-ammonia-water. For thin layer chromatography the Desaga-apparatus and Kaldey's technique (1964) were used. Rf values, UV absorption traces, colour spots obtained with Salkowski reagent and autoradiographic results were compared and from them the amounts of radioactivity in various spots determined.

The results from the Salkowski colour evaluation test and the autoradiographic determinations on the chromatograms (fig. 1) show that the externally applied IAA-C<sup>14</sup> is at least partially transported as IAA in hypocotyls. It passes even under the TIBA paste ring proving thereby that TIBA does not block its transport. These results support our earlier view (Vardar, 1959) and show that IAA-C<sup>14</sup> is at least partly responsible for the radioactivity observed in the hypocotyl tissue under the TIBA paste ring. At the same time a comparison of the radioactivity evaluations in thin layer chromatograms (table 1) shows that in the various spots the activity was generally higher in TIBA treated hypocotyl tissues than in the controls. This supports the view that the TIBA treatment enhances the rate of accumulation of externally applied IAA-C<sup>14</sup> within the hypocotyls as pointed out by various workers (Goldsmith and Thimann, 1962; Christie and Leopold, 1965; Vardar, 1959; Winter, 1967).

To explain the physiological phenomena occurring during growth in length (induced by IAA) it is necessary to identify the substances derived from IAA. Several investigations (Andreae and Good, 1955; Klambt, 1960; Rausendorff-Bargen, 1962) show that IAA applied externally to the coleoptile is transformed into various substances depending on the concentration and the length of time of application. Klambt (1960) found 3-indole aspartic acid, 3-indole carboxylic acid and

3-indole acetic acid as the characteristic substances. Our studies confirmed these results, three spots on the chromatograms corresponding to 3-indole aspartic acid (0.07), 3-indole carboxylic acid (0.25) and 3-indole acetic acid (0.35); in addition to these we found a fourth radioactive derivate, the Rf value of which (0.45) corresponds roughly to 3-indole propionic acid or 3-indole butyric acid (Audus, 1959). With UV we noticed a derivate which approximately corresponds to 3-indole acetamide (Rf 0.80) which was also mentioned by Klambt; it did not show any radioactivity in our experiments.

In conclusion we may say that IAA-C<sup>14</sup>, applied externally to young green seedlings of *Helianthus annuus*, is transformed into different metabolic derivatives during the transport and that TIBA accelerates these changes. This might explain the action of TIBA on plant growth but additional biochemical experiments are necessary before drawing definite conclusions.

Table 1  
Proportional radioactivity in different zones of chromatograms

| Spot Rf |  | TIBA-treated |            | Control    |            |
|---------|--|--------------|------------|------------|------------|
|         |  | upper part   | lower part | upper part | lower part |
| 0.07    | 3-indole aspartic acid                         | 295          | 82         | 180        | 30         |
| 0.25    | 3-indole carboxylic acid                       | 45           | 18         | 28         | 15         |
| 0.35    | 3-indole acetic acid                           | 54           | 21         | 50         | 12         |
| 0.45    | 3-indole propionic or<br>3-indole butyric acid | 85           | 15         | 64         | 7          |

### Zusammenfassung

C<sup>14</sup>-markierte Indolylessigsäure (IAA, Keimlingen von *Helianthus annuus* äusserlich verabreicht) wird während des Transports in verschiedene verwandte Verbindungen (Tab. 1) umgewandelt. Trijodbenzoesäure (TIBA) bewirkt eine Anreicherung im Hypokotyl und intensiviert die Umwandlungsreaktionen.

This work was supported by the Turkish Atomic Energy Commission.

### References

- Andreae W.A. and M.E. Good. 1955. The formation of indole acetic acid in plant seedlings. *Plant Physiol.* **30**, 380-382.
- Audus L.J. 1959. *Plant growth substances*. Leonard Hill Ltd.
- Christie A.E. and A. Leopold. 1965. On the manner of triiodobenzoic acid inhibition of auxin transport. *Plant and Cell Physiol.* **6**, 337-345.
- Goldsmith M.H.M. and K.V. Thimann. 1962. Some characteristics of movement of indole acetic acid in immobilization and distribution of IAA during basipetal translocation. *Plant Physiol.* **37**, 492-505.
- Kaldewey H. 1964. Papier- und dünnschichtchromatographische Trennung, Farbreaktionen und biologischer Test einiger Wachstumsregulatoren aus Fruchtsielen von *Fritillaria meleagris* L. *Coll. Int. CNRS Paris*. Nr. **123**, 412-443.
- Klambt H.D. 1960. Wachstumsinduktion und Wuchsstoffmetabolismus in Weizenkoleoptilylindern. *Planta* **56**, 309-320.
- 1968. Cytokine aus *Helianthus annuus*. *Ibid.* **82**, 170-178.
- Kuse G. 1953. Effect of 2,3,5-triiodobenzoic acid on the growth of lateral bud and on tropism of petiole. *Mem. College of Sci. Kyoto Univ., Series B/20*, 207-215.
- Nitsch J.P. and C. Nitsch. 1955. The separation of natural plant growth substances by paper chromatography. *Beitr. Biol. Pfl.* **31**, 387-391.
- Rausendorff-Bargen G.V. 1962. Indole derivate in apple. *Planta* **58**, 471-476.
- Vardar Y. 1959. Some confirmatory experiments performed with IAA-C<sup>14</sup> concerning the effect of TIBA upon auxin transport. *Rev. Fac. Univ. Istanbul. Ser. B. XXIV, Fasc. 1-2*, 133-250.
- Winter A. 1967. The promotion of the immobilization of auxin in *Avena* coleoptiles by triiodobenzoic acid. *Physiol. Plant.* **20**, 330-336.