

Noronhia buxifolia

Objekttyp: **Chapter**

Zeitschrift: **Boissiera : mémoires de botanique systématique**

Band (Jahr): **70 (2016)**

PDF erstellt am: **27.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

9. *Noronhia buxifolia* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2 : 302. 1949.

Typus: MADAGASCAR. Prov. Toliara: Forêt d'Analavelona (bassin du Fiherenana), [22°41'S 44°08'E], 1000-1200 m, 15-19.XII.1946, Humbert 19719 (holo-: P [P00418069]!; iso-: LISC [LISC002881] image seen, P [P00418070]!, S [S09-36002] image seen, WAG [WAG0002490] image seen).

= *Noronhia buxifolia* var. *quadriloba* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2 : 302. 1949. **Typus:** MADAGASCAR. Prov. Toliara: E d'Ampandrandava, [24°05'S 45°42'E], 1200 m, XI.1943, Seyrig 373 (= Herb. Jard. Bot. 5390) (holo-: P [P04046973]!; iso-: [P00701282]!).

Description Trees to 15 m tall, trunk to 15 cm diameter; young twigs cylindrical, 0.7-1.5 mm diameter, glabrous; bark medium gray to yellowish, smooth to slightly rugose. Leaves opposite, persistent; bud scales persistent; blades dark green above, lighter below, ovate, 4-8.5 × 1.5-3.5 cm, coriaceous, glabrous, domatia absent, base rounded to acute, margin flat to slightly revolute and undulate, apex acuminate, the acumen 3-11 mm long, midrib flat to slightly sunken above, raised below, secondary veins conspicuous, 6-10 per side, 4.5-11 mm apart, looping 1-4 mm from the margin; petiole light gray, 3-9 × 1-1.5 mm, entirely woody, glabrous. Flowers fasciculate; pedicel 2.5-21 mm long, glabrous; calyx very sparsely pubescent outside, glabrous inside, lobes triangular, 0.8-3 × 0.7-2.5 mm; corolla red, urceolate, 2.5-9 mm long, glabrous on both sides, the tube 0.5-6 mm long, lobes triangular, apex obtuse; corona present, (0-)1-2 mm long, undivided to slightly 4-lobed; stamens 1.2-2.8 mm long, anthers obconical to orbicular, 0.7-1.9 mm long; pistil 1.2-2.7 mm long, stigma capitate. Fruiting pedicel 3-15 × 0.6-1.3 mm; young fruits green, dark red when mature, ovoid, 12-14.5 × 9.5-11.5 mm, surface smooth, apex apiculate to flat, with the persistent style; dry pericarp 0.5-0.7 mm thick; endocarp woody; seed 8.5-10.5 × 6.5-10 mm.

Distribution, ecology and phenology *Noronhia buxifolia* occurs in low- to high-elevation dry forests from Antsalova in the west to Andohahela in the south (Fig. 3). It produces flowers and fruits from November to June.

Conservation status Based on 21 collections representing 15 localities, the assessment indicated an EOO of 96,461 km², an AOO of 60 km² (although the true AOO is probably larger than 2,000 km²), and 11 subpopulations representing 10 locations, of which seven occur within protected areas (Amoron'i Onilahy, Analavelona, Andohahela, Behara-Tranomavo, Bemaraha, Corridor Parcel I-II Andohahela, and Menabe-Antanamena). Therefore, *N. buxifolia* is assigned a preliminary status of "Near Threatened".

Notes *Noronhia buxifolia* can be recognized by its ovate, acuminate leaf blades and fasciculate, long-pedicellate red flowers. It differs from *N. tefyana* Hong-Wa by its ovate (vs. broadly lanceolate) leaves and its urceolate (vs. cupuliform) and red (vs. pinkish tinged green) flowers. PERRIER DE LA BATHIE (1949, 1952) distinguished the variety *N. buxifolia* var. *quadriloba* H. Perrier based on differences in leaf, corolla and corona size and the margin of the corona. These differences, however, are only part of the range of variations within this species and do not justify the recognition of a distinct entity below the species level or at any other rank. Therefore, the variety *N. buxifolia* var. *quadriloba* is placed in synonymy.

Additional specimens examined

MADAGASCAR. **Prov. Mahajanga:** Tsingy de Bemaraha N of Manambolo river, 19°09'S 44°49'E, 50 m, 28.XI.1996, Jongkind et al. 3262 (MO, TAN, WAG); Besaraha, [18°41'13"S 44°15'11"E], 0-20 m, 18.XII.1952, Leandri et al. 2257 (G, MO, P); Antsalova, [18°13'S 44°35'E], 150-750 m, 11.I.1959, Réserves Naturelles 10210 (G, MO, P, TEF). **Prov. Toliara:** Morondava, forêt de Marosalaza, [20°02'S 44°33'E], 9.VI.1974, Abraham 79 (MO, P); Sakaraha, Mahaboboka, Marotsiraka-Betsileo, 22°38'36"S 44°10'17"E, 1285 m, 7.XI.2007, Andriamihajarivo et al. 1485 (MO, P, TAN); Ampandrandava, [24°05'S 45°42'E], 1200 m, XI.1943, Herb. Jard. Bot. Tan. 5390 (P); Forêt d'Analavelona (bassin du Fiherenana), [22°41'S 44°08'E], 1000-1200 m, 15-19.XII.1946, Humbert 19754bis (P); Massif de l'Andohahela et abords, au N de Imonty (RN XI), [24°49'S 46°41'E], 400-1900 m, 3.XII.1960, Leandri & Saboureau 4218 (P); Haut bassin de la Mananara et abords (SE), [24°46'S 46°43'E], 400-1900 m, 3.XII.1960, Leandri & Saboureau 4437 (G, MO, P); Forêt d'Analavelona, près d'Antanimena, 22°40'42"S 44°11'30"E, 1050 m, 12.III.1998, Messmer 597 (G, MO); Andranohinaly, [23°16'30"S 43°58'30"E], 20.VIII.1960, Peltier 2570 (P); Forêt d'Analavelona, près de la source de la rivière Manasay, 22°36'50"S 44°08'00"E, 1000-1348 m, 7.XI.2000, Randriatafika 220 (G, MO, P); ibid. loc., Randriatafika 228 (G, MO); Analavelona, [22°36'50"S 44°08'00"E], 800-1348 m, 10.XII.1955, Service Forestier 15578 (P, TEF).