

Zeitschrift: Boissiera : mémoires de botanique systématique
Band: 70 (2016)

Artikel: A taxonomic revision of the genus *Noronhia* Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
Kapitel: *Noronhia richardii*
Autor: Hong-Wa, Cynthia
DOI: <https://doi.org/10.5169/seals-1036126>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 19.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

71. *Noronhia richardii* Hong-Wa, *spec. nova* (Fig. 57).

Typus: MADAGASCAR. **Prov. Toliara:** Anosy, Fort-Dauphin, Iaboko, Antsofso, forêt Ivohibe, 24°34'26"S 47°11'42"E, 261 m, XII.2005, Razakamalala *et al.* 2666 (holo-: MO-6615556!; iso-: G [G00341619]!, P [P06490472] image seen, TAN).

Diagnosis *Noronhia richardii* Hong-Wa can be distinguished from other congeneric species by its lanceolate leaves, and its long and diffuse inflorescences with red flowers.

Description Small trees to 7 m tall; young twigs cylindrical, 0.7-1.4 mm diameter, glabrous; bark medium gray, smooth. Leaves opposite, persistent; bud scales rarely persistent; blades medium green above, yellowish below, lanceolate to oblong, 9-11 × 2-4 cm, chartaceous, glabrous, domatia absent, base acute to rounded, margin slightly revolute, apex acuminate, the acumen 5-9 mm long, midrib flat above, slightly raised below, secondary veins conspicuous, 9-12 per side, 9-17 mm apart, looping 1.3-3 mm from the margin; petiole medium gray, 6-11 × 1-2.5 mm, entirely woody, glabrous. Thyrses geminate, pauciflorous, diffuse; peduncle 4-30 mm long, glabrous; pedicel 11-25 mm long, glabrous; calyx sparsely pubescent to glabrescent outside, glabrous inside, lobes ovate, 1-2.5 × 1-2 mm; corolla red, cupuliform, 3-7 mm long, glabrous on both sides, the tube 1-5 mm long lobes widely ovate, apex acute; corona absent; stamens 2-3 mm long, anthers widely oblong to almost square, 1.6-1.9 mm long; pistil 2.2-3.5 mm long, stigma capitate. Fruits unseen.

Etymology This name honors Richard Razakamalala, a botanist at the Missouri Botanical Garden in Madagascar and one of the most knowledgeable Malagasy field botanists, for collecting the type specimen and for providing invaluable preliminary identification on countless field collections.

Distribution, ecology and phenology *Noronhia richardii* occurs in low-elevation humid forests on basement rocks at Bemangidy-Ivohibe north of Ste Luce in the south (Fig. 49). It has been collected in flowers in November and December.

Conservation status Only three collections, representing three localities, were available and were all obtained from a single location within the recently established protected area of Tsitongambarika. The subpopulation occurs on the northeastern edge of the protected area, where significant human activities still put some pressure on the forest, resulting in continuing decline in habitat quality. Thus, with an AOO of just 8 km², a single location, and lingering threats that could rapidly affect the area, *N. richardii* is assigned a preliminary status of "Vulnerable" [VU D2].




Fig. 57.

Noronhia richardii Hong-Wa. **A.** Flowering branch; **B.** Abaxial side of leaf blade; **C.** Adaxial side of leaf blade; **D.** Flower; **E.** Inner side of corolla; **F.** Pistil; **G.** Stamen.

[Razakamalala 2424, TAN] Drawings: R.L. Andriamiarisoa

Notes *Noronhia richardii* closely resembles *N. capuronii*, from which it differs by its glabrous (vs. pubescent) petioles, lanceolate to oblong (vs. ovate) leaf blades with an acute to rounded (vs. cordate) base, long peduncles and pedicels (up to 30 mm vs. 8 mm), and red (vs. greenish) flowers. It can be recognized by its lanceolate leaves and long, diffuse inflorescences with red flowers.

Paratypes MADAGASCAR. **Prov. Toliara:** Fort-Dauphin, Iaboko, Antsotso, forêt d'Ivohibe, 24°34'14"S 47°12'04"E, 265 m, 28.XI.2005, *Razakamalala et al.* 2424 (G, MO, P, TAN); *ibid. loc.*, 24°34'16"S 47°12'06"E, 271 m, 8.XII.2007, *Razakamalala et al.* 3777 (MO, P, TAN).