

# Chromosome numbers of some gentianaceae from the Caucasus

Autor(en): **Gagnidze, Revas / Küpfer, Philippe / Yuan, Yong-Ming**

Objekttyp: **Article**

Zeitschrift: **Bulletin de la Société Neuchâteloise des Sciences Naturelles**

Band (Jahr): **115 (1992)**

PDF erstellt am: **12.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-89351>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# CHROMOSOME NUMBERS OF SOME *GENTIANACEAE* FROM THE CAUCASUS

by

REVAS GAGNIDZE<sup>1</sup>, PHILIPPE KÜPFER<sup>2</sup> and YONG-MING YUAN<sup>2</sup>

WITH 1 TABLE

---

## INTRODUCTION

*Gentianaceae* is a fairly cosmopolitan family comprizing about 80 genera and 900 species, mostly represented in temperate zones and mountainous regions of the tropics. Various aspects of most species belonging to the genus *Gentiana* have been extensively studied throughout the world. According to HO and LIU (1990), the genus consists of 15 sections and 361 species with the highest concentration of species in SW China and NE Burma. However, the European sections namely *Gentiana* and *Ciminalis* (Adams.) Dumort. are very different from the Asiatic ones. Chromosome numbers of all the European species have been determined (e.g. FAVARGER 1949, 1952; RORK 1949; SKALINSKA 1951; LÖVE 1953; LÖVE and LÖVE 1975, 1986; KÜPFER 1980; MÜLLER 1982). Some regional karyological studies in Asia have also been carried out, in Japan (SHIGENOBU 1983, 1984), the Himalayas (VASUDEVAN 1975) and China (KÜPFER and YUAN, YUAN, YUAN and KÜPFER, to be published). Nevertheless, some transitional areas have been less studied, though detailed studies of these regions may reveal more concerning the relationships among taxa of different regions. The Caucasus linking W.Asia and E.Europe, is a typical transitional area. About 40 *Gentianaceae* have been recorded in this area (GROSSHEIM 1967). Very few of them have been observed karyologically.  $2n = 42$  has been reported for *Swertia iberica* Fisch. and Mey. from this area (SOKOLOVSKAYA and STRELKOVA 1948 cited from AGAPOVA 1990), but recently it has been counted as  $2n = 26$  (GAGNIDZE et al. 1986).  $2n = 26$  has also been reported for *Gentiana pyrenaica* L. from this area by the above two authors (cited from AGAPOVA 1990).

<sup>1</sup> Institute of Botany of the Georgian Academy, 380007 Tbilisi, Georgia.

<sup>2</sup> Institut de botanique, Université de Neuchâtel, CH-2007 Neuchâtel.

The present short report is a summary of our preliminary observations on the karyology of *Gentianaceae* from the Caucasus, concerning 11 species from 3 genera. Chromosome numbers for 6 species were determined for the first time.

#### MATERIALS AND METHODS

The species examined along with their origins are listed in Table 1. All the voucher specimens have been deposited in the Herbarium of the University of Neuchâtel (NEU). Chromosome numbers were determined from mitosis of root-tips, young ovaries and meiosis of pollen mother cells as well as from pollen mitosis. For observations on mitosis of root-tips, seeds collected in the field were germinated in petri dishes following a 24 hour treatment with 0,3 % gibberellic acid ( $GA_3$ ) (MÜLLER 1977). Root-tips pretreated with a saturated aqueous solution of  $\alpha$ -bromonaphthalene for 1 hour and 20 minutes, were stained and squashed with the aceto-orcein method (KRÄHENBÜHL and KÜPFER, to be published). SNOW'S (1963) method was used for observations on meiosis and the mitosis of young ovaries and pollens.

#### RESULTS AND DISCUSSION

The chromosome numbers of the species examined are shown in Table 1. Among them, chromosome numbers for *Gentiana angulosa* ( $2n = 30$ ), *Gentiana gelida* ( $2n = 26$ ), *Gentiana kolakovskiyi* ( $2n = 26$ ), *Gentianella caucasea* ( $2n = 36$ ), *Gentianella biebersteinii* ( $2n = 36$ ) and *Gentianella umbellata* ( $2n = 18$ ) were determined for the first time. The other numbers confirmed previous reports.

In the genus *Gentiana*,  $2n = 48$  has been reported for *G. aquatica* from Siberia (BELAEVA and SIPLIVINSKY 1977 cited from AGAPOVA 1990) and *G. atlantica* Litard. and Maire from North Africa (KÜPFER 1980). Our present observations revealed that *G. aquatica* was fairly stable karyologically with the same number of  $2n = 48$  in individuals from three populations in the Caucasus, the western limits of this species' geographical distribution. Although *G. aquatica* and *G. atlantica* share the same chromosome number and belong to the same section (*Chondrophyllae* Bunge), they are quite distinct geographically and morphologically. Whether the same number arised independently in the two species still needs to be confirmed by more studies. *G. pyrenaica* of the section *Chondrophyllae* was confirmed as  $2n = 26$  from the Caucasus, near the eastern limit of its distribution. Different numbers of  $2n = 36$  and 44 have been reported for *G. asclepiadea*. Our results revealed  $2n = 44$ , a figure consistent with most reports on the species. *G. gelida*, *G. kolakovskiyi* and *G. septemfida* were all  $2n = 26$  confirming the fairly uniform chromosome number of the section *Pneumonanthae* (Gled.) Gaudin to which the above three species belong. *G. angulosa* was shown to have  $2n = 30$  similar to other parapatrically distributed taxa of the section *Calathianae* Froelich in that area, such as *G. verna* subsp. *balcanica* Pritchard, *G. pontica* Soltok. and *G. oschtenica* (Kusn.) G. Woron. (MÜLLER 1982).

TABLE 1 *Chromosome numbers of some Gentianaceae from the Caucasus*

Taxon	Origin and altitude	Material	Chrom. No.	Previous report
<i>Gentiana aquatica</i> L.	C, Kazbegi, 2900 m	root-tips	2n=48	2n=48 (cited from AGAPOVA 1990)
	C, Djvari Pass, 2400 m	root-tips	2n=48	
	C, vicinity of Djvari, 2500 m	root-tips	2n=48	
<i>G. asclepiadea</i> L.	T, Bakuriani, 1900 m	ovaries	2n=44	2n=44 (FAVARGER 1949, RORK 1949, SKALINSKA 1951, MAJOVSKY; 1974a) 2n=36 (SKALINSKA 1950, MAJOVSKY 1974b)
<i>G. angulosa</i> Bieb.	C, E. of Djvari Pass, 2900 m	root-tips	2n=30*	
<i>G. gelida</i> Bieb.	T, between Tbilisi and Tsalka, Trialetskiy Range, 1600 m	ovaries	2n=26	
<i>G. kolakovskiyi</i> Doluch.	C, Inguri Valley, 1000 m	ovaries	2n=26*	
<i>G. pyrenaica</i> L.	C, E. of Djvari Pass, 2900 m	root-tips	2n=26	2n=26 (FAVARGER and KÜPFER 1968, KÜPFER and FAVARGER 1967, LÖVE and LÖVE 1975, 1986)
	C, Terek Valley, 2100 m	ovaries, anthers	2n=26, n=13	
<i>G. septemfida</i> Pall.	C, Terek Valley, 2100 m	root-tips	2n=26	2n=26 (ZHUKOVA 1967, AGAPOVA 1990)
		ovaries, anthers	2n=26, n=13	
<i>Gentianella caucasea</i> (Loddiges ex Sims) Holub	C, Kazbek, 3200 m	root-tips	2n=36*	
<i>G. biebersteinii</i> (Bunge) Holub	C, Terek Valley, 2100 m	root-tips	2n=36*	
<i>G. umbellata</i> (Bieb.) Holub	C, Djvari Pass, 2150 m	root-tips	2n=18*	
		root-tips	2n=18*	
<i>Lomatogonium carinthiacum</i> (Wulfen) A. Braun	C, Terek Valley, 2100 m	root-tips	2n=48	2n=40 (FUERNKRANZ 1965); 2n=32 (KROGULEVICH 1978, LÖVE and LÖVE 1986); n=24 (VASUDEVAN 1975)

C: Caucasus (Bolshoy Kavkaz) T: Transcaucasia

\* first report of chromosome number for the species.

In the genus *Gentianella*, most of the previous reports are of  $2n = 36$  for most species (e.g. WEAVER and RÜDENBERG 1975), except that  $2n = 18$  has been reported for *G. acuta* Michx. from Siberia (KROGULEVICH 1978), *G. amarella* L. from California (POST 1983) and *G. moorcroftiana* from the Himalayas (MEHRA and VASUDEVAN 1972; VASUDEVAN 1975). Our present study revealed  $2n = 36$  for *G. caucasea* and *G. biebersteinii* and  $2n = 18$  for *G. umbellata*. About *G. caucasea* FAVARGER (personal communication) found, but did not publish, the same number of  $2n = 36$  in *Gentiana caucasica* Bieb. from Erevan (Armenia). Therefore *Gentianella caucasea* would be synonymous to *Gentiana caucasica* var. *markoviczii* Kusn. The relationship between *Gentianella caucasea* and *Gentiana caucasica* would be close.

The basic number of  $x = 9$  for the genus *Gentianella* is now well established. Most species known karyologically are tetraploid, a few of them are diploid. Tetraploid species are distributed in North America, the Himalayas and Caucasus, which suggests that the polyploidisation is polytopical.

Our present observation confirmed the chromosome number of  $2n = 48$  for *Lomatogonium carinthiacum* which was previously reported as  $n = 24$  from Kashmir in the western Himalayas (VASUDEVAN 1975). A similar but not published result of  $n = 23-24$  was obtained by FAVARGER (personal communication) on material from the Swiss Alps (Grisons, Val-d'Avers, Bergalgabach, 2000 m). Some other results were also published:  $2n = 40$  from the Austrian Alps (FUERNKRANZ 1965),  $2n = 32$  from Siberia (KROGULEVICH 1978) and the Mts Bucegi in Roumania (LÖVE and LÖVE 1986). We suppose the basic number of the species to be  $x = 8$  instead of  $x = 6$  as was suggested by VASUDEVAN (1975), and furthermore, an intraspecific polyploidy exists; the tetraploid cytotype ( $2n = 32$ ) and the hexaploid cytotype ( $2n = 48$ ) may be distributed parapatrically. The number  $2n = 40$  from the Austrian Alps would represent a pentaploid type. The phytogeographical value of this result has yet to be demonstrated.

---

### Summary

Chromosome numbers were determined for the first time for *Gentiana angulosa* ( $2n = 30$ ), *Gentiana gelida* ( $2n = 26$ ), *Gentiana kolakovskiyi* ( $2n = 26$ ), *Gentianella caucasea* ( $2n = 36$ ), *Gentianella biebersteinii* ( $2n = 36$ ) and *Gentianella umbellata* ( $2n = 18$ ); chromosome numbers were confirmed for *Gentiana aquatica* ( $2n = 48$ ), *Gentiana asclepiadea* ( $2n = 44$ ), *Gentiana pyrenaica* ( $2n = 26$ ) and *Gentiana septemfida* ( $2n = 26$ ). The chromosome number  $2n = 48$  was confirmed for *Lomatogonium carinthiacum* and the basic number of  $x = 8$  was suggested for the species.

### Résumé

Les nombres chromosomiques de *Gentiana angulosa* ( $2n = 30$ ), *Gentiana gelida* ( $2n = 26$ ), *Gentiana kolakovskiyi* ( $2n = 26$ ), *Gentianella caucasea* ( $2n = 36$ ), *Gentianella biebersteinii* ( $2n = 36$ ) et *Gentianella umbellata* ( $2n = 18$ ) ont été déterminés pour la première fois. Dans le domaine eurasiatique, *Gentianella umbellata* représente l'espèce la plus occidentale du genre à être diploïde. Toutes les espèces européennes étudiées jusqu'ici se sont révélées tétraploïdes. Le nombre chromosomique ( $2n = 48$ ) du *Lomatogonium carinthiacum* laisse penser que les populations caucasiennes sont hexaploïdes, avec  $x = 8$  comme nombre de base, comme les populations du Cachemire et peut-être aussi comme celles des Alpes, en partie tout au moins.

### REFERENCES

- AGAPOVA, N. D. (ed.) — (1990). Numeri Chromosomatum Magnoliophytorum Florae URSS. *Leninopoli*.
- FAVARGER, C. — (1949). Contribution à l'étude caryologique et biologique des Gentianaceae. *Bull. Soc. Bot. Suisse* 59: 62-86.
- (1952). Contribution à l'étude caryologique et biologique des Gentianaceae. II. *Bull. Soc. Bot. Suisse* 62: 244-257.
- FAVARGER, C. and KÜPFER, Ph. — (1968). Contribution à la cytotaxonomie de la flore alpine des Pyrénées. *Collect. Bot. (Barcelona)* 7: 325-357.
- FUERNKRANZ, D. — (1965). Einige Chromosomenzahlen von Pflanzen aus den Österreichischen Alpen. *Österr. Bot. Z.* 112: 421-423.
- GAGNIDZE, R., GVINIASCHVILI, T. and TSCHURADSE, M. — (1986). Numeri chromosomatum specierum nonnullarum endemicarum florum Caucasi. *Notul. Syst. Geograph. Inst. Bot. Tbilissiensis* 41: 80-84.
- GROSSHEIM, A. A. — (1967). Flora Kavkaza. Vol. VII, pp. 202-217. *Baku*.
- HO, T.-N. and LIU, S.-W. — (1990). Infrageneric classification of *Gentiana* (Gentianaceae). *Bull. Brit. Mus. Nat. Hist. (Bot.)* 20: 168-192.
- KRÄHENBÜHL, M. and KÜPFER, Ph. — (to be published). Basic chromosome number and systematic position of the genus *Molopospermum* Koch in the Umbelliferae. *Bauhinia*.
- KROGULEVICH, R. E. — (1978). Kariologicheskij analiz vidov flory vostochnogo Sajana. V. Flora Pribajkalja., pp. 19-48. *Nauka, Novosibirsk*.
- KÜPFER, Ph. — (1980). A propos de quelques gentianes de la section *Chondrophyllae* Bunge. *Bol. Soc. Brot. Sér. 2*, 53: 537-547.
- KÜPFER, Ph. and FAVARGER, C. — (1967). Premières prospections caryologiques dans la flore orophile des Pyrénées et de la Sierra Nevada. *Compt. Rend. Hebd. Séances Acad. Sci.* 264: 2463-2465.
- KÜPFER, Ph. and YUAN, Y.-M. — (to be published). Contribution to cytotaxonomy and cytogeography of *Gentiana* (Gentianaceae) from China. II. Karyological studies on sect. *Chondrophyllae* Bunge. *Pl. Syst. Evol.*
- LÖVE, D. — (1953). Cytotaxonomical remarks on the Gentianaceae. *Hereditas* 39: 225-235.
- LÖVE, A. and LÖVE, D. — (1975). The spanish gentians. *Anal. Inst. Bot. Cavanilles* 32, II: 221-232.
- (1986). In IOPB chromosome number reports XCIII. *Taxon* 35: 897-903.

- MAJOVSKY, J. (ed.) — (1974 *a*). Index of chromosome number of Slovakian flora (part 3). *Acta Fac. Rerum Nat. Univ. Comenianae, Bot.* 22: 1-20.
- (1974 *b*). Index of chromosome number of Slovakian flora (part 4). *Acta Fac. Rerum Nat. Univ. Comenianae, Bot.* 23: 1-23.
- MEHRA, P. N. and VASUDEVAN, K. N. — (1972). In IOPB chromosome number reports XXXVI. *Taxon* 21: 333-346.
- MÜLLER, G. — (1977). Action de l'acide gibbérellique sur la germination des gentianes de la section *Cyclostigma*. *Bull. Soc. Neuchâtel. Sci. Nat.* 100: 121-126.
- (1982). Contribution à la cytotaxonomie de la section *Cyclostigma* Griseb. du genre *Gentiana*. *Feddes Repert.* 93: 625-722.
- POST, D. M. — (1983). In IOPB chromosome number reports LXXX. *Taxon* 32: 509.
- RORK, C. L. — (1949). Cytological studies in the Gentianaceae. *Amer. J. Bot.* 36: 687-723.
- SHIGENOBU, Y. — (1983). Karyomorphological studies in some genera of Gentianaceae II. *Gentiana* and its allied four genera. *Bull. Coll. Child Develop., Kochi Women's Univ.* 7: 65-84.
- (1984). Karyomorphological studies in some genera of Gentianaceae II. *Gentiana* and its allied four genera. *Bull. Coll. Child Develop., Kochi Women's Univ.* 8: 55-104.
- SKALINSKA, M. — (1950). Studies in chromosome numbers of Polish angiosperms. *Acta Soc. Bot. Poloniae* 20: 45-68.
- (1951). Cytological studies in *Gentiana* species from the Tatra and Pieniny Mts. *Bull. Acad. Polon. Sci., Sér. Sci. Biol.* 1951: 119-136.
- SNOW, R. — (1963). Alcoholic hydrochloric acid-carmines as a stain for chromosomes in squash preparations. *Stain Techn.* 38: 9-13.
- VASUDEVAN, K. N. — (1975). Contribution to cytotaxonomy and cytogeography of the flora of Western Himalayas (with an attempt to compare it with the flora of the Alps). Part I. *Ber. Schweiz. Bot. Ges.* 85: 57-84.
- WEAVER, R. E. and RÜDENBERG, L. — (1975). Cytotaxonomic notes on some Gentianaceae. *J. Arnold Arb.* 56: 211-222.
- YUAN, Y.-M. — (to be published). Contribution to cytotaxonomy and cytogeography of *Gentiana* (Gentianaceae) from China. I. General introduction and karyological studies on sect. *Cruciata* Gaudin. *Caryologia*.
- YUAN, Y.-M. and KÜPFER, Ph. — (to be published). Contribution to cytotaxonomy and cytogeography of *Gentiana* (Gentianaceae) from China. III. Karyological studies on sect. *Frigida* Kusn. s. l. and sect. *Stenogyne* Franch. *Nord. J. Bot.*
- ZHUKOVA, P. G. — (1967). Karyology of some plants, cultivated in the Arctic Botanical Garden. In: AVRORIN, N. A. (ed.), *Plantarum in Zonam Polarem Transportatio*. II. *Leningrad*, pp. 139-149.
-