

Zeitschrift: Bulletin de la Société Neuchâteloise des Sciences Naturelles
Band: 125-1 (2002)

Artikel: A new species of Freycinetia (Pandanaceae) from the Santa Cruz Islands
Autor: Huynh, Kim-Lang
DOI: <https://doi.org/10.5169/seals-89569>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 05.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A NEW SPECIES OF *FREYCINETIA* (PANDANACEAE) FROM THE SANTA CRUZ ISLANDS

KIM-LANG HUYNH

Evolutionary Botany Division, Botanical Institute, University of Neuchâtel, PO Box 2, CH-2007 Neuchâtel, Switzerland.

Mots-clés: Pandanacées, *Freycinetia*, Taxonomie, îles de Santa Cruz

Keywords: Pandanaceae, *Freycinetia*, Taxonomy, Santa Cruz Islands

Résumé

Une espèce nouvelle de *Freycinetia* des îles de Santa Cruz est décrite (*F. santacruzensis*).

Summary

A new species of *Freycinetia* is described from the Santa Cruz Islands (*F. santacruzensis*).

Zusammenfassung

Eine neue *Freycinetia*-Art aus den Santa Cruz Inseln wird beschrieben (*F. santacruzensis*).

INTRODUCTION

The Santa Cruz Islands are a group of melanesian islands situated north of Vanuatu (the New Hebrides) and south-east of the Solomon Islands. A new species of *Freycinetia* has been found in the Santa Cruz Islands: it is described below.

DESCRIPTION

Freycinetia santacruzensis Huynh, sp. nova (sect. *Oligostigma*) (fig. 1)

Folia infra infructescentiam 75-95 cm longa, 5.5-6 cm lata in medio 5 cm in basi (auriculis non inclusis), ensiformia, in apice abrupte attenuata, 1.5-2 cm caudata, imbricata, in basi amplexicaulia; in sicco subcoriacea, pallida, striata in pagina adaxiali; venis longitudinalibus distinctis in pagina abaxiali minus in adaxiali; venis transversalibus utrinque invisibilibus; marginibus armatis in 5-8 cm summis solum, denticulis usque ad 2/3 mm longis; costa media armata in 3-5 cm summis, denticulis usque ad 2/3 mm longis, raro cum 2 vel 3 denticulis in medio usque ad 1.3 mm longis; auriculis omnibus conservatis, 15 cm longis, 1.2 cm latis, in apice adnatis, pallidis, submembranaceis, inermibus, utrinque longitudinaliter nerviis sed non striatis. Infructescentia terminalis, 2 (?) spicis praedita, pedunculo communi 9 cm longo, 8 mm crasso; bracteis floralibus 10-11 cm longis, 5 cm latis, navicu-

laribus, breviter acuminatis, tote inermibus, utrinque dense longitudinaliter nerviis, atrobrunneis; syncarpiis cylindratis apice basique rotundatis, 8 cm longis, 3.8-4 cm latis, maturis, pedunculis 4 cm longis, 5-6 mm latis, laevibus. Baccae filiformes, 15 mm longae, 1 mm latae, sclerenchymate centrali in pileo et longissimis fasciculis fibrarum, ex apice ad basim non interruptis, non fusiformibus/ellipticis, praeditae; pileo 5 mm longo, 1 mm lato, anguste pyramidalis, fusco; ovario 10 mm longo; stigmatibus 2, distinctis et oppositis, interdum 4, areola stigmatica annulo perspicuo cincta. Semina 1.23 mm longa, 0.42 mm lata in medio, recta; strophio indistincto; endospermio 0.39 mm lato; raphe 0.03 mm lata, cellulis raphidiphoris praedita sed crystalliferis destituta, cellulis circum fasciculum vasorum lignosis.

Typus: *Yen 804* (holo-BISH!); Vanikoro Island (Santa Cruz Islands), south coast, permanently wet coastal swamp extending inland from dune, alt. 300 m, September 1978; common in swamp; low tree, 6-12 ft tall.

Note: Vanikoro Island is at about 11°42'S 166°50'E in its central position.

Freycinetia santacruzensis seems to be the first species of *Freycinetia* described from the Santa Cruz Islands. Although its occurrence in these islands was reported some years ago (HUYNH, 1996: 540), it is described for the first time in the present paper.

Freycinetia santacruzensis is noteworthy by its leaves which are armed in a short apical part only, and by its floral bracts which are entirely unarmed. These two features are not common in *Freycinetia*.

Freycinetia santacruzensis is a species of sect. *Oligostigma* sensu Warburg (1900), non STONE (1968: 367). In Stone's classification, it has its position in sect. *Filiformicarpae* (STONE, 1968: 366). This latter section may be recognized by the following characteristics: berries filiform;

stigmas mostly 2 and opposite, sometimes 4; in particular, the berries have long fibre-strands extending in a single stretch from the apex to the base. These fibre-strands were to date observed in sect. *Filiformicarpae* only.

Freycinetia santacruzensis appears closest to *F. tessellata* (Solomon Islands) and *F. filiformis* (Bismarck Archipelago) - which also belong to sect. *Filiformicarpae* (HUYNH, 1996: 539) - , considering in particular the size of their leaves and the length of their mature syncarps. *F. tessellata* differs from *F. santacruzensis* in its leaves which are tessellate and armed in the basal part and the apical part of the margins; its syncarp peduncles are scabrid; in particular, its syncarps are clavato-cylindric (MERRILL & PERRY, 1939: 149), a rare feature in the genus *Freycinetia*. *F. filiformis* differs from *F. santacruzensis* in its leaves which are tessellate, rigid-coriaceous, and armed in the basal part and the apical part of the margins; its leaf auricles are striate between the longitudinal veins; its pileus is 2 mm long for a berry of 13 mm (HUYNH, 1996: 537).




Figure 1: *Freycinetia santacruzensis* (*Yen 804*): Upper part of leaf flattened horizontally, viewed at the adaxial face.

ACKNOWLEDGMENTS

The author is grateful to Bishop Herbarium (BISH!) for the loan of material.

REFERENCES

- HUYNH, K.-L. 1996. The genus *Freycinetia* (Pandanaeae) in New Guinea (part 1). *Bot. Jahrb. Syst.* 118: 529-545.
- MERRILL, E. D. & PERRY, L. M. 1939. On the Brass collections of Pandanaeae from New Guinea. *J. Arnold Arbor.* 20: 139-186.
- STONE, B. C. 1968. Materials for a monograph of *Freycinetia* Gaud. IV. Subdivision of the genus, with fifteen new sections. *Blumea* 16: 361-372.
- WARBURG, O. 1900. Pandanaeae. In: A. ENGLER (ed.), *Das Pflanzenreich* 3 (= IV.9). *Engelmann, Leipzig.*