

Zeitschrift: Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber: Conservatoire et Jardin botaniques de la Ville de Genève
Band: 42 (1987)
Heft: 1

Artikel: Rumex gangotrianus : a new species of Polygonaceae from India
Autor: Aswal, B.S. / Srivastava, S.K.
DOI: <https://doi.org/10.5169/seals-879952>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 24.01.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Rumex gangotrianus — a new species of Polygonaceae from India

C.D.R.I. Communication No. 4010

B. S. ASWAL
&
S. K. SRIVASTAVA

RÉSUMÉ

ASWAL, B. S. & S. K. SRIVASTAVA (1987). *Rumex gangotrianus* — une espèce nouvelle de Polygonacées d'Inde. *Candollea* 42: 389-391. En anglais, résumé français.

Une nouvelle espèce *Rumex gangotrianus* Aswal & Srivastava spec. nov. (Polygonaceae), récoltée au voisinage de Gangotri, Uttarkashi, Uttar Pradesh, Inde, est décrite et illustrée.

ABSTRACT

ASWAL, B. S. & S. K. SRIVASTAVA (1987). *Rumex gangotrianus* — a new species of Polygonaceae from India. *Candollea* 42: 389-391. In English, French abstract.

A new species *Rumex gangotrianus* Aswal & Srivastava spec. nov. (Polygonaceae) collected from the vicinity of Gangotri, Uttarkashi, Uttar Pradesh, India, is described with illustration.

During a recent plant collection tour to the upper Bhagirathi valley in Uttarkashi district, Uttar Pradesh, India, one of the authors (BSA) collected an interesting specimen of *Rumex* growing near the population of *Rumex nepalensis* on moist shady slopes. A critical study of the literature and comparison with the herbarium specimens of the known species of *Rumex* at the Indian herbaria (BSD, CAL, CDRI, DD, LWG) and at the Royal Botanic Gardens, Kew, England (K) proved it to be a hitherto unrecognized species. Hence, it is described here as a new species.

Rumex Linn. is represented by ca. 200 species in the world, distributed specially in all the temperate and some tropical countries. About 15 species are known from India, mostly from the Himalayan region.

***Rumex gangotrianus* Aswal & Srivastava spec. nov.** (Figs. 1-6).

Typus: India: Uttar Pradesh, Uttarkashi district, between Lanka and Gangotri, 2800 m, 5.6.1986, *B. S. Aswal 11370A* (holo, CDRI); *B. S. Aswal 11370B, C* (iso, CDRI).

R. nepalensis Spreng. proxime affinis, sed differt foliis basalibus permajoribus, basibus late cordatis, floribus bisexualibus, confertim dispositis in paniculis densis; tepalorum segmentis interioribus orbiculari-ovatis, reticulatim venosis; staminibus cum filamentis brevibus, antheris apicibus acutis.

This species is closely allied to *R. nepalensis* Spreng., but differs in having much larger basal leaves with broad cordate base; flowers bisexual, compactly arranged in dense panicle; tepals — inner segments orbicular-ovate, reticulately veined; stamens with short filament, anthers with acute apex.

Erect, annual herbs, ca. 65-110 cm tall. Stems stout, fistular, branched, ribbed, minutely puberulous often swollen at nodes. Leaves simple, alternate, 6-15(-32) × 2-8(-15) cm, ovate, ovate-




Fig. 1. — *Rumex gangotrianus* Aswal & Srivastava.

1, habit. 2, flower (cut open). 3, tepal (outer). 4, tepal (inner). 5a, stamen (lateral view). 5b, stamen (dorsal view). 6, pistil.

oblong or triangular, glabrous, apex acute to acuminate, margin entire or somewhat undulate, midrib distinct on both the surfaces, lateral veins 10-12 pairs, basal leaves long petioled much larger with broad cordate base, upper leaves short petioled; petioles 0.6-1.8 cm long, minutely puberulous; stipules membranous, fringed. Flowers 1.5 cm across, purplish green, bisexual, whorled, compactly arranged in dense panicles; pedicels 5 mm long, filiform. Perianth 6 cleft, biseriate, tepals distinct (2 whorls of 3 tepals), inner segments 3×2 mm, orbicular-ovate, reticulately veined, outer segments 2×1 mm, oblong-ovate, margin incurved. Stamens 6, each enclosed in front of each tepal, anthers oblong, 1 mm long, filament short. Pistil 2 mm long; ovary ovoid, triangular ca. 1 mm in diam.; styles 3, filiform, 0.5 mm long, stigma fimbriate.

Flowering: May-July.

Etymology: The specific epithet has been derived from the locality "Gangotri", the source of the holy river Ganga.

ACKNOWLEDGEMENTS

The authors are thankful to the Director, Central Drug Research Institute, Lucknow for facilities; to Dr. B. N. Mehrotra, Head, Botany Division for encouragement; and to Dr. Ved Prakash, Scientist, Botany Division for matching the specimen at the herbarium of the Royal Botanic Gardens, Kew, England.

Thanks are also due to Dr. N. C. Majumdar, Scientist, Botanical Survey of India, Howrah for providing Latin diagnosis.

