

# Cancellation Properties of H-Spaces

Autor(en): **Mislin, Guido**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **49 (1974)**

PDF erstellt am: **16.08.2024**

Persistenter Link: <https://doi.org/10.5169/seals-37988>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## Cancellation Properties of $H$ -Spaces

GUIDO MISLIN<sup>1)</sup>

One defines the genus of a homotopy type  $X$  to be the set  $G(X)$  of all homotopy types  $Y$  such that the  $p$ -localizations  $X_p$  and  $Y_p$  are homotopy equivalent for all primes  $p$  [11]. To have a good notion of  $p$ -localization, we will work throughout this paper in the homotopy category of nilpotent, connected, pointed CW-complexes [1, 7, 14]. Notice that all  $H$ -spaces are nilpotent spaces whereas  $H'$ -spaces are in general not nilpotent unless they are 1-connected. We call  $X$  *quasi-finite*, if  $H_i(X; \mathbf{Z})$  is finitely generated for all  $i$  and zero for  $i$  big enough. It is not known whether a quasi-finite  $H$ -complexes is necessarily of the homotopy type of a finite complex.

It was observed by many authors [5, 8, 9, 10, 11, 13, 15] that  $H$ -spaces, cancellation phenomena and the genus of a space are closely related. The oldest example expressing this relation is the  $H$ -manifold  $E_{7w}$  of [6] for which one has

$$G(E_{7w}) = G(Sp(2)) \quad (\text{see [11]})$$

and

$$E_{7w} \times S^3 \simeq Sp(2) \times S^3 \quad (\text{see [6]}).$$

More complicated examples of the same nature, involving the Lie group  $G_2$ , were discussed in [8].

Our main results (Proposition 1 and 2) give conditions under which one can cancel factors in a product or summands in a wedge respectively. When applied to  $H$ -spaces, this yields a converse of a theorem of Zabrodsky [15], giving a very simple characterization of the genus of an  $H$ -complex (Theorem A below).

For a quasi-finite  $H$ -complex  $X$  one has by Hopf's theorem  $H^*(X; \mathbf{Q}) \cong H^*(S^{n_1} \times \cdots \times S^{n_k}; \mathbf{Q})$ ; the array  $(n_1, \dots, n_k) = \tau(X)$  is called the *type* of  $X$ . We will prove

**THEOREM A.** *Let  $X$  and  $Y$  be quasi-finite  $H$ -complexes. Then the following are equivalent*

$$A1: G(X) = G(Y)$$

$$A2: X \times \left( \prod_{\tau(X)} S^{n_i} \right) \simeq Y \times \left( \prod_{\tau(Y)} S^{m_j} \right)$$

---

<sup>1)</sup> The author was partially supported by NSF Grant GP-29544 X.

Dually, if  $X$  is a quasi-finite nilpotent  $H'$ -complex then  $\pi_*(X) \otimes \mathbf{Q} \cong \pi_*(S^{m_1} \vee \dots \vee S^{m_k}) \otimes \mathbf{Q}$  and we call  $(m_1, \dots, m_k) = \tau'(X)$  the *cotype* of  $X$ ; notice that if  $X$  is not 1-connected, then  $X$  is of the form  $S^1 \vee Y$ , and the nilpotency of  $X$  implies  $X \simeq S^1$ . Then one has

**THEOREM B.** *Let  $X$  and  $Y$  be quasi-finite nilpotent  $H'$ -complexes. Then the following are equivalent*

**B1:**  $G(X) = G(Y)$

**B2:**  $X \vee (\bigvee_{\tau'(X)} S^{n_i}) \simeq Y \vee (\bigvee_{\tau'(Y)} S^{m_j})$

It was proved in [15] that in the 1-connected case A1 implies A2 and B1 implies B2 (under the conditions stated). Since these proofs easily extend to the nilpotent case, we will not repeat the arguments here. In order to get the converse, we proceed very much in the way one would argue in an “abelian” situation [2, 4]. We show that if the homotopy endomorphism set  $[X, X]$  has a suitable “local” structure, then  $X$ -factors or  $X$ -summands may be cancelled. More precisely

**DEFINITION 1<sup>2)</sup>.** The set  $\text{End } X = [X, X]$  is called  $H$ -local (or  $\pi$ -local) if for all  $f, g \in \text{End } X$  which are not homotopy equivalences,  $H_* f + H_* g \neq \text{Id}$  in  $\text{End}(H_*(X; \mathbf{Z}))$  (or  $\pi_* f + \pi_* g \neq \text{Id}$  in  $\text{End}(\pi_* X)$  respectively).

For example it is obvious that  $S_p^n$  and  $K(\mathbf{Z}_p, n)$  are both  $\pi$ -local and  $H$ -local for all  $n \geq 1$  and all primes  $p$ . ( $\mathbf{Z}_p$  denotes the integers localized at  $(p)$ ).

**LEMMA 1.** *Let  $A$  be a retract of  $X \times Y$  and suppose that  $\text{End } A$  is  $\pi$ -local. Then  $A$  is a retract of  $X$  or  $Y$ .*

*Proof.* There are maps  $f: A \rightarrow X \times Y$  and  $g: X \times Y \rightarrow A$  such that  $g \circ f \simeq \text{Id}_A$ . Let  $f_X = pr_X \circ f$ ,  $f_Y = pr_Y \circ f$  and  $g_X = g \mid X$ ,  $g_Y = g \mid Y$ . Then  $\pi_*(g_X \circ f_X) + \pi_*(g_Y \circ f_Y) = \text{Id}$ . Hence, since  $\text{End } A$  is  $\pi$ -local,  $g_X \circ f_X$  or  $g_Y \circ f_Y$  must be a homotopy equivalence and therefore  $A$  is a retract of  $X$  or  $Y$ .

Dually one has

**LEMMA 2.** *Let  $A$  be a retract of  $X \vee Y$  and suppose that  $\text{End } A$  is  $H$ -local. Then  $A$  is a retract of  $X$  or  $Y$ .*

The proof is completely analogous to that of Lemma 1, with  $\pi_*$  replaced by  $H_*$ .

**DEFINITION 2.**  $X$  is called irreducible, if  $X$  has no non-trivial retracts.

<sup>2)</sup>  $H_*$  denotes reduced homology.

Examples of irreducible spaces are  $S^n$ ,  $S_p^n$ ,  $K(\mathbf{Z}, n)$ ,  $K(\mathbf{Z}_p, n)$ ,  $CP^n$ ,  $HP^n$ ,  $BS^3$ ,  $SU(r)$  etc.

**DEFINITION 3.**  $X$  is called completely reducible (resp. completely coreducible) if  $X$  is a finite product (resp. finite wedge) of irreducible spaces.

Notice that if  $X$  is an  $H$ -complex and  $Y$  a retract of  $X$  with retraction map  $\varphi: X \rightarrow Y$ , then  $X \simeq \text{fib}(\varphi) \times Y$  and  $Y$  as well as the "fiber"  $\text{fib}(\varphi)$  are  $H$ -spaces. Hence, if in addition  $X$  is quasi-finite, one arrives after finitely many steps at a decomposition  $X \simeq X_1 \times \cdots \times X_n$  with  $X_i$  irreducible for  $1 \leq i \leq n$ . By essentially the same argument we see that for a quasi-finite  $H$ -complex  $X$  the  $p$ -localization  $X_p$  is completely reducible. Similarly, if  $Y$  is a quasi-finite  $H'$ -complex then  $Y$  and  $Y_p$  are both completely coreducible.

**DEFINITION 4.**  $X$  is called balanced if for all  $f, g \in \text{End } X$  with  $f \circ g$  a homotopy equivalence both  $f$  and  $g$  are homotopy equivalences.

If  $X$  is nilpotent and of finite type (i.e.  $\pi_i X$  or equivalently  $H_i X$  is finitely generated for all  $i$ ) then  $X$  is balanced. This follows easily by checking induced maps in homology. Similarly, if  $X$  is nilpotent and of finite type, then  $X_p$  is balanced for  $p$  a prime of 0.

**PROPOSITION 1.** *Let  $Y$  be a completely reducible space and  $A$  a balanced space with  $\text{End } A$  a  $\pi$ -local set. Then  $X \times A \simeq Y \times A$  implies  $X \simeq Y$ .*

*Proof.* For a map  $g: U \times V \rightarrow S \times T$  we write  $g(U, T): U \rightarrow T$  for  $pr_T \circ g \circ in_U$ . Let  $f: X \times A \rightarrow Y \times A$  be a homotopy equivalence. We will distinguish two cases

1)  $f(A, A)$  is a homotopy equivalence. Then one has in homotopy

$$\begin{aligned} f_*(A, A) \circ f_*^{-1}(Y, A) + f_*(X, A) \circ f_*^{-1}(Y, X) &= 0 \\ f_*(A, Y) \circ f_*^{-1}(Y, A) + f_*(X, Y) \circ f_*^{-1}(Y, X) &= Id \end{aligned}$$

Hence

$$(f_*(X, Y) - f_*(A, Y) (f_*(A, A))^{-1} f_*(X, A)) \circ f_*^{-1}(Y, X) = Id.$$

Similarly

$$f_*^{-1}(Y, X) \circ (f_*(X, Y) - f_*(A, Y) (f_*(A, A))^{-1} f_*(X, A)) = Id.$$

It follows that  $f_*^{-1}(Y, X): Y \rightarrow X$  is a homotopy equivalence.

2)  $f(A, A)$  is not a homotopy equivalence. Then, since  $A$  is a balanced space,  $f_*^{-1}(A, A) \circ f_*(A, A)$  is not a homotopy equivalence. But  $f_*^{-1}(A, A) \circ f_*(A, A) + f_*^{-1}(Y, A) \circ f_*(A, Y) = Id$  and  $\text{End } A$  is  $\pi$ -local. Therefore  $f_*^{-1}(Y, A) \circ f_*(A, Y)$  is a homotopy equivalence and we conclude that  $A$  is a retract of  $Y$ . Now  $Y$  is completely reducible:  $Y \simeq Y_1 \times \cdots \times Y_n$  with  $Y_i$  irreducible  $1 \leq i \leq n$ . It follows by

Lemma 1 and induction that  $A$  is a retract of one of the  $Y_i$ 's. More precisely, for some  $i_0$ ,  $\varphi = pr_{i_0} \circ f(A, Y): A \rightarrow Y_{i_0}$  has a homotopy left inverse. Since  $Y_{i_0}$  is irreducible, we conclude that  $\varphi$  is actually a homotopy equivalence. We consider now the automorphism

$$\theta: Y \times A \rightarrow Y \times A, \theta(y_1, \dots, y_{i_0}, \dots, y_n, a) = (y_1, \dots, \varphi a, \dots, y_n, \varphi^{-1} y_{i_0})$$

and form the new homotopy equivalence  $f' = \theta \circ f$ . It is immediate that  $f'(A, A) = \varphi^{-1} \circ \varphi = Id_A$ . Hence, by replacing  $f$  by  $f'$ , we are back in case 1.

If we replace, in this proof, induced maps in homotopy by induced maps in homology and using the nilpotency of the spaces involved, we will get the following dual proposition.

**PROPOSITION 2.** *Let  $Y$  be a completely coreducible space and  $A$  a balanced space with  $\text{End} A$  an  $H$ -local set. Then  $X \vee A \simeq Y \vee A$  implies  $X \simeq Y$ .*

*Remark.* In the proofs of Lemma 1 and 2 as well as Proposition 1 we made no use of the nilpotency of the spaces involved. Therefore these results remain true if the spaces are only assumed to be pointed connected CW-complexes. However for Proposition 2 and the following applications we have to restrict to nilpotent spaces.

**COROLLARY 1.** *Let  $Y$  be a finite product of quasi-finite  $H$ -complexes and spheres. Then  $G(X \times S^n) = G(Y \times S^n)$  implies  $G(X) = G(Y)$ .*

*Proof.* Suppose  $G(X \times S^n) = G(Y \times S^n)$ . Then  $X_p \times S_p^n \simeq Y_p \times S_p^n$  with  $Y_p$  completely reducible,  $\text{End} S_p^n$   $\pi$ -local and  $S_p^n$  balanced. Hence we conclude by Proposition 1 that  $X_p \simeq Y_p$  for all primes  $p$ .

**COROLLARY 2.** *A2 implies A1 in Theorem A.*

Namely one has only to observe that A2 implies  $\tau(X) = \tau(Y)$ , by checking rational cohomology. Then the result follows by iterated application of Corollary 1.

Similarly one has in the dual situation

**COROLLARY 3.** *Let  $Y$  be a quasi-finite  $H'$ -complex. If  $G(X \vee S^n) = G(Y \vee S^n)$  for some  $n > 1$ , then  $G(X) = G(Y)$ .*

**COROLLARY 4.** *B2 implies B1 in Theorem B.*

Notice that Theorem B is trivial in case  $Y$  is not 1-connected, because then  $Y \simeq S^1$ . We can also use Proposition 1 to find new  $H$ -complexes. For instance one has:

**COROLLARY 5.** *Let  $Y$  be a quasi-finite  $H$ -complex. Then  $X \times S^{n_1} \times \dots \times S^{n_k} \simeq Y \times S^{n_1} \times \dots \times S^{n_k}$  implies that  $X$  is a quasi-finite  $H$ -complex.*

*Proof.* Corollary 1 implies  $G(X) = G(Y)$  or  $X \in G(Y)$ . Notice that  $X$  is of finite type. Since  $Y$  is a quasi-finite  $H$ -complex, the same is true for  $X$  (cf [7]).

It is known [3] that for a finite 1-connected  $H$ -complex  $X$  there can be only finitely many finite  $H$ -complexes  $Y \in G(X)$ . The following examples should illustrate that the set  $G(X)$  can still be quite big.

EXAMPLE 1 (compare also [10] and [16]). Let  $p$  denote a prime and consider the principal fibration  $SU(p) \rightarrow SU(p+1) \rightarrow S^{2p+1}$  classified by  $\gamma = \alpha + \beta \in \pi_{2p}SU(p)$ , where  $\beta$  is of order  $p$  and  $\alpha$  of order prime to  $p$ . Define  $X_n$  as the total space of the principal  $SU(p)$ -fibration over  $S^{2p+1}$  classified by  $\alpha + n\beta$ . Then it is easily seen that  $X_n \simeq X_m$  if and only if  $n \equiv \pm m(p)$  and, if  $n \not\equiv 0(p)$  then  $X_n \in G(SU(p+1))$ . Hence we conclude

$$|G(SU(p+1))| \geq \frac{p-1}{2}$$

and, by Theorem A,  $X_n \times S^3 \times S^5 \times \dots \times S^{2p+1} \simeq SU(p+1) \times S^3 \times S^5 \times \dots \times S^{2p+1}$  if  $n \not\equiv 0(p)$ .

EXAMPLE 2 (compare also [12]). Let  $p$  be a prime and  $Y_m = S^3 \bigcup_{m\gamma} e^{2p+1}$  where  $\gamma \in \pi_{2p}S^3$  is of order  $p$ . Then  $Y_m$  is a  $H'$ -space since  $\gamma$  is a primitive element. Furthermore it is easy to see that  $Y_m \simeq Y_n$  if and only if  $m \equiv \pm n(p)$ . Hence

$$|G(Y_1)| \geq \frac{p-1}{2}$$

and, by Theorem B,  $Y_1 \vee S^3 \vee S^{2p+1} \simeq Y_m \vee S^3 \vee S^{2p+1}$  if  $m \not\equiv 0(p)$ .

## REFERENCES

- [1] BOUSFIELD, A. K. and KAN, D. M., *Homotopy limits, completions and localizations*, Lecture Notes in Mathematics 304, Springer Verlag (1972).
- [2] COHEN, J. M., *Stable homotopy*, Lecture Notes in Mathematics 165, Springer Verlag (1970).
- [3] CURJEL, C. R. and DOUGLAS, R. R., *On  $H$ -spaces of finite dimension*, *Topology* 10 (1971), 385–389.
- [3] FREYD, P., *Stable homotopy*. Proceedings of the conference on categorical algebra (La Jolla, 1965). Springer Verlag (1966).
- [5] HILTON, P. J., *On the Grothendieck group of compact polyhedra*, *Fund. Math.* 61 (1967), 199–214.
- [6] HILTON, P. J. and ROITBERG, J., *On principal  $S^3$ -bundles over spheres*, *Ann. of Math.* 90 (1969), 91–107.
- [7] HILTON, P. J., MISLIN, G. and ROITBERG, J. (to appear).
- [8] —,  *$H$ -spaces of rank 2 and non-cancellation phenomena*, *Inv. Math.* 16 (1972), 325–334.
- [9] —, *Sphere bundles over spheres and non-cancellation phenomena*, *J. Lond. Math. Soc.* (2), 6 (1972), 15–23.

- [10] MISLIN, G., *H-spaces mod  $p$* , *H-spaces conference*, Neuchatel 1970, Lecture Notes in Mathematics 196, Springer Verlag (1971).
- [11] —, *The genus of an H-space*, Lecture Notes in Mathematics 249, Springer Verlag (1971), 75–83.
- [12] MOLNAR, E. A., *Relation between wedge cancellation and localization for complexes with two cells*, *J. Pure Appl. Algebra* 3 (1973), 141–158.
- [13] SIERADSKI, A., *Square roots up to homotopy type*, *Amer. J. Math.* 94 (1972), 73–81.
- [14] SULLIVAN, D., *Geometric Topology, part I: Localization, periodicity and Galois symmetry*, MIT, June 1970, (mimeographed notes).
- [15] ZABRODSKY, A., *On the genus of finite CW H-spaces*, to appear.
- [16] —, *On the homotopy type of principal classical group bundles over spheres*, *Israel J. Math.* 11 (3) 1972, 315–325.

*E.T.H. Leonard-Str.*  
8006 Zürich

Received November 28, 1973