

# On the Covariant Differential of an Almost Hermitian Structure.

Autor(en): **Terrier, J.M.**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **50 (1975)**

PDF erstellt am: **11.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-38800>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## On the Covariant Differential of an Almost Hermitian Structure

J. M. TERRIER

This paper deals with the covariant differential  $\nabla J$  of an almost hermitian structure  $J$  on a Riemannian manifold  $M$  with metric  $g$ . The connection with respect to which  $\nabla J$  is defined is the Riemannian connection on  $M$ . The extension of the notion of antisymmetrization and symmetrization from tensor fields of type  $(0, q)$  to those of type  $(p, q)$  for  $p > 0$  allows us to decompose  $\nabla J$  in an antisymmetric part  $A$  and a symmetric part  $S$ . In this way, we find new formulas which have the feature of stressing the relationship between  $\nabla J$  and the torsion  $\tau$  of  $J$  as well as the fundamental 2-form  $\omega$  or better, its exterior derivative  $d\omega$ . The results are essentially based first on the Palais formula which gives the exterior derivative of a  $q$ -form via Lie product and covariant derivative and second on a theorem ([2], p. 149) which gives the exterior derivative  $d\alpha$  of a  $q$ -form  $\alpha$  as the antisymmetric part of the covariant differential  $\nabla\alpha$  of  $\alpha$ , provided the connection has vanishing torsion. As an application of our formulas we give a characterization of so called nearly Kähler manifolds ([1]) via the fundamental 2-form  $\omega$ . We also give a very simple proof of the characterization of a Kähler manifold given by the vanishing of  $\nabla J$  or of  $d\omega$  and  $\tau$ . We finally prove a lemma which gives a nice interpretation of the torsion of  $J$  when the fundamental 2-form is closed, that is, in the case of an almost Kähler manifold.

### §1. The Covariant Differential of a Tensor Field

Let  $t$  be a given tensor field of type  $(p, q)$  on a  $C^\infty$  manifold  $M$ . We shall simply write  $t \in T_M(p, q)$  or  $t \in T(p, q)$ .

Suppose there is also a linear connection  $\nabla$  given on  $M$ . Then, as in [2], we can define the covariant differential of  $t$  as the tensor field  $\nabla t \in T(p, q+1)$  defined by

$$\nabla t(X_1, \dots, X_q, X) = (\nabla_X t)(X_1, \dots, X_q) \tag{1.1}$$

where  $\nabla_X t$  denotes the covariant derivative of the tensor field  $t$  and  $X_1, \dots, X_q, X$  are in the Lie algebra  $\mathfrak{X}(M)$  of vector fields on  $M$ .

Because  $\nabla_X$  is a derivation commuting with every contraction, we have

**THEOREM 1.1** ([2], p. 124). *If  $t \in T(p, q)$  then, for  $X_i, X \in \mathfrak{X}(M)$*

$$\nabla t(X_1, \dots, X_q, X) = \nabla_X(t(X_1, \dots, X_q)) - \sum_{k=1}^q t(X_1, \dots, \nabla_{X_i} X_k, \dots, X_q).$$

EXAMPLE 1. Take  $t=g$  a Riemann metric on  $M$ .  $g$  is in  $T(0, 2)$ , so  $\nabla g$  is in  $T(0, 3)$  and for  $X, Y, Z, \in \mathfrak{X}(M)$  we have

$$\nabla g(X, Y, Z) = (\nabla_Z g)(X, Y) = Zg(X, Y) - g(\nabla_Z X, Y) - g(X, \nabla_Z Y).$$

One of the features of a Riemannian metric  $g$  is to be parallel i.e.  $\nabla g = 0$ . For convenience we shall write  $\langle X, Y \rangle$  instead of  $g(X, Y)$ .

EXAMPLE 2. Take  $t=J$  an almost complex structure on  $M$ . This is a tensor field of type  $(1, 1)$  whose square  $J^2$  equals minus the identity.  $\nabla J$  is in  $T(1, 2)$  and

$$\nabla J(X, Y) = (\nabla_Y J)X = \nabla_Y(JX) - J\nabla_Y X.$$

One of the features of a Kähler structure  $J$  on  $M$  is to be parallel with respect to  $\nabla$ :  $\nabla J = 0$ .

## §2. Extension of Antisymmetrization and Symmetrization

It is known [1] how to define for a covariant tensor field  $t$  in  $T(0, q)$  the *alternation*  $At$  of  $t$ . It is a tensor field of the same type defined by

$$(At)(X_1, \dots, X_q) = \frac{1}{q!} \sum_{\pi \in P_q} \varepsilon(\pi) t(X_{\pi(1)}, \dots, X_{\pi(q)}) \quad (2.1)$$

where  $P_q$  is the group of permutations of  $\{1, 2, \dots, q\}$  and  $\varepsilon(\pi)$  is the signe of the permutation  $\pi$ . Notice that  $At$  is a skew-symmetric tensor field and  $t$  is skew-symmetric if and only if  $At = t$ .

On the other hand one also defines for  $t \in T(0, q)$  the *symmetrization*  $St$  of  $t$  by

$$St(X_1, \dots, X_q) = \frac{1}{q!} \sum_{\pi \in P_q} t(X_{\pi(1)}, \dots, X_{\pi(q)}). \quad (2.2)$$

Here is  $St$  a symmetric tensor field and  $t$  is symmetric if and only if  $St = t$ .

We now make the straightforward extension of the above notions to tensor fields of type  $(p, q)$  for  $p > 0$ . In this case  $t(X_1, \dots, X_q)$  is no longer a real function on  $M$ , but a  $p$ -contravariant tensor field. Nevertheless, all algebraic operations needed for definitions (2.1) and (2.2) still make sense in the module of  $p$ -contravariant tensor fields.

EXAMPLE 1. Take  $t=\tau \in T(1, 2)$  the torsion of an almost complex structure  $J$  defined by

$$\tau(X, Y) = [X, Y] + J[JX, Y] + J[X, JY] - [JX, JY].$$

Here we have  $A\tau = \tau$  and  $S\tau = 0$ .

EXAMPLE 2. Take  $t = K \in T(1, 3)$  where  $K(X, Y, Z) = R(X, Y)Z$  and  $R \in T(1, 3)$  is defined by  $R(X, Y) = [\nabla_X, \nabla_Y] - \nabla_{[X, Y]}$ .

Because  $R(X, Y) = -R(Y, X)$  we have

$$(AK)(X, Y, Z) = \frac{1}{3}\xi(K(X, Y, Z))$$

( $\xi$  means cyclic sum) and  $SK = 0$ . Furthermore if the torsion of  $\nabla$  vanishes, we have  $AK = 0$  in view of the first Bianchi identity.

### §3. Almost Hermitian Manifolds

From now on, we assume that  $(M, J, g)$  is an almost hermitian manifold, that is, the almost complex structure  $J$  on  $M$  gives in each point  $p$  of  $M$  an isometry of  $T_p(M)$ , the tangent space at  $M$  in  $p$ , i.e.

$$\langle JX, JY \rangle = \langle X, Y \rangle \quad \forall X, Y \in T_p(M). \tag{3.1}$$

Denoting by  $\nabla$  the Riemannian connection on  $M$ , we can compute  $\nabla J$  which is a tensor field of type  $(1, 2)$  and we can write the following decomposition

$$\nabla J = A(VJ) + S(VJ). \tag{3.2}$$

Needless to say, this is only possible because we are in  $T(p, q)$  with  $q = 2$ . Example 2 above shows what can happen with  $q \neq 2$ .

For convenience we shall write  $A$  for  $A(\nabla J)$  and  $S$  for  $S(\nabla J)$  and establish several formulas relating  $A$  and/or  $S$  with various tensor fields one can define on an almost hermitian manifold.

#### I. The Torsion $\tau$ and $A$

THEOREM 3.1. *If  $A$  denotes the antisymmetric part of the covariant differential  $\nabla J$  of the almost hermitian structure  $J$  on  $M$ , then*

$$\frac{1}{2}J\tau(X, Y) = A(X, Y) - A(JY, JY). \tag{3.3}$$

*Proof.* By definition  $A(X, Y) = \frac{1}{2}\{\nabla J(X, Y) - \nabla J(Y, X)\}$ . So

$$2\{A(X, Y) - A(JX, JY)\} = \nabla J(X, Y) - \nabla J(Y, X) - \nabla J(JX, JY) + \nabla J(JY, JX).$$

In view of (1.1), the right hand side becomes

$$(\nabla_Y J)X - (\nabla_X J)Y - (\nabla_{JY} J)(JX) + (\nabla_{JX} J)(JY).$$

which is, according to Theorem 1.1, the same as

$$\nabla_Y JX - J\nabla_Y X - \nabla_X JY + J\nabla_X Y + \nabla_{JY} X + J\nabla_{JY} JX - \nabla_{JX} Y - J\nabla_{JX} JY. \quad (3.4)$$

The Riemannian connection having torsion zero, we have  $\nabla_X Y - \nabla_Y X = [X, Y]$  and multiplication of (3.4) by  $J$  gives (3.3).

From Theorem 3.1 we trivially get the following

**COROLLARY 3.2.** *If the covariant differential  $\nabla J$  of an almost hermitian structure  $J$  is symmetric then  $J$  is integrable.*

## II. The Fundamental 2-form $\omega$ and $A$

The fundamental 2-form  $\omega$  on an almost hermitian manifold  $(M, J, g)$  is defined by

$$\omega(X, Y) = \langle JX, Y \rangle \quad \text{for } X, Y \in \mathfrak{X}(M). \quad (3.5)$$

The torsion of the Riemann connection being zero, we know ([2], chap. III) that  $d\omega = A(\nabla\omega)$ . For vector fields  $X, Y$ , and  $Z$  on  $M$ , we have therefore:

$$6d\omega(X, Y, Z) = 6A(\nabla\omega)(X, Y, Z) = \mathfrak{s}(\nabla\omega(X, Y, Z)) - \mathfrak{s}(\nabla\omega(Y, X, Z)) \quad (3.6)$$

where  $\mathfrak{s}$  denotes cyclic sum. By definition of the covariant differential and (3.5) we have

$$\begin{aligned} \nabla\omega(X, Y, Z) &= (\nabla_Z\omega)(X, Y) = Z\omega(X, Y) - \omega(\nabla_Z X, Y) - \omega(X, \nabla_Z Y) \\ &= \langle \nabla_Z JX, Y \rangle + \langle JX, \nabla_Z Y \rangle - \langle J\nabla_Z X, Y \rangle - \langle JX, \nabla_Z Y \rangle \\ &= \langle \nabla_Z JX, Y \rangle - \langle J\nabla_Z X, Y \rangle = \langle (\nabla_Z J)X, Y \rangle = \langle \nabla J(X, Z), Y \rangle. \end{aligned}$$

Hence

$$\nabla\omega(X, Y, Z) = \langle \nabla J(X, Z), Y \rangle \quad (3.7)$$

In view of (2.1) and (3.2) we have the following

**THEOREM 3.3.** *On an almost hermitian manifold  $(M, J, g)$  the exterior differential  $d\omega$  of the fundamental 2-form  $\omega$  and the antisymmetric part of the covariant differential  $\nabla J$  are related by the formula*

$$3d\omega(X, Y, Z) = -\mathfrak{s}(\langle A(X, Y), Z \rangle). \quad (3.8)$$

Together with Corollary 3.2 this result implies

**COROLLARY 3.4.** *If the covariant differential  $\nabla J$  of an almost hermitian structure*

*J is symmetric, then the fundamental 2-form  $\omega$  is closed and J is integrable, i.e.  $(M, J, g)$  is a Kähler manifold.*

There is another interesting relation between  $d\omega$ ,  $A$  and  $\nabla J$ ; namely, we have

**THEOREM 3.5.** *On any almost hermitian manifold  $(M, J, q)$  we have for the fundamental 2-form  $\omega$ :*

$$3d\omega(X, Y, Z) = -2\langle A(X, Y), Z \rangle + \langle \nabla J(X, Z), Y \rangle. \quad (3.9)$$

*Proof.* By the Palais formula, one has for any 2-form  $\omega$

$$3d\omega(X, Y, Z) = X\omega(Y, Z) - Y\omega(X, Z) + Z\omega(X, Y) - \omega([X, Y], Z) + \omega([X, Z], Y) - \omega([Y, Z], X).$$

By the definition of  $\omega$  and because  $g$  is parallel with respect to  $\nabla$ , together with the fact, once again, that the torsion of  $\nabla$  vanishes, we get the stated result.

### III. A Relation Between $A$ and $S$ and an Identity for $S$

From Theorem 3.3 and Theorem 3.5 above we deduce the

**COROLLARY 3.6.** *If  $A$  (resp.  $S$ ) is the antisymmetric (resp. symmetric) part of the covariant differential  $\nabla J$  of an almost hermitian structure then, for vector fields  $X, Y, Z$  on  $M$ , we have*

$$\langle S(X, Y), Z \rangle = \langle A(X, Z), Y \rangle + \langle A(Y, Z), X \rangle. \quad (3.10)$$

*Proof.* From (3.2), (3.8) and (3.9) one has

$$\mathfrak{s}(\langle A(X, Y), Z \rangle) = 2\langle A(X, Y), Z \rangle + \langle A(X, Z), Y \rangle + \langle S(X, Z), Y \rangle.$$

But  $A$  is antisymmetric, so  $A(X, Z) = -A(Z, X)$  and (3.10) follows by permuting  $Y$  and  $Z$ .

The antisymmetry of  $A$  has another consequence:

**COROLLARY 3.7.** *The symmetric part  $S$  of the covariant differential  $\nabla J$  of an almost hermitian structure satisfies the following identity*

$$\mathfrak{s}(\langle S(X, Y), Z \rangle) = 0 \quad (3.11)$$

where  $\mathfrak{s}$  denotes cyclic sum.

*Proof.* Write the left hand side of (3.11) with (3.10) and use the antisymmetry of  $A$ .

#### §4. A few Remarks

1. Because of (3.11) we can rewrite Theorem 3.3 in the following form:

**THEOREM 3.3'.** *On an almost hermitian manifold  $(M, J, g)$  the exterior differential  $d\omega$  of the fundamental 2-form  $\omega$  and the covariant differential  $\nabla J$  of  $J$  are related by the formula*

$$3d\omega(X, Y, Z) = -s(\langle \nabla J(X, Y), Z \rangle). \quad (4.1)$$

2. An almost hermitian manifold  $(M, J, g)$  for which  $S=0$  is already known [1] as *nearly Kähler manifold*. An alternative condition is given by the following

**THEOREM 4.1.** *A nearly Kähler manifold  $(M, J, g)$  is characterized by the condition*

$$\nabla\omega = d\omega. \quad (4.2)$$

*Proof.* We have to show that this condition is equivalent to  $S=0$ . Suppose (4.2) is true. Then  $\nabla\omega$  is antisymmetric. By (3.7)  $\nabla\omega(X, Y, Z) = \langle \nabla J(X, Z), Y \rangle$  and  $\nabla\omega(Z, Y, X) = \langle \nabla J(Z, X), Y \rangle = -\langle \nabla J(X, Z), Y \rangle$  which implies that  $\nabla J$  itself is antisymmetric, i.e.  $S=0$ .

On the other hand,  $S=0$  implies  $\nabla J = A$  and by (3.10)  $\langle \nabla J(X, Z), Y \rangle + \langle \nabla J(Y, Z), X \rangle = 0$  which in turn gives by antisymmetry of  $\nabla J$ :  $\langle \nabla J(Z, X), Y \rangle + \langle \nabla J(Z, X), Y \rangle = 0$  or with (3.7),  $\nabla\omega(Z, Y, X) + \nabla\omega(Z, X, Y) = 0$ . But (even if  $S \neq 0$ ) one has  $\nabla\omega(X, Y, Z) + \nabla\omega(Y, X, Z) = 0$ , and (3.6) gives the result.

3. Of the antisymmetric part  $A$  and the symmetric part  $S$  of  $\nabla J$ , the former plays the most important role. It allows us to give as an application a very simple proof of the following theorem. Compare with [3] (chap. IX).

**THEOREM 4.2.** *An almost hermitian manifold  $(M, J, g)$  is a Kähler manifold (i.e.  $\tau=0$  and  $d\omega=0$ ) if and only if the covariant differential  $\nabla J$  vanishes.*

*Proof.* If  $\nabla J=0$  then  $A=0$  which implies  $\tau=0$  by (3.3) and  $d\omega=0$  by (3.8). On the other hand, from Lemma 4.3 below and  $d\omega=0$  we get  $A=0$  and by (3.10),  $S=0$ .

**LEMMA 4.3.** *For an almost Kähler manifold, the antisymmetric part  $A$  of the covariant differential  $\nabla J$  is essentially the torsion of  $J$ : more precisely we have*

$$4A = J\tau.$$

*Proof.*  $d\omega=0$  implies by (3.9)

$$\langle \nabla J(X, Z), Y \rangle = 2\langle A(X, Y), Z \rangle.$$

Substituting  $JX$  to  $X$  and  $JY$  to  $Y$  and adding, we get

$$\langle \nabla J(X, Z), Y \rangle + \langle \nabla J(JX, Z), JY \rangle = 2 \langle A(X, Y) + A(JX, JY), Z \rangle. \quad (4.3)$$

But the left hand side vanishes because  $\nabla J(JX, Y) = -J\nabla J(X, Y)$  as it is easy to see and  $J$  is an isometry. To get the desired result one just has to add (4.3) to (3.3).

#### REFERENCES

- [1] GRAY A., *Nearly Kähler manifolds*, J. Differential Geometry 4 (1970), 283–309.
- [2] KOBAYASHI S. and NOMIZU K., *Foundations of Differential Geometry*, Vol. I. Interscience Publishers, New York, 1963.
- [3] —, *Foundations of Differential Geometry*, Vol. II. Interscience Publishers, New York, 1969.

*Department of Mathematics  
University of Montreal  
Montreal, Canada*

*Forschungsinstitut für Mathematik  
E.T.H. Zürich Switzerland*

Received September 23, 1974.



