Zeitschrift:	Commentarii Mathematici Helvetici
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	57 (1982)
Artikel:	An algebro-geometric interpretation of the Bäcklund-transformation for
	the Koneweg-de vites equation.
Autor:	Ehlers, Fritz / Knörrer, Horst

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 01.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

An algebro-geometric interpretation of the Bäcklundtransformation for the Korteweg-de Vries equation

FRITZ EHLERS and HORST KNÖRRER

1. Introduction

In 1882 G. Darboux has developed a method that associates to any onedimensional Schrödinger-operator $Z = -d^2/dx^2 + u(x)$ infinitely many other such operators the spectra of which are intimately related to the spectrum of Z itself (cf. [4] §408). This transformation is defined in the following way: Let c be a constant, let $\psi(x)$ be a solution of the differential equation $Z \cdot \psi = c\psi$

with $\psi \neq 0$ and put $v := \psi_x/\psi$. One easily verifies that $u = v^2 + v_x + c$, and we put

$$\tilde{u}:=v^2-v_x+c=u-2v_x.$$

The differential operator

$$\tilde{Z}:=-\frac{d^2}{dx^2}+\tilde{u}$$

is called the Darboux-Bäcklund-transform of Z. It depends on the choice of c and ψ , a change of ψ by a constant factor does not effect the transformation.

It is well known that Korteweg-de Vries-equation

$$u_t = 3uu_x - \frac{1}{2}u_{xxx}$$

is equivalent to the Lax-equation

$$\frac{\partial Z}{\partial t} = -[K, Z]$$

with

$$Z := -\frac{d^2}{dx^2} + u, \qquad K := 2\frac{d^3}{dx^3} - 3u\frac{d}{dx} - \frac{3}{2}u_x$$

(For a discussion of the properties of the KdV-equation cf. [10]).

The transformation of Z described above is compatible with this Laxequation, so it yields a transformation that produces new solutions of the KdV-equation out of given ones. Transformations of this type are called Bäcklund-transformations after A. Bäcklund who developed such a transformation for the sine-Gordon-equation in the last century (cf. [8]). For the KdVequation the transformation described above has been worked out by R. Miura [13].

Successive application of the Bäcklund-transformation with diminishing values of $c \in \mathbb{R}$, c < 0, produces out of the solution $u \equiv 0$ of the KdV-equation the multisoliton-solutions of this equation.

In 1976 I. M. Krichever developed a method for constructing one-dimensional Schrödinger operators starting with certain line bundles on hyperelliptic curves (cf. [12], [14]; parts of this construction will be described in chapter 2). Since Krichever's construction is purely formal we work with differential operators with coefficients in the ring $\mathbb{C}[[x]]$ of complex formal power series in x (i.e. with elements of $\mathbb{C}[[x]][d/dx]$). The object of this paper is to describe how the Darboux-Bäcklund-transformation, applied to such a Schrödinger operator, changes the hyperelliptic curve and the line bundle on it. The precise result will be given in Chapter 3; its essential content is the following:

Let Z be a Schrödinger-operator constructed by Krichever's method from a line bundle on the hyperelliptic curve $y^2 + F(z) = 0$ (F a polynomial of odd degree), and let \tilde{Z} be a Bäcklund-transform of Z as described above. If ψ was chosen generically⁽¹⁾ in the c-eigenspace of the linear map $Z: C[[x]] \rightarrow C[[x]]$ then \tilde{Z} corresponds to a line bundle on the curve $y^2 + (z-c)^2 F(z) = 0$, whereas for some special choices of ψ the operator \tilde{Z} can be obtained from a line bundle on the curve $y^2 + F(z) = 0$ or $y^2 + (z-c)^{-2}F(z) = 0$.

Our work on this problem was motivated by discussions in a seminar on geometrical methods in mathematical physics, run by members and guests of the SFB 40 in Bonn in 1979 and 1980. We would like to thank all its participants, especially M. Adler, for many helpful and interesting suggestions.

2. Krichever's construction, as described by Mumford

Krichever's construction provides a relation between certain line bundles on hyperelliptic curves (see the end of this chapter) and certain Schrödinger

¹ The notion "generically" has the meaning used in algebraic geometry. As the eigenspaces of Z are two-dimensional it means that ψ has to avoid some finite number of one-dimensional subspaces.

operators. These Schrödinger operators have the property that they commute with some other differential operator Y of odd degree.⁽²⁾ Conversely, any Schrödinger operator with this property can be obtained from a line bundle on some hyperelliptic curve. In this chapter we will describe how the hyperelliptic curve and the line bundle on it can be reconstructed from the operator Z. In the exposition of these results we follow the approach of Mumford [14], §2.

Let Y be a differential operator of odd degree commuting with Z, and assume further that its degree r is minimal with respect to this property. It follows from the commutativity that its leading coefficient is constant and thus we may assume that it equals one. Y and Z generate the ring $R \subset \mathbb{C}[[x]][d/dx]$ of all differential operators commuting with Z. As in [14], p. 133 one sees that there are polynomials $F_1(z), F_2(z) \in \mathbb{C}[z]$ such that $Y^2 + 2F_1(Z) \cdot Y + F_2(Z) = 0$, deg $F_1 \leq (r-1)/2$, deg $F_2 = r$. Replacing Y by $Y + F_1(Z)$ one obtains $Y^2 + F(Z) = 0$ where $F(z) := F_2(z) - F_1(z)^2$ has leading coefficient one. It can be shown that

 $\mathbb{C}[z, y]/(y^2 + F(z)) \rightarrow R, \quad z \mapsto Z, \quad y \mapsto Y$

is an isomorphism.

Let C_0 be the affine curve $\{(x, y) \in C^2 \mid y^2 + F(X) = 0\}$. This is the spectrum Spec R of the commutative ring R in the sense of algebraic geometry. There is a unique compact curve $C \supset C_0$, such that the points of $C - C_0$ are nonsingular on C. Since deg F is odd, $C - C_0$ consists only of one point, P.

In some neighbourhood of P, $\zeta := y \cdot z^{-(r+1)/2}$ is a local parameter of C. If $D \in R$ is considered as a rational function on C then its pole order at P is its order as a differential operator.

In order to construct the line bundle on C we notice that for every $z \in \mathbb{C}$ the operator Y maps the two-dimensional space ker $(Z - z \cdot id)$ to itself. Its eigenvalues on this space are the values $y \in \mathbb{C}$ with $y^2 + F(z) = 0$. So for each point $Q \in C_0$ the "eigenspace of R corresponding to Q"

$$E_{O} := \ker (Z - z \cdot id) \cap \ker (Y - y \cdot id)$$

has dimension ≥ 1 .

Remark. dim $E_{O} = 1$ for all $Q \in C_{O}$.

Proof. Let Q = (z, y) be a point of C_0 . If E_0 were not one-dimensional then

² The existence of such an operator Y implies that u is a so called finite gap potential.

the two-dimensional space ker $(Z - z \cdot id)$ would be contained in ker $(Y - y \cdot id)$. This implies that $(Y - y \cdot id) = Y'$. $(Z - z \cdot id)$ with some operator Y' of degree r-2 (cf. [9], 5.4). Since $Y - y \cdot id$ and $Z - z \cdot id$ commute with Z the same is true for Y', in contradiction to the assumption that the degree of Y was minimal.

The spaces E_Q can be glued to a line bundle on the affine curve C_0 . In order to describe its extension to C we use the following construction indicated by Mumford [14]:

Consider $M := \mathbb{C}[d/dx]$ as a right *R*-module with multiplication

$$X \cdot D = (X \circ D)_0, \quad X \in M, \quad D \in R, \quad \left(\sum a_i(x) \frac{d^i}{dx^i}\right) = \sum a_i(0) \frac{d^i}{dx^i}$$

M is finitely generated, so it is the space of sections of some coherent sheaf \mathscr{L}_0 on C_0 , unique up to isomorphism. \mathscr{L}_0 extends to a sheaf \mathscr{L} on *C* in such a way that the space of sections of $\mathscr{L} \otimes \mathcal{O}(nP)$ corresponds to the space of differential operators of order at most *n*.

Mumford shows that the pairing

$$\begin{array}{l} M \times E_Q \to \mathbf{C} \\ (X, f) \mapsto (Xf)(0) \end{array} (Q \in C_0) \end{array}$$

induces an isomorphism

 $\operatorname{Hom}_{\mathbf{C}}(\mathfrak{L}_{\mathbf{Q}}/m_{\mathbf{Q}}\mathfrak{L}_{\mathbf{Q}},\mathbf{C})\cong E_{\mathbf{Q}}$

where m_0 denotes the maximal ideal of $Q \in C_0$.

Since E_Q is one-dimensional for all $Q \in C_0$ this implies that \mathscr{L} is locally free of rank 1, hence \mathscr{L} is the sheaf of sections of a line bundle L on C. (Obviously $L|_{C_0}$ is the dual of the line bundle obtained by glueing the E_Q 's.) By construction we have that dim $H^0(C, L(nP)) = \dim M_{n-1} = n$ for all $n \in \mathbb{N}$, by Riemann-Roch this is equivalent to dim $H^0(C, L) = \dim H^1(C, L) = 0$. Hence L is a line bundle of degree g-1, where g = (r-1)/2 is the arithmetic genus of C.

We have thus obtained the following algebraic-geometric data:

(i) a hyperelliptic curve C in normal form $y^2 + F(z) = 0^{(3)}$, where F is a polynomial of odd degree with leading coefficient one.

(ii) a line bundle L on C with dim $H^0(C, L) = \dim H^1(C, L) = 0$.

³ i.e. a hyperelliptic curve C together with two rational functions y, z on C that generate the function field of C and that fulfil the identity $y^2 + F(z) = 0$.

As we mentioned in the beginning one can recover the operator Z from these data. Furthermore it can be shown that the time-variation of the operator

$$Z_t = -\frac{d^2}{dx^2} + u(x, t)$$

according to the KdV-equation induces a linear variation of L in the Jacobian of line bundles of degree g-1 on C.

3. The Bäcklund-transformation in the algebraic-geometric picture

Let $Z = -(d^2/dx^2) + u(x)$ be a formal Schrödinger operator, and let R be the ring of all differential operators commuting with Z.

ASSUMPTION. R contains a differential operator of odd order.

As in the introduction let $\tilde{Z} = -(d^2/dx^2) + \tilde{u}(x)$ be Darboux-Bäcklund-transform of Z with parameters $c \in \mathbb{C}$, $\psi \in \ker (Z - c \cdot id)$, $\psi(0) \neq 0$. Denote by \tilde{R} the ring of differential operators commuting with \tilde{Z} .

ASSERTION. \tilde{R} also contains a differential operator of odd order.

As in Chapter 2 we associate to R resp. \tilde{R} hyperelliptic curves C and \tilde{C} with normal forms $y^2 + F(z) = 0$ resp. $y^2 + \tilde{F}(z) = 0$ and line bundles L and \tilde{L} on C. resp. \tilde{C} . Let \mathscr{L} resp. $\tilde{\mathscr{L}}$ be the sheaf of sections of L resp. \tilde{L} , and let P resp. \tilde{P} be the point at infinity of C resp. \tilde{C} . We call $E := \bigcup_{Q \in C-P} E_Q$ the set of common eigenfunctions of R.

THEOREM.

(I) If $\psi \notin E$ then \tilde{C} is the hyperelliptic curve in the normal form $y^2 + (z-c)^2 \cdot F(z) = 0$ (i.e. $\tilde{F}(z) = (z-c)^2 \cdot F(z)$). If $\pi: C \to \tilde{C}$ denotes the map induced by $(z, y) \mapsto (z, (z-c)y)$ then $\tilde{\mathcal{L}}$ is naturally isomorphic to the subsheaf of $\pi_* \mathcal{L} \otimes \mathcal{O}(\tilde{P})$ of sections s with $\langle s_0, \psi \rangle = 0$.

(ii) If $\psi \in E_Q$ for some smooth point $Q \in C$ then the curves C and \tilde{C} coincide (i.e. $F = \tilde{F}$) and $\tilde{L} \cong L \otimes \mathcal{O}(P) \otimes \mathcal{O}(-Q)$.

(iii) If $\psi \in E_Q$ for a singular point Q of C then \tilde{C} is the hyperelliptic curve with the normal form $y^2 + (z-c)^{-2}F(z) = 0$. If $\tau: \tilde{C} \to C$ denotes the map induced by $(z, y) \mapsto (z, (z-c)y)$ then $\tilde{L} \cong \tau^*L \otimes O(-\tilde{P})$. Remarks.

1. For almost all choices of $\psi \in \ker (Z - c \cdot id)$, ψ is not contained in *E*. In this case \tilde{L} is a line bundle on \tilde{C} with $\pi^*(\tilde{L}) \otimes \mathbb{O}(-P) \cong L$. By [15], V. 12 the Jacobian⁽⁴⁾ of \tilde{C} is a central extension of the Jacobian of *C*:

 $e \to G \to \operatorname{Jac}(\tilde{C}) \to \operatorname{Jac}(C) \to 0, \qquad \tilde{L} \to \pi^*(\tilde{L}) \otimes \mathbb{O}(-P)$

with $G = \mathbb{C}^*$ or $G = \mathbb{C}$, *e* the neutral element of *G*. Given a line bundle *L* on *C* with $H^0(L) = H^1(L) = 0$ we have an isomorphism between *G* and the set of elements $[\psi] \in \mathbb{P}(\ker (Z - c \cdot id))$ such that $\psi \notin E$. The condition $\psi(0) \neq 0$ corresponds to the condition $H^0(\tilde{L}) = H^1(\tilde{L}) = 0$.

2. If $\psi \in E_Q$ for a singular point $Q \in C$ (which then has the coordinates (c, 0)) the Bäcklund transformation from Z to \tilde{Z} is the inverse of a Bäcklund transformation from \tilde{Z} to Z as described in part (i) of the theorem and in the remark above.

3. The "transference" in Jac (C) caused by the Bäcklund transformation with $\psi \in E_Q$ for a smooth point $Q \in C$ has been considered in greater generality by Burchnall and Chaundy [3], §VI, VII.

4. J. Drach noticed in 1912 that the Darboux-Backfund transformation does not increase the effective genus of the associated hyperelliptic curve (see [6]).⁽⁵⁾

5. In [1] it is shown that iterated application of the Bäcklund transformation with c = 0 to the solution $u \equiv 0$ of the KdV-equation leads to the rational solutions discovered by Airault-McKean-Moser [2]. So the theorem shows that these solutions belong to curves of the form $y^2 + z^{2n+1} = 0$.

Similarly one can construct the multi-soliton-solutions by repeated application of the Bäcklund-transformation with diminishing values of c (cf. [5]). This gives a simple proof that these solutions belong to curves of the form $y^2 + (z-c_1)^2 \cdots (z-c_n)^2 z = 0$ (see also [11]).

6. For a real valued Schwartz function u(x) the effect of the Bäcklund-transformation on the spectral data has been described in [7].

4. Proofs

Replacing u by u-c we may assume that c=0. 4.1. The interpretation of the Bäcklund-transformation given in [1], [5] shows

⁴ By abuse of notation we denote by Jac (C) the moduli space of line bundles of degree g-1 on C. As base point we take the point at infinity.

⁵ This reference was shown to us by D. V. and G. V. Chudnovsky during the 1980 Arbeitstagung of the SFB 40 in Bonn.

that \tilde{Z} is obtained from Z by conjugation in the ring PsD[[x]] of pseudodifferential operators. (For the definition of PsD[[x]] see [14], p. 140.)

LEMMA 1. $\tilde{Z} = A \circ Z \circ A^{-1}$ where A is the differential operator

$$A:=\psi\circ\frac{d}{dx}\circ\psi^{-1}=\frac{d}{dx}-\frac{\psi_x}{\psi}=\frac{d}{dx}-v$$

Proof. Since $Z\psi = -\psi_{xx} + u\psi = 0$ we have $u = \psi_{xx}/\psi$ and hence $Z = A^* \circ A$ where $A^* = -(d/dx) - v = -\psi^{-1} \circ d/dx \circ \psi$ denotes the adjoint of A. Then one easily verifies that $\tilde{Z} = A \circ A^*$, which proves the lemma.

The ring R is generated by Z and some operator Y with $Y^2 + F(Z) = 0$. $\psi \in E$ if and only if ψ is an eigenvector of Y.

LEMMA 2. Let $\kappa: D \to A \circ D \circ A^{-1}$ be the conjugation by A in PsD[[x]]. (i) If $\psi \notin E$, let $R':= \mathbb{C}[Z, Y \circ Z] \subset R$. Then $\kappa(R) \cap \mathbb{C}[[x]][d/dx] = \kappa(R')$. (ii) $\kappa(R) \subset \mathbb{C}[[x]][d/dx]$ if and only if $\psi \in E$. In this case, call R' = R.

Proof. Let $X \in \mathbb{C}[[x]][d/dx]$ be any formal differential operator. Then

$$X \circ \psi = X\psi + X' \circ \frac{d}{dx}$$

where X' is some differential operator. It follows that

$$A \circ X \circ A^{-1} = \psi \circ \frac{d}{dx} \circ \psi^{-1} \circ X \psi \circ \left(\frac{d}{dx}\right)^{-1} \circ \psi^{-1} + \psi \circ \frac{d}{dx} \circ \psi^{-1} \circ X' \circ \psi^{-1}.$$

Thus $A \circ X \circ A^{-1} \in \mathbb{C}[[x]][d/dx]$ if and only if $\psi^{-1} \cdot X\psi$ is a constant function, i.e. if ψ is an an eigenfunction of X. This proves (ii) and the " \supset "-part of (i). Since $\dim_{\mathbb{C}} R/R' \leq 1$ the lemma is proved.

The assertion in Chapter 3 follows from Lemma 1 and 2. We now want to compare $\kappa(R') = \kappa(R) \cap \mathbb{C}[[x]][d/dx]$ with the ring \tilde{R} of all operators commuting with \tilde{Z} . Obviously $\kappa(R) \cap \mathbb{C}[[x]][d/dx] \subset \tilde{R}$, and we have

LEMMA 3. If $\kappa(R) \cap \mathbb{C}[[x]][d/dx] \neq \tilde{R}$ then $\psi \in E_Q$ for some singular point $Q \in C$.

Proof. We put $Y' := \kappa(Y \circ Z)$ if $\psi \notin E$ and $Y' := \kappa(Y)$ if $\psi \in E$. In the first case $Y'^2 + Z^2 \circ F(\tilde{Z}) = 0$, in the second case $Y'^2 + F(\tilde{Z}) = 0$. In both cases $\kappa(R) \cap C[[x]][d/dx] = C[Y', \tilde{Z}]$. Since $\tilde{R} \supset C[Y', \tilde{Z}]$, $\tilde{R} = C[\tilde{Y}, \tilde{Z}]$ with some differential operator \tilde{Y} fulfilling an equation of the form $\tilde{Y}^2 + \tilde{F}(\tilde{Z}) = 0$ with a polynomial \tilde{F} of odd degree. Then $Y' = g(\tilde{Z}) \circ \tilde{Y}$ with $g(z) \in C[z]$.

First we show that $g(z) = az^m$ for some $a \in \mathbb{C}$, $n \in \mathbb{N}$. Otherwise Y' splits $Y' = (\tilde{Z} - b) \circ Y''$ with $b \neq 0$, $Y'' \in \tilde{R} - R$. But this implies that $b \cdot \kappa^{-1}(Y'') = \kappa^{-1}(\tilde{Z} \cdot Y'') - \kappa^{-1}(Y') = A^* \circ Y'' \circ A - \kappa^{-1}(Y')$ is a differential operator in R, thus $Y'' \in R'$.

Put $C'_0 :=$ Spec R', $\tilde{C}_0 :=$ Spec \tilde{R} . Since $\kappa : R[Z^{-1}] \xrightarrow{\cong} \kappa(R')[\tilde{Z}^{-1}] = \tilde{R}[\tilde{Z}^{-1}]$ the maps $\pi, \tilde{\pi}$ in the diagram below are isomorphisms outside of $\{(z, y) \in C'_0 | z = 0\}$



From dim $R/R' \leq 1$ and the condition $\kappa(R') \neq \tilde{R}$ it follows that there exists a regular map $\pi_1: \tilde{C}_0 \to C_0$ such that $\pi \circ \pi_1 = \tilde{\pi}$. Thus $\kappa(R)$ is contained in \tilde{R} and by Lemma 2, $\psi \in E_Q$ for some point $Q = (0, y) \in C_0$. If Q were a regular point of C_0 then $\pi_1: \tilde{C}_0 \to C_0$ were an isomorphism and $\kappa(R) = \tilde{R}$.

Remark. An application of Lemma 2 (i) shows that in this case Z is obtained from \tilde{Z} by a Bäcklund-transformation with parameters $0, \psi^{-1}$.

4.2. Let us suppose that $\kappa(R) \cap \mathbb{C}[[x]][d/dx] = \tilde{R}$, i.e. $\tilde{R} = A \circ R' \circ A^{-1}$. If $\psi \notin E$, then \tilde{C} is given by $y^2 + z^2 F(z) = 0$, if $\psi \in E$ then $\tilde{C} = C$. Let $\phi : \tilde{R} \to R$ be the map $D \mapsto A^{-1} \circ D \circ A$ and $\pi : C \to \tilde{C}$ the dual map. If $\psi \notin E$ then π is the map induced by $(z, y) \mapsto (z, zy)$; if $\psi \in E$ then it is the identity map of C.

In order to determine the line bundle on \tilde{C} the space $M = \mathbb{C}[d/dx]$ will be considered as a module over various rings (cf. Chapter 2). The ring in question will be indicated by an index: M_R , $M_{\tilde{R}}$ etc. As in Chapter 2 we have the following natural identifications

$$\Gamma(C-P,L) \cong M_R, \qquad \Gamma(\tilde{C}-\tilde{P},\tilde{L}) \cong M_{\tilde{R}}, \qquad \Gamma(\tilde{C}-\tilde{P},\pi_*\mathcal{L}) = M_{\phi(\tilde{R})}.$$

(By $M_{\phi(\tilde{R})}$ we denote the \tilde{R} -module M where the scalar-multiplication is given by $X \cdot D := (X \circ \phi(D))_0, X \in M, D \in \tilde{R}$.)

LEMMA 4. The map $\phi: M_{\tilde{R}} \to M_{\phi(\tilde{R})}, X \mapsto (X \circ A)_0$ is \tilde{R} -linear.

Proof. Let $D \in \overline{R}$, $X \in M_{\overline{R}}$. Then

$$\Phi(X \cdot D) = ((X \circ D)_0 \circ A)_0 = (X \circ D \circ A)_0 =$$

= $(X \circ A \circ A^{-1} \circ D \circ A)_0 = ((X \circ A)_0 \circ (A^{-1} \circ D \circ A))_0 = \Phi(X) \cdot \phi(D).$

(The above identities hold because $(X \circ X')_0 = (X_0 \circ X')_0$ for any two differential operators $X, X' \in \mathbb{C}[[x]][d/dx]$.)

LEMMA 5.
$$\Phi(M_{\tilde{R}}) = \{X \in M_{\phi(\tilde{R})} / (X\psi)(0) = 0\}.$$

Proof. Since A has order 1 the image $\Phi(M_{\tilde{R}})$ has codimension 1 in $M_{\phi(\tilde{R})}$. On the other hand the non-trivial map $X \mapsto (X\psi)(0)$ vanishes on $\Phi(M_{\tilde{R}})$ since $A\psi = 0$.

Second proof: By an argument as in the proof of Lemma 2 one sees that $(X\psi)(0) = 0$ if and only if $(X \circ A^{-1})_0$ is a differential operator.

Let \mathcal{F} be the sheaf of sections s of $\pi_* \mathcal{L}$ with $\langle s_0, \psi \rangle = 0$. Obviously

$$\Phi(M_{\tilde{R}}) \subset \Gamma(\tilde{C} - \tilde{P}, \mathcal{F}) \subset M_{\phi(\tilde{R})} = \Gamma(\tilde{C} - \tilde{P}, \pi_* \mathcal{L}).$$

Since $\Phi(M_{\tilde{R}})$ has codimension 1 in $M_{\phi(\tilde{R})}$ and $\Gamma(\tilde{C}-\tilde{P}, \mathcal{F}) \neq \Gamma(\tilde{C}-\tilde{P}, \pi_*\mathfrak{L})$ it follows that $\Phi(M_{\tilde{R}}) = \Gamma(\tilde{C}-\tilde{P}, \mathcal{F})$, hence

$$\Gamma(\tilde{C}-\tilde{P},\tilde{\mathscr{L}})\cong\Gamma(\tilde{C}-\tilde{P},\mathfrak{F}).$$

Because A has order 1 Φ induces a sheaf map $\tilde{\mathscr{L}} \to \pi_*\mathscr{L}$ with a pole of order 1 at \tilde{P} . Hence $\tilde{\mathscr{L}} \cong \mathfrak{F} \otimes \mathfrak{O}(\tilde{P})$, whenever $\kappa(R) \cap \mathbb{C}[[x]][d/dx] = \tilde{R}$. This, together with Lemma 3, proves part (i) and (ii) of the theorem.

4.3. Now suppose that $\psi \in E_Q$ for a singular point $Q \in C$ (which then has the coordinates (0, 0)). If $\kappa(R) \cap \mathbb{C}[[x]][d/dx]$ were equal to \tilde{R} then by the results of 4.2 $\tilde{\mathscr{L}}$ were isomorphic to $\mathcal{F} \otimes \mathbb{O}(\tilde{P})$. But this sheaf is not invertible at Q, whereas $\tilde{\mathfrak{L}}$ is invertible.

So $\kappa(R) \cap \mathbb{C}[[x]][d/dx] \subset \tilde{R}$. In the remark after Lemma 3 we noticed that \tilde{C} has the normal form $y^2 + 1/z^2 F(z) = 0$ and that u is obtained by a Bäcklund transformation applied to \tilde{u} . By part (i) of the theorem this implies that $\tilde{L} = \tau^*(L) \otimes \mathbb{O}(-P)$, where $\tau: \tilde{C} \to C$ is the map $(z, y) \mapsto (z, yz)$. Thus also part (ii) of the theorem is proved.

REFERENCES

- [1] ADLER, M. and MOSER, J., On a class of polynomials connected with the Korteweg-De Vries equation. Comm. Math. Physics 61 (1978) 1-30.
- [2] AIRAULT, H., MCKEAN, H. and MOSER, J., Rational and elliptic solutions of the Korteweg-De Vries equation and a related many body problem. Comm. Pure Appl. Math. 30 (1977) 95-148.
- [3] BURCHNALL, J. and CHAUNDY, T., Commutative ordinary differential operators. Proc. Roy. Soc. London, Ser. A, 118 (1928) 557-583.
- [4] DARBOUX, G., Leçons sur la théorie générale des surfaces II. Gauthier-Villars, Paris 1889.
- [5] DEIFT, P., Applications of a Commutation formula. Duke Math. Journ. 45, (1978) 267-310.
- [6] DRACH, J., Sur l'intégration par quadrature de l'équation $d^2y/dx^2 = [\phi(x)+h] \cdot y$. C. R. Acad. Sci. Paris 168, (1919) 337-340.
- [7] FLASCHKA, H. and MCLAUGHLIN, D., Some comments on Bäcklund transformations, canonical transformations, and the inverse scattering method. In: Bäcklund Transformations, pp. 253–295. Lecture Notes in Mathematics 515, Springer, Berlin, Heidelberg, New York 1976.
- [8] GOURSAT, E., Le Problème de Bäcklund. Gauthiers-Villars, Paris 1925.
- [9] INCE, E., Ordinary Differential Equations. Dover, New York 1956.
- [10] MCKEAN, H., Integrable systems and algebraic curves. In: Global Analysis, pp. 83-200. Lecture Notes in Mathematics 755, Springer, Berlin, Heidelberg, New York 1979.
- [11] MCKEAN, H., Theta functions, solitons, and singular curves. In: Partial Differential Equations and Geometry, pp. 237–254. Lecture Notes in Pure and Applied Mathematics 48, Marcel Dekker, New York, Basel 1979.
- [12] KRICHEVER, I. M., Methods of algebraic geometry in the theory of non-linear equations. Russ. Math. Surveys 32 (1977) 185-213.
- [13] MIURA, R., Korteweg-De Vries equation and generalizations I. A remarkable explicit nonlinear transformation. Journ. Math. Phys. 9 (1969) 1202-1204.
- [14] MUMFORD, D., An algebro-geometric construction of commuting operators and of solutions to the Toda lattice equation, Korteweg-De Vries equation and related non-linear equations. Proc. Intern. Symp. Algebraic Geometry Kyoto 1977, 115–153.
- [15] SERRE, J. P., Groupes algébriques et corps de classes. Herrmann, Paris 1959.

Mathematisches Institut der Universität Wegelerstr. 10 D 5300 Bonn 1 W.-Germany

Mathematisch Instituut Rijksuniversiteit Leiden Wassenaarseweg 80 2300 RA Leiden The Netherlands

Received May 11, 1981/November 4, 1981