

A sharp four dimensional isoperimetric inequality.

Autor(en): **Croke, Christopher B.**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **59 (1984)**

PDF erstellt am: **11.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-45390>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

A sharp four dimensional isoperimetric inequality

CHRISTOPHER B. CROKE*

Introduction

Let $(M, \partial M)$ be a compact n -dimensional Riemannian manifold (with boundary) of non positive sectional curvature. Assume further that every geodesic ray in M minimizes length up to the point it hits the boundary. In this paper we show:

THEOREM. *If $(M, \partial M)$ is as above ($n \geq 3$) then $\text{Vol}(\partial M)^n \geq C(n) \text{Vol}(M)^{n-1}$ where*

$$C(n) = \frac{\alpha(n-1)^{n-1}}{\alpha(n-2)^{n-2} \left\{ \int_0^{\pi/2} \cos(t)^{n/n-2} \sin(t)^{n-2} dt \right\}^{n-2}}$$

and $\alpha(n)$ represents the volume of the unit n sphere. If $n \neq 4$ equality never holds. If $n = 4$ equality holds if and only if M is isometric to a flat ball.

This answers a long standing conjecture in dimension 4. The conjecture states that for $(M, \partial M)$ a compact domain in a complete simply connected manifold of non-positive curvature (which implies the condition of the theorem) we have $\text{Vol}(\partial M)^n \geq \bar{C}(n) \text{Vol}(M)^{n-1}$ for $\bar{C}(n) = n^{n-1} \alpha(n-1)$, with equality holding if and only if M is isometric to a flat ball. It is an easy computation to see that $C(4) = \bar{C}(4)$. The conjecture was proved in dimension 2 by Beckenbach and Radó (see [B–R]) in 1933, and is open in all dimensions except 2, and now 4.

This conjecture is a special case of a more general conjecture (see [A]) where an upper bound K (not necessarily 0) on the curvature is assumed. The more general conjecture was proved in dimension 2 by Aubin (see [A]), and for constant curvature in all dimensions by Schmidt (see [Sc]).

The isoperimetric constants are related to Sobolev constants (see [A], [Bo], and [FF] among other). In particular we see that for a domain D in a simply

* Supported by NSF Grant #MCS-01780

connected n -dimensional Riemannian manifold of non-positive curvature and for any $g \in H_0^1(M)$ we have

$$\int_D \|dg\| \geq C(n) \left\{ \int_D |g|^{n/n-1} \right\}^{(n-1)/n}$$

where $C(4)$ is the flat constant.

Such isoperimetric inequalities are interesting even with non-sharp constants. Previous non-sharp versions of the theorem are consequences of results in [H-S] and [C]. The constants $C(n)$ given here are the best known to the author in all dimensions (greater than 2). In particular $C(3) = 32\pi$ while $\bar{C}(3) = 36\pi$.

Notation and definitions

We will use the notation of [C]. Let $UM \xrightarrow{\pi} M$ represent the unit sphere bundle with the canonical (local product) measure. For $v \in UM$, let γ_v be the geodesic with $\gamma'_v(0) = v$ and let $\xi^t(v)$ represent the geodesic flow (i.e. $\xi^t(v) = \gamma'_v(t)$). For $v \in UM$ we let $l(v) = \max\{t \mid \gamma_v(t) \in M\}$. Note $\xi^t(v)$ is defined for $t \leq l(v)$ and $\gamma_v(l(v)) \in \partial M$.

For $p \in \partial M$ let N_p be the inwardly pointing unit normal vector to ∂M at p . Let $U^+\partial M \xrightarrow{\pi} \partial M$ be the bundle of inwardly pointing unit vectors (i.e. $U^+\partial M = \{u \in UM \mid \pi(u) \in \partial M \text{ and } \langle u, N_{\pi(u)} \rangle > 0\}$). We let $U_p^+\partial M$ represent $\pi^{-1}(p)$. For $u \in U^+\partial M$ we will use $\cos(u)$ to represent $\langle u, N_{\pi(u)} \rangle$. The measure on $U^+\partial M$ is the local product measure du where the measure of the fibre is that of the unit upper hemisphere.

The proof

The main tool in the proof is a formula due to Santalo:

$$(i) \int_{UM} f(v) dv = \int_{U^+\partial M} \int_0^{l(u)} f(\xi^t(u)) \cos(u) dt du$$

for all integrable functions f . The formula takes this form in our case since all geodesics in M hit ∂M . For a proof see [Sa] pp. 336–338 or [B] p. 286.

From this we derive:

LEMMA 1. a) $\text{Vol}(M) = 1/\alpha(n-1) \int_{U^+\partial M} l(u) \cos(u) du$
 b) For all integrable functions g

$$\int_{U^+\partial M} g(u) \cos(u) du = \int_{U^+\partial M} g(\text{ant}(u)) \cos(u) du$$

where $\text{ant}(u) = -\gamma'_u(l(u))$.

Proof. Part a) follows directly from (i) by letting $f(v) \equiv 1$ and integrating the t . That is

$$\alpha(n-1) \text{Vol}(M) = \int_{UM} dv = \int_{U^+\partial M} \int_0^{l(u)} \cos(u) dt du = \int_{U^+\partial M} l(u) \cos(u) du.$$

To prove part b) we first note that (i) says that the geodesic flow ξ is a measure preserving map from Q to UM where $Q = \{(u, t) \mid u \in U^+\partial M \text{ and } 0 \leq t \leq l(u)\}$ is given the measure $\cos(u) dt du$. ξ has an inverse (smooth almost everywhere) ξ^{-1} which is also measure preserving, for $v \in UM$ $\xi^{-1}(v) = (-\gamma'_{-v}(l(-v)), l(-v))$. Since the antipodal map $-1 : UM \rightarrow UM$ is also measure preserving we have $\xi^{-1} \circ (-1) \circ \xi : Q \rightarrow Q$ is measure preserving. Since $\xi^{-1} \circ (-1) \circ \xi(u, t) = (\text{ant}(u), l(u) - t)$ we see that for every integrable $G : Q \rightarrow \mathbb{R}$ we have:

$$\int_{U^+\partial M} \int_0^{l(u)} G(u, t) \cos(u) dt du = \int_{U^+\partial M} \int_0^{l(u)} G(\text{ant}(u), l(u) - t) \cos(u) dt du$$

To complete the proof of part b) simply take $G(u, t) = g(u)/l(u)$ and integrate the t (note: $l(\text{ant } u) = l(u)$).

LEMMA 2. a) $\int_{U^+\partial M} l(u)^{n-1} / \cos(\text{ant } u) du \leq \text{Vol}(\partial M)^2$ with equality holding if and only if M is flat and convex.

b) $\int_{U^+\partial M} (\cos(\text{ant } u))^{1/n-2} (\cos(u))^{n-1/n-2} du \leq \text{Vol}(\partial M) \cdot C_2(n)$ where $C_2(n) = \alpha(n-2) \int_0^{\pi/2} \cos^{n/n-2}(t) \sin^{n-2}(t) dt$. Equality holds if and only if $\cos(u) = \cos(\text{ant } u)$ almost everywhere.

Remark. “almost everywhere” above can be replaced with “everywhere” but it is not worth going into.

Proof. Let dx be the volume form on M and dp the volume form on ∂M . Let $q \in \partial M$. In normal polar coordinates (u, r) about q in the region $\text{Exp}\{tu \mid u \in U_q^+ \partial M \text{ and } 0 \leq t \leq l(u)\}$ we have $dx = F(u, r) du dr$ for some function $F(u, r)$. Let $A = \text{Exp}\{tu \mid t = l(u)\}$. Then $A \subset \partial M$ and dp on A is precisely $(F(u, l(u))/\cos(\text{ant } u)) du$. Thus we see

$$\int_{U_q^+ \partial M} \frac{F(u, l(u))}{\cos(\text{ant } u)} du = \text{Vol}(A) \leq \text{Vol}(\partial M).$$

Equality holds if and only if $A = \partial M$. That is, M is (geodesically) star shaped from q .

Integrating over q we get

$$\int_{U^+ \partial M} \frac{F(u, l(u))}{\cos(\text{ant } u)} du \leq \text{Vol}(\partial M)^2$$

with equality holding if and only if M is convex. Part a) now follows since M having non-positive curvature implies $F(u, l(u)) \geq l(u)^{n-1}$ with equality if and only if the sectional curvatures of all sections containing $\gamma'_u(t)$ for some t , are 0 (see [B-C] Section 11.10).

To prove part b) we apply a Schwarz inequality and Lemma 1b.

$$\begin{aligned} & \int_{U^+ \partial M} (\cos(\text{ant } u))^{1/n-2} (\cos(u))^{n-1/n-2} du \\ &= \int_{U^+ \partial M} (\cos(\text{ant } u))^{1/n-2} (\cos(u))^{1/n-2} \cos(u) du \\ &\leq \left\{ \int_{U^+ \partial M} (\cos(\text{ant } u))^{2/n-2} \cos(u) du \right\}^{1/2} \cdot \left\{ \int_{U^+ \partial M} (\cos(u))^{2/n-2} \cos(u) du \right\}^{1/2} \\ &= \int_{U^+ \partial M} (\cos(u))^{n/n-2} du = \int_{\partial M} \left(\int_{U^+ \partial M} (\cos(u))^{n/n-2} du \right) dq \\ &= \text{Vol}(\partial M) \cdot C_2(n) \end{aligned}$$

In order for equality to hold we need to have equality in the inequality, i.e. $\cos(\text{ant } u) = K \cos(u)$ almost everywhere for some constant K . Since the maximum values of both $\cos(\text{ant } u)$ and $\cos(u)$ are 1 it is clear that K must be 1.

Proof of the theorem. By Lemma 1a and a Hölder inequality we have

$$\begin{aligned} \text{Vol}(M) &= \frac{1}{\alpha(n-1)} \int_{U^+\partial M} l(u) \cos(u) \, du \\ &= \frac{1}{\alpha(n-1)} \int_{U^+\partial M} \frac{l(u)}{(\cos(\text{ant } u))^{1/n-1}} (\cos(\text{ant } u))^{1/n-1} \cos(u) \, du \\ &\leq \frac{1}{\alpha(n-1)} \left\{ \int_{U^+\partial M} \frac{l(u)^{n-1}}{\cos(\text{ant } u)} \, du \right\}^{1/n-1} \\ &\quad \cdot \left\{ \int_{U^+\partial M} (\cos(\text{ant } u))^{1/n-2} (\cos(u))^{n-1/n-2} \, du \right\}^{n-2/n-1}. \end{aligned}$$

Applying Lemmas 2a and 2b we get

$$\text{Vol}(M) \leq \frac{1}{\alpha(n-1)} (\text{Vol}(\partial M))^{2/n-1} \cdot (\text{Vol}(\partial M))^{n-2/n-1} C_2(n)^{n-2/n-1}.$$

hence

$$C(n) \text{Vol}(M)^{n-1} = \frac{\alpha(n-1)^{n-1}}{C_2(n)^{n-2}} (\text{Vol}(M))^{n-1} \leq (\text{Vol}(\partial M))^n.$$

In order for equality to hold we must have equality in Lemmas 2a and 2b as well as the above Hölder inequality. By Lemma 2a we see that M must be flat and hence the theorem follows from the classical result in \mathbb{R}^n , since $C(4)$ is sharp and $C(n)$ for $n \neq 4$ is not. One can also see the $n = 4$ case directly. Equality in the Hölder inequality gives

$$\frac{l^3(u)}{\cos(\text{ant } u)} = K \cos(\text{ant } u)^{1/2} \cos(u)^{3/2}$$

almost everywhere for some constant K . By the equality condition in Lemma 2b we see $l(u) = 2r \cos(u)$ for some constant r . It is now easy to see (since M is flat) that M is a ball of radius r .

Remark. A similar equality analysis for $n \neq 4$ would require:

- 1) M flat and convex
- 2) $\cos(\text{ant } u) = \cos(u)$
- 3) $l(u) = K \cos^{2/n-2}(u)$.

No such M exists.

REFERENCES

- [A] T. AUBIN, *Problèmes isopérimétriques et espaces de Sobolev*, J. Differential Geometry 11 (1976), 573–598.
- [B] M. BERGER, *Lectures on geodesics in Riemannian geometry*, Tata Institute, Bombay, 1965.
- [B–R] E. F. BECKENBACH and T. RADÓ, *Subharmonic functions and surfaces of negative curvature*, Trans. Amer. Math. Soc. 35 (1933), 662–674.
- [B–C] R. BISHOP and R. CRITTENDEN, *Geometry of manifolds*, Academic Press, 1964.
- [Bo] E. BOMBIERI, *Theory of minimal surfaces, and a counterexample to the Bernstein conjecture in high dimension*, Lecture Notes, Courant Inst., 1970.
- [C] C. CROKE, *Some isoperimetric inequalities and eigenvalue estimates*, Ann. Scient. Éc. Norm. Sup., 4^e série, t. 13, 1980, 419–435.
- [F–F] H. FEDERER and W. FLEMMING, *Normal integral currents*, Ann. of Math. 72 (1960), 458–520.
- [H–S] D. HOFFMAN and J. SPRUCK, *Sobolev and isoperimetric inequalities for Riemannian submanifolds*, Communications Pure Appl. Math. 27 (1974), 715–727; correction, ibid. 28 (1975), 765–766.
- [Sa] L. A. SANTALÓ, *Integral geometry and geometric probability*, (Encyclopedia of Mathematics and Its Applications), Addison-Wesley, London–Amsterdam–Don Mills, Ontario–Sydney–Tokyo, 1976.
- [Sc] E. SCHMIDT, *Beweis der isoperimetrischen Eigenschaft der Kugel in hyperbolischen und sphärischen Raum jeder Dimensionenzahl*, Math. Z. 49 (1943/44), 1–109.

University of Pennsylvania
Philadelphia, PA 19104 USA

Received March 28, 1983