

# On the embedding of 1-convex manifolds with 1-dimensional exceptional sets.

Autor(en): **Coltui, Mihnea**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **60 (1985)**

PDF erstellt am: **16.08.2024**

Persistenter Link: <https://doi.org/10.5169/seals-46326>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## On the embedding of 1-convex manifolds with 1-dimensional exceptional set

MIHNEA COLTOIU

### Introduction

Let  $X$  be a 1-convex manifold and  $S \subset X$  its exceptional set.  $X$  is called embeddable if there exists a holomorphic embedding of  $X$  into  $\mathbb{C}^k \times \mathbb{P}^l$  for suitable  $k, l \in \mathbb{N}$ . When  $X$  has dimension 2 a result of C. Bănică [1], proved also by Vo Van Tan [13c], asserts that  $X$  is embeddable (in fact in this case we may allow  $X$  to have singularities).

The purpose of the present paper is to generalize this result to higher dimensions. We consider a 1-convex manifold  $X$  such that its exceptional set  $S$  is an irreducible curve. Under the assumption that  $S$  is not rational (i.e. its normalization is not  $\mathbb{P}^1$ ) we prove that  $X$  is embeddable. A similar result holds if we assume that  $S \cong \mathbb{P}^1$  and  $\dim X \neq 3$  (see Theorem 5).

The technique of proof enables us to obtain also the following result:

If  $X$  is a complex manifold (not necessarily 1-convex) and  $S \subset X$  is an irreducible exceptional curve with the above properties then the fundamental class of  $S$  in  $X$  does not vanish (see Theorem 6).

### 1. Preliminaries

Throughout this paper we shall not distinguish between holomorphic line bundles and invertible sheaves.

If  $X$  is a complex manifold and  $L$  is a holomorphic line bundle on  $X$  given by transition functions  $\{g_{kl}\}$  corresponding to an open covering  $\{U_k\}$  of  $X$ , a hermitian metric on  $L$  is a system  $\{h_k\}$  of  $C^\infty$  functions  $h_k: U_k \rightarrow (0, \infty)$  such that  $h_k/h_l = |g_{kl}|^2$  on  $U_k \cap U_l$ .

$L$  is said to be Nakano semipositive if there exists a hermitian metric  $h = (h_k)$  on  $L$  such that  $-\log h_k$  is plurisubharmonic on  $U_k$  for any  $k$ .

Let now  $X$  be a 1-convex manifold and  $S \subset X$  its exceptional set.  $X$  is said to be embeddable if it can be realized as a closed analytic submanifold of some  $\mathbb{C}^k \times \mathbb{P}^l$ .

The following theorem of M. Schneider [12], proved also by Vo Van Tan [13a], gives sufficient and necessary conditions for a 1-convex manifold to be embeddable.

**THEOREM 1.** *Let  $X$  be a 1-convex manifold and  $S \subset X$  its exceptional set. Then  $X$  is embeddable iff there exists a holomorphic line bundle  $L$  on  $X$  such that  $L|_S$  is ample.*

If  $X$  is a complex manifold we denote by  $K = K_X$  the canonical line bundle on  $X$ . In order to prove our results we shall need also the following “precise vanishing theorems”:

**THEOREM 2** [10] [13b]. *Let  $X$  be a 1-convex manifold with exceptional set  $S$  and let  $L$  be a holomorphic line bundle on  $X$  such that  $L|_S$  is ample. Then  $H^q(X, K \otimes L) = 0$  for  $q \geq 1$ .*

**THEOREM 3** [5]. *Let  $X$  be a Kählerian manifold and  $L$  a Nakano semipositive line bundle on  $X$ . If  $D \subset X$  is a relatively compact strongly pseudoconvex domain with smooth boundary then  $H^q(D, K \otimes L) = 0$  for  $q \geq 1$ .*

## 2. Main results

**DEFINITION.** Let  $S$  be an irreducible curve and  $\pi: \tilde{S} \rightarrow S$  its normalization.  $S$  is called a rational curve iff  $\tilde{S} = \mathbb{P}^1$ .

The following theorem explains us the behaviour of the canonical bundle in the neighbourhood of an exceptional irreducible curve.

**THEOREM 4.** *Let  $X$  be a 1-convex manifold and assume that its exceptional set  $S$  is an irreducible curve. Suppose that:*

- a)  $S$  is not a rational curve or
- b)  $S \cong \mathbb{P}^1$  and  $\dim X \geq 4$

*Then  $K|_S$  is ample.*

The proof of Theorem 4 is based on several lemmas.

**LEMMA 1.** *Let  $X$  be a 1-convex manifold,  $S \subset X$  its exceptional set and  $k = \dim S$ . Then for every  $\mathcal{F} \in \text{Coh}(X)$  it follows that  $H^q(X, \mathcal{F}) = 0$  for  $q > k$ .*

*Proof.* By a theorem of Narasimhan [9]  $H^q(X, \mathcal{F}) \cong H^q(S, \mathcal{F}|_S)$  for any  $q > 0$ .

Here  $\mathcal{F}|_S$  denotes the topological restriction of  $\mathcal{F}$  to  $S$ , hence  $\mathcal{F}|_S$  is not a coherent sheaf on  $S$ . However, by a result of Reiffen [11 Satz 2] the cohomology groups  $H^q(S, \mathcal{F}|_S)$  vanish for  $q > k$  and the lemma is proved.

**LEMMA 2.** *Let  $X$  be a 1-convex manifold such that its exceptional set  $S$  is 1-dimensional. Then  $S$  has a Kählerian neighbourhood.*

A proof of this lemma can be found in [10 p. 165]. In fact it is shown that  $S$  has an embeddable neighbourhood.

If  $S$  is an irreducible curve we denote by  $\pi: \tilde{S} \rightarrow S$  its normalization. There is an injective morphism of sheaves  $\mathcal{O}_S \xrightarrow{i} \pi_*\mathcal{O}_{\tilde{S}}$  where  $\pi_*\mathcal{O}_{\tilde{S}}$  is the 0-direct image of  $\mathcal{O}_{\tilde{S}}$  (i.e. the sheaf of weakly holomorphic functions on  $S$ ). Let  $\mathbb{R}_S$  be the sheaf on  $S$  of locally constant real valued functions and similarly define  $\mathbb{R}_{\tilde{S}}$  on  $\tilde{S}$ . If  $\mathbb{R}_S \xrightarrow{j} \mathcal{O}_S$  is the natural inclusion map then  $k = i \circ j$  is an injective morphism of sheaves. Let  $k^*: H^1(S, \mathbb{R}_S) \rightarrow H^1(S, \pi_*\mathcal{O}_{\tilde{S}})$  denote the induced map on cohomology.

**LEMMA 3.** *The map  $k^*$  is surjective.*

*Proof.* Consider first the commutative diagram

$$\begin{array}{ccc} H^1(\tilde{S}, \mathbb{R}_{\tilde{S}}) & \xrightarrow{\alpha} & H^1(\tilde{S}, \mathcal{O}_{\tilde{S}}) \\ \uparrow \gamma & & \uparrow \delta \\ H^1(S, \pi_*\mathbb{R}_{\tilde{S}}) & \xrightarrow{\beta} & H^1(S, \pi_*\mathcal{O}_{\tilde{S}}) \end{array}$$

Remark that:

the map  $\delta$  is bijective since  $R^q\pi_*(\mathcal{O}_{\tilde{S}}) = 0$  for  $q > 0$  ( $\pi$  is a finite morphism).

the map  $\gamma$  is bijective since  $R^q\pi_*(\mathbb{R}_{\tilde{S}}) = 0$  for  $q > 0$

(if  $U \subset S$  is contractible it follows easily that  $H^q(\pi^{-1}(U), \mathbb{R}_{\tilde{S}}) = 0$  for  $q > 0$ ; since any point in  $S$  has a fundamental system of contractible open neighbourhoods we deduce that  $R^q\pi_*(\mathbb{R}_{\tilde{S}}) = 0$  for  $q > 0$ ).

the map  $\alpha$  is bijective since  $\tilde{S}$  is Kählerian.

It follows from the commutativity of this diagram that  $\beta$  is bijective.

Consider now the commutative diagram:

$$\begin{array}{ccc} H^1(S, \pi_*\mathbb{R}_{\tilde{S}}) & \xrightarrow{\beta} & H^1(S, \pi_*\mathcal{O}_{\tilde{S}}) \\ \uparrow v & \nearrow k^* & \uparrow i^* \\ H^1(S, \mathbb{R}_S) & \xrightarrow{j^*} & H^1(S, \mathcal{O}_S) \end{array}$$



The map  $v$  is surjective because  $\text{supp}(\pi_*\mathbb{R}_{\tilde{S}}/\mathbb{R}_S)$  is a finite set. Hence  $k^*$  is surjective and Lemma 3 is proved.

**LEMMA 4.** *Let  $S$  be an irreducible curve and  $\pi : \tilde{S} \rightarrow S$  its normalization. Let  $L$  be a holomorphic line bundle on  $S$  which is topologically trivial. Then there exists a holomorphic line bundle  $L'$  on  $S$  which can be given by constant transition functions  $\{g_{kl}\}$  with  $|g_{kl}| = 1$  and such that  $\pi^*(L \otimes L')$  is the trivial line bundle on  $\tilde{S}$ .*

*Proof.* Let  $\mathcal{U} = \{U_i\}$  be a finite open covering of  $S$  such that  $L|_{U_i}$  is trivial and all intersections  $U_{i_0} \cap \dots \cap U_{i_r}$  are connected and contractible. Let  $h_{kl} \in \mathcal{O}^*(U_k \cap U_l)$  denote the transition functions for  $L$ . Since  $L$  is topologically trivial and the covering  $\mathcal{U}$  is topologically acyclic we can find holomorphic functions  $\lambda_{kl} \in \mathcal{O}(U_k \cap U_l)$  such that  $\exp(2\pi i \lambda_{kl}) = h_{kl}$  and  $\lambda_{kl} + \lambda_{ls} + \lambda_{sk} = 0$  on  $U_k \cap U_l \cap U_s$  for any  $k, l, s$ . Hence  $\{\lambda_{kl}\}$  defines a cocycle in  $Z^1(\mathcal{U}, \mathcal{O}_S)$ . Set:  $\hat{U}_i = \pi^{-1}(U_i)$ ,  $\hat{\mathcal{U}} = \{\hat{U}_i\}$  and  $\hat{\lambda}_{kl} = \lambda_{kl} \circ \pi \cdot \{\hat{\lambda}_{kl}\}$  is a cocycle in  $Z^1(\hat{\mathcal{U}}, \pi_*\mathcal{O}_{\tilde{S}})$ . Consider now the commutative diagram:

$$\begin{CD} H^1(\mathcal{U}, \mathbb{R}_S) @>p>> H^1(\mathcal{U}, \pi_*\mathcal{O}_{\tilde{S}}) \\ @VVV @VVV \\ H^1(S, \mathbb{R}_S) @>k^*>> H^1(S, \pi_*\mathcal{O}_{\tilde{S}}) \end{CD}$$

Note that:

- the map  $k^*$  is surjective by Lemma 3
- the map  $m$  is bijective because  $\mathcal{U}$  is topologically acyclic
- the map  $n$  is injective

It follows that  $p$  is surjective. This implies that one can find a cocycle  $\{c_{kl}\} \in Z^1(\mathcal{U}, \mathbb{R}_S)$  and holomorphic functions  $f_k \in \mathcal{O}(\hat{U}_k)$  such that  $\hat{\lambda}_{kl} - f_k + f_l = c_{kl}$  on  $\hat{U}_k \cap \hat{U}_l$  for any  $k, l$ .

If  $L'$  is the holomorphic line bundle on  $S$  with transition functions  $g_{kl} = \exp(-2\pi i c_{kl})$  it follows from our construction that  $\{\exp(2\pi i f_k)\}$  defines a nonvanishing section in  $\pi^*(L \otimes L')$ , hence  $\pi^*(L \otimes L')$  is the trivial line bundle and Lemma 4 is completely proved.

**LEMMA 5.** *Let  $S$  be an irreducible curve and  $\pi : \tilde{S} \rightarrow S$  its normalization. Suppose that there exists a holomorphic line bundle  $L$  on  $S$  such that  $H^1(S, L) = 0$  and  $\pi^*L$  is the trivial line bundle on  $\tilde{S}$ . Then  $S$  is a rational curve.*

*Proof.* There is a canonical morphism of sheaves  $L \xrightarrow{\phi} \pi_*\pi^*L$ . If we set  $\mathcal{F}_1 = \ker \phi$  and  $\mathcal{F}_2 = \text{Im } \phi$  we get an exact sequence

$$0 \rightarrow \mathcal{F}_1 \rightarrow L \rightarrow \mathcal{F}_2 \rightarrow 0$$

Since  $H^1(S, L) = 0$  by hypothesis and  $H^2(S, \mathcal{F}_1) = 0$  because  $\dim S = 1$  it follows from the long exact sequence of cohomology that  $H^1(S, \mathcal{F}_2) = 0$ .

Consider now the exact sequence

$$0 \rightarrow \mathcal{F}_2 \rightarrow \pi_*\pi^*L \rightarrow \frac{\pi_*\pi^*L}{\mathcal{F}_2} \rightarrow 0$$

Since  $\text{supp}(\pi_*\pi^*L/\mathcal{F}_2)$  is a finite set it follows that  $H^1(S, \pi_*\pi^*L/\mathcal{F}_2) = 0$ , hence  $H^1(S, \pi_*\pi^*L) = 0$ . But  $H^1(S, \pi_*\pi^*L) \cong H^1(\tilde{S}, \pi^*L)$  because  $\pi$  is a finite morphism. We deduce that  $H^1(\tilde{S}, \mathcal{O}_{\tilde{S}}) = 0$  and consequently  $\tilde{S} \cong \mathbb{P}^1$ , i.e.  $S$  is a rational curve. Lemma 5 is completely proved.

We are now in a position to prove Theorem 4.

a) Suppose first that  $S$  is an irreducible curve which is not rational. We prove that  $K|_S$  is ample.

It is easy to verify that  $H^2(S, \mathbb{Z}) \cong H^2(\tilde{S}, \mathbb{Z}) \cong \mathbb{Z}$  for any irreducible curve and if  $F$  is a holomorphic line bundle on  $S$  then  $F$  is ample iff  $c(F)$  (the Chern class of  $F$ ) corresponds under the above isomorphisms to a strictly positive integer. Consequently we have to prove that  $c(K|_S) > 0$ .

We remark first that  $c(K|_S) \geq 0$ . Indeed, if  $c(K|_S) < 0$  then  $K^{-1}$  (the dual of  $K$ ) is ample when restricted to  $S$ . By Theorem 2 we obtain  $H^1(X, K \otimes K^{-1}) = 0$ , hence  $H^1(X, \mathcal{O}_X) = 0$ . If  $\mathcal{I}$  denotes the ideal sheaf of  $S$  there is an exact sequence of sheaves on  $X$ :

$$0 \rightarrow \mathcal{I} \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_X/\mathcal{I} \rightarrow 0$$

Since  $H^1(X, \mathcal{O}_X) = 0$  and  $H^2(X, \mathcal{I}) = 0$  (by Lemma 1) we deduce from the long exact sequence of cohomology that  $H^1(S, \mathcal{O}_S) = 0$  which implies  $S \cong \mathbb{P}^1$ . This contradicts our assumption that  $S$  is not a rational curve. So we must have  $c(K|_S) \geq 0$ .

In order to prove Theorem 4 in case a) we have only to verify that  $c(K|_S) \neq 0$ .

Suppose that  $c(K|_S) = 0$ , hence  $L := K|_S$  is topologically trivial. If  $\pi: \tilde{S} \rightarrow S$  denotes the normalization of  $S$  from Lemma 4 there exists a holomorphic line bundle  $L'$  on  $S$  which can be given by constant transition functions  $\{g_{kl}\}$  with  $|g_{kl}| = 1$  and such that  $\pi^*(L \otimes L')$  is the trivial line bundle on  $\tilde{S}$ .

By Lemma 2  $S$  has an open neighbourhood  $U$  which is Kählerian and shrinking  $U$  if necessary we may assume that there exists a continuous retract  $\rho: U \rightarrow S$ . Let  $S \subset U' \Subset U$  be a strongly pseudoconvex neighbourhood of  $S$  with smooth boundary and let  $\mathcal{V} = \{V_j\}$  be an open covering of  $S$  such that  $L'$  is given on  $V_k \cap V_l$  by the constants  $g_{kl}$  with  $|g_{kl}| = 1$ . Set  $\tilde{V}_k := \rho^{-1}(V_k) \subset U$  and on  $\tilde{V}_k \cap \tilde{V}_l$  consider the transition functions  $\tilde{g}_{kl} := g_{kl}$ . Since  $g_{kl}$  are constants it follows that

the cocycle  $\{\tilde{g}_{kl}\}$  defines a holomorphic line bundle  $\tilde{L}'$  on  $U$  and  $\tilde{L}'|_S = L'$ . Moreover  $\tilde{L}'$  is Nakano semipositive because  $|\tilde{g}_{kl}| = 1$  for any  $k, l$ . From Theorem 3 of Grauert and Riemenschneider we get  $H^1(U', K \otimes \tilde{L}') = 0$ .

Now consider the exact sequence on  $U'$ :

$$(*) \quad 0 \rightarrow \mathcal{I} \rightarrow \mathcal{O}_{U'} \rightarrow \mathcal{O}_{U'}/\mathcal{I} \rightarrow 0$$

where  $\mathcal{I}$  is the ideal sheaf of  $S$ . From  $(*)$  we get the exact sequence on  $U'$ :

$$(**) \quad 0 \rightarrow K \otimes \tilde{L}' \otimes \mathcal{I} \rightarrow K \otimes \tilde{L}' \rightarrow K \otimes \tilde{L}' \otimes \mathcal{O}/\mathcal{I} \rightarrow 0.$$

By Lemma 1  $H^2(U', K \otimes \tilde{L}' \otimes \mathcal{I}) = 0$ . Since  $\tilde{L}'|_S = L'$  the long exact sequence of cohomology implies that  $H^1(S, K|_S \otimes L') = 0$ . But  $\pi^*(K|_S \otimes L')$  is the trivial line bundle on  $\tilde{S}$  and from Lemma 5 it follows that  $S$  is a rational curve which contradicts our hypothesis. Consequently a) is proved.

b) Assume that  $S \cong \mathbb{P}^1$  and  $n = \dim X \geq 4$ . We shall prove that  $K|_S$  is ample.

Let  $N_{S|X}$  denote the normal bundle of  $S$  in  $X$  and  $K_S$  the canonical line bundle of  $S$ . If we use the adjunction formula  $K|_S = K_S \otimes \det(N_{S|X}^*)$  we obtain the following formula for the Chern class of  $K|_S$ :

$$c(K|_S) = c(K_S) - c(\det(N_{S|X}))$$

Since  $S \cong \mathbb{P}^1$  we have  $c(K_S) = -2$ . On the other hand a result of Laufer [6] gives the following estimation:  $c(\det(N_{S|X})) \leq -n + 1$ . Hence we obtain  $c(K|_S) \geq n - 3 > 0$  and Theorem 4 is completely proved.

*Remark.* If  $\dim X = 3$  and  $S \cong \mathbb{P}^1$  it may happen that  $K$  is trivial in the neighbourhood of  $S$ . If  $N_{S|X} = \mathcal{O}(c_1) \otimes \mathcal{O}(c_2)$ ,  $c_1 \leq c_2$ , is the decomposition of  $N_{S|X}$  into line bundles and  $K$  is trivial in the neighbourhood of  $S$  then  $(c_1, c_2) \in \{(-1, -1), (-2, 0), (-3, 1)\}$  (see Laufer [6]). Hence Theorem 4 does not hold if  $\dim X = 3$  and  $S \cong \mathbb{P}^1$ . If  $\dim X = 2$  and  $S \cong \mathbb{P}^1$  easy examples show us that  $K|_S$  may even be negative.

**THEOREM 5.** *Let  $X$  be a 1-convex manifold such that its exceptional set  $S$  is an irreducible curve. Assume that:*

a)  *$S$  is not a rational curve*

or

b)  *$S \cong \mathbb{P}^1$  and  $\dim X \neq 3$ .*

*Then  $X$  is embeddable.*

*Proof.* In case a) it follows from Theorem 4 that  $K|_S$  is ample. By Theorem 1  $X$  is embeddable. A similar argument shows us that  $X$  is embeddable if  $S \cong \mathbb{P}^1$

and  $\dim X \geq 4$ . If  $X$  has dimension 2 then  $S$  is a divisor and if we denote by  $[S]$  the corresponding line bundle it follows that  $[S]^{-1}$  (the dual of  $[S]$ ) is ample when restricted to  $S$ . Again by Theorem 1 we deduce that  $X$  is embeddable.

*Remark.* It seems very likely that Theorem 5 should hold for any curve  $S$ .

Let now  $X$  be a complex manifold,  $S \subset X$  an irreducible, compact curve and  $\pi: \tilde{S} \rightarrow S$  its normalization. The image of the fundamental class of  $\tilde{S}$  in  $H_2(X, \mathbb{Z})$  is called the fundamental class of  $S$  in  $X$ . A straightforward consequence of Theorem 4 is the following topological result:

**THEOREM 6.** *Let  $X$  be a complex manifold and  $S \subset X$  an irreducible exceptional curve such that:*

a)  $S$  is not a rational curve

or

b)  $S \cong \mathbb{P}^1$  and  $\dim X \neq 3$

*Then the fundamental class of  $S$  in  $X$  does not vanish.*

*Remarks.* i) In [13b] Vo Van Tan has proved that any 1-convex manifold with 1-dimensional exceptional set is Kählerian. Unfortunately, as we shall see, there is a gap in a main step of his proof.

According to his notations let  $\pi: X \rightarrow Y$  be the Remmert reduction of  $X$ . We assume also that the exceptional set  $S$  is a smooth curve and let  $T$  be any point of  $S$  and set  $Z := X \setminus T$ ,  $\check{S} := S \setminus T$ . If  $\hat{E}$  is a holomorphic line bundle on  $Y$  we set  $E := \pi^*(\hat{E})$  and  $L := E|_Z$ . The author asserts that if  $\hat{E}$  is positive then there exists a metric  $\{h_i\}$  on  $L$  such that:

$$(*) \quad \begin{cases} -\partial\bar{\partial} \log h_i(x) > 0 & \text{on } T_{\check{S},x} \\ -\partial\bar{\partial} \log h_i(x) \geq 0 & \text{on } N_{\check{S},x} \\ -\partial\bar{\partial} \log h_i(z) > 0 & \text{on } T_{Z,z} \text{ if } z \in Z \setminus \check{S} = X \setminus S \end{cases}$$

where  $T_{\check{S},x}$  is the tangent space to  $\check{S}$  at  $x$  and  $N_{\check{S},x}$  is the complement space of  $T_{\check{S},x}$  in  $T_{Z,x}$ .

We shall show that  $(*)$  does not hold. We take  $\hat{E}$  to be the trivial line bundle on  $Y$  which is positive since  $Y$  is Stein. It follows that  $L$  is also the trivial line bundle on  $Z$  and  $(*)$  implies the existence of a  $C^\infty$  function  $h: Z \rightarrow (0, \infty)$  such that  $-\log h$  is strongly plurisubharmonic on  $Z \setminus \check{S}$  and  $-\log h|_{\check{S}}$  is strongly plurisubharmonic. Since  $-\log h$  is strongly plurisubharmonic on  $Z \setminus \check{S}$  it follows from the continuity of second derivatives that  $-\log h$  is plurisubharmonic on  $Z$ . By a well known result concerning the extension of plurisubharmonic functions (see Grauert-Remmert [4]) there exists a plurisubharmonic function  $p$  on  $X$  such that  $p|_Z = -\log h$ . The maximum principle for plurisubharmonic

functions implies that  $p|_S = \text{constant}$ , hence  $-\log h|_S = \text{constant}$ . This contradicts the fact that  $-\log h|_S$  is strongly plurisubharmonic.

The gap in the proof of Vo Van Tan is the following: since  $\check{S} := S \setminus T$  is Stein the metric  $\{h_i\}$  can be suitably modified such that  $L|_{\check{S}}$  is Nakano positive [8] but this can be done only on  $\check{S}$  and there is no control outside  $\check{S}$ .

ii) Under the assumptions of Lemma 5 it follows that  $S$  is a rational curve with  $\dim_{\mathbb{C}} H^1(S, \mathcal{O}_S) \leq 1$ . This can easily be deduced from Riemann–Roch theorem for singular curves. Consequently all our theorems hold if we assume that  $S$  is a rational curve with  $\dim_{\mathbb{C}} H^1(S, \mathcal{O}_S) \geq 2$ .

## REFERENCES

- [1] BANICA, C., Sur les fibres infinitésimales d'un morphisme propre d'espaces complexes. Seminaire F. Norguet. Fonctions de plusieurs variables complexes. IV Springer-Verlag Lec. Notes in Math. 807 (1980).
- [2] COLTOIU, M. and MIHALACHE, N., Strongly plurisubharmonic exhaustion functions on 1-convex spaces. To appear in Math. Ann.
- [3] GRAUERT, H., Über Modifikationen und exzeptionelle analytische Mengen. Math. Ann. 146 (1962) p. 331–368.
- [4] GRAUERT, H. and REMMERT, R., Plurisubharmonische Funktionen in komplexen Raumen. Math. Z. 65 (1956) p. 175–194.
- [5] GRAUERT, H. and RIEMENSCHNEIDER, O., Kählersche Mannigfaltigkeiten mit hyper-q-konvexen Rand. Problems in Analysis (Lectures at the Sympos. in honor of Solomon Bochner, Princeton Univ., Princeton, N.J., 1969), Princeton Univ. Press, Princeton, N.J., 1970, p. 61–79.
- [6] LAUFER, H., On  $CP_1$  as exceptional set. Recent developments in S.C.V. Ann. Math. Studies, Princeton, N.J. (1981) p. 261–275.
- [7] LOJASIEWICZ, S., Triangulation of semi-analytic sets. Ann. Scuola Norm. Sup. Pisa 18 (1964) 449–474.
- [8] NAKANO, S., Vanishing theorems for weakly 1-complete manifolds. II, Publ. Res. Inst. Math. Sci., Kyoto Univ. 10 (1974) p. 101–110.
- [9] NARASIMHAN, R., The Levi problem for complex spaces. II, Math. Ann. 146 (1962) p. 195–216.
- [10] PETERNELL, T., On strongly pseudo-convex Kähler manifolds. Invent. Math. 70 (1982) p. 157–168.
- [11] REIFFEN, H. J., Riemannsche Hebbarkeitssätze für Cohomologieklassen mit kompakten Träger. Math. Ann. 164 (1966) 272–279.
- [12] SCHNEIDER, M., Familien negativer Vektorbündel und 1-convexe Abbildungen. Abh. Math. Sem. Univ. Hamburg 47 (1978) p. 150–170.
- [13] VO VAN TAN, (a) On the embedding problem for 1-convex spaces. Trans. AMS 256 (1979) p. 185–197; (b) Vanishing theorems and Kählerity for strongly pseudoconvex manifolds. Trans. AMS 261 (1980) p. 297–302; (c) Embedding theorems and Kählerity for 1-convex spaces. Commen. Mathe. Helve. 57 (1982) p. 196–201.

National Institute for Scientific  
and Technical Creation Dept. of Mathematics  
Bd. Păcii 220  
77538 Bucharest Romania

Received January 3, 1985