

# On the periodic spectrum of the 1-dimensional Schrödinger operator.

Autor(en): **Kappeler, Th.**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **65 (1990)**

PDF erstellt am: **17.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-49708>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.


Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# On the periodic spectrum of the 1-dimensional Schrödinger operator

TH. KAPPELER



## 1. Introduction

Let us consider Hill's equation

$$-y''(x) + q(x)y(x) = \lambda y(x) \quad (x \text{ in } \mathbf{R}) \tag{1}$$

$$y(x + 2) = y(x) \quad (x \text{ in } \mathbf{R}) \tag{2}$$

where  $q$  is a potential in  $L^2[0, 1]$ , periodically extended to all of  $\mathbf{R}$ . It is well known that those  $\lambda$ 's for which (1)–(2) admits a non zero solution, form a non decreasing sequence of real numbers  $\lambda_k = \lambda_k(q)$  ( $k \geq 0$ ), written with multiplicities.  $(\lambda_k)_{k \geq 0}$  is called the periodic spectrum of  $q$ . Observe that, for convenience, the period in (2) has been chosen equal to 2 rather than 1 in order to include the so called antiperiodic eigenvalues as well. For  $q$  in  $L^2[0, 1]$ , the isospectral set  $\text{Iso}(q)$  is defined to be the set of all potentials  $p$  in  $L^2[0, 1]$  such that  $\lambda_k(p) = \lambda_k(q)$  ( $k \geq 0$ ) and  $G(q)$  denotes the set of all potentials  $p$  in  $L^2[0, 1]$  with the same gaps as  $q$ , i.e.  $\lambda_0(p) = \lambda_0(q)$  and  $\lambda_{2k}(p) - \lambda_{2k-1}(p) = \lambda_{2k}(q) - \lambda_{2k-1}(q)$  ( $k \geq 1$ ). Then  $\text{Iso}(q) \subseteq G(q)$ .

This paper presents an elementary proof of a result due to J. Garnett and E. Trubowitz [GT1] which says that the converse inclusion  $G(q) \subseteq \text{Iso}(q)$  also holds:

**THEOREM** (Garnett, Trubowitz). *For all  $q$  in  $L^2[0, 1]$ ,  $\text{Iso}(q) = G(q)$ .*

In [GT1], this theorem is proved by applying harmonic measure arguments to the identification, due to Marcenko and Ostrovskii [MO], of band configurations with certain slit quarter planes. In this paper it is shown that the theorem is a direct consequence of the spectral theory for even potentials  $q$  in  $L^2[0, 1]$  (i.e.  $q(x) = q(1 - x)$ ), as it is presented in the beautiful paper [GT2], using analysis in Hilbert space.

## 2. Proof of theorem

First observe that due to the fact that  $\lambda_k(q+c) = \lambda_k(q) + c$  ( $k \geq 0$ ;  $c$  real) it suffices to prove the theorem for potentials  $q$  in  $V$  where  $V$  is given by  $V := \{q \in L^2[0, 1] : \lambda_0(q) = 0\}$ .

Let  $q$  be a fixed element in  $V$ . Clearly, for  $p$  in  $G(q)$ ,  $\text{Iso}(p) \subset G(q)$  and thus  $G(q) = \bigcup \text{Iso}(p)$  where the union extends over all  $p$  in  $G(q)$ .

Denote by  $\mu_n = \mu_n(p)$  ( $n \geq 1$ ) and  $\nu_n = \nu_n(p)$  ( $n \geq 0$ ) the Dirichlet and Neuman spectrum of  $p$  in  $L^2[0, 1]$ , that is the spectrum of (1) for the boundary conditions  $y(0)=0, y(1)=0$  and  $y'(0)=0, y'(1)=0$  respectively. It is well known (cf. e.g. [MW]) that  $\nu_0 \leq \lambda_0$  and  $\lambda_{2n-1} \leq \mu_n, \nu_n \leq \lambda_{2n}$  ( $n \geq 1$ ). By the lemma below one can find for a given  $p$  in  $V$  an element  $p_{\max}$  in  $\text{Iso}(p) \cap E$  with the properties that  $\mu_n(p_{\max}) = \lambda_{2n}(p)$  ( $n \geq 1$ ),  $\nu_0(p_{\max}) = \lambda_0(p)$  and  $\nu_n(p_{\max}) = \lambda_{2n-1}(p)$  ( $n \geq 1$ ) where  $E$  denotes the subspace of  $L^2[0, 1]$  of all even potentials  $p$  (i.e.  $p(x) = p(1-x)$ ). Thus for all  $n \geq 1$ ,  $\mu_n(p_{\max}) - \nu_n(p_{\max}) = \lambda_{2n}(q) - \lambda_{2n-1}(q)$ . For  $p$  in  $L^2[0, 1]$ , define  $\sigma(p)$  to be the sequence  $(\mu_n(p) - \nu_n(p))_{n \geq 1}$ . From the asymptotics (cf. e.g. [PT])  $\mu_n(p), \nu_n(p) = n^2\pi^2 + \int_0^1 p(x) dx + a_n, b_n$ , where  $\sum (a_n^2 + b_n^2) < \infty$ , one concludes that  $\sigma(p)$  is an element in  $l^2$ . In [GT2] it is proved that the restriction of  $\sigma$  to  $V \cap E$  is 1-1. Therefore  $p_{\max} = q_{\max}$  for all  $p$  in  $G(q)$  and thus  $\text{Iso}(p) = \text{Iso}(q)$ . This implies that  $G(q) = \text{Iso}(q)$ .

**LEMMA.** *Let  $p$  be in  $L^2[0, 1]$ . Then there exists a potential  $p_{\max}$  in  $\text{Iso}(p)$ , such that*

- (1)  $\mu_n(p_{\max}) = \lambda_{2n}(p)$  ( $n \geq 1$ )
- (2)  $\nu_0(p_{\max}) = \lambda_0(p)$  and  $\nu_n(p_{\max}) = \lambda_{2n-1}(p)$  ( $n \geq 1$ )
- (3)  $p_{\max}$  is even, i.e. an element in  $E$ .

*Proof.* For  $p$  in  $L^2[0, 1]$  with only a finite number of simple periodic eigenvalues, the existence of  $p_{\max}$  with property (1) together with  $\|p_{\max}\|_{L^2} = \|p\|_{L^2}$  is a direct consequence of results presented in [M, M]. By standard arguments one proves (2) and (3). To be more precise, denote by  $y_1(x, \lambda)$  and  $y_2(x, \lambda)$  the fundamental solutions of (1), i.e. the solutions  $y(x, \lambda)$  of (1) with the initial conditions  $y(0, \lambda) = 1, y'(0, \lambda) = 0$  and  $y(0, \lambda) = 0, y'(0, \lambda) = 1$  respectively. For  $\lambda_{2n}(p_{\max}) = \mu_n(p_{\max})$ ,  $y_2(x, \mu_n)$  is the corresponding eigenfunction and thus  $y_2'(2, \mu_n) = 1$ . By investigating the Floquet matrix

$$F(\lambda) = \begin{pmatrix} y_1(1, \lambda) & y_2(1, \lambda) \\ y_1'(1, \lambda) & y_2'(1, \lambda) \end{pmatrix}$$

one concludes that  $|y_2'(1, \mu_n)| = 1$ . Combining Corollary 2.2 and Lemma 3.4 in [PT], it follows that  $p_{\max}$  is even. Using this fact together with reflection one now

verifies that  $\nu_n(p_{\max})$  is a periodic eigenvalue of  $p_{\max}$  ( $n \geq 0$ ). From  $\nu_0 \leq \lambda_0$  and  $\lambda_{2n-1} \leq \mu_n$ ,  $\nu_n \leq \lambda_{2n}$  ( $n \geq 1$ ) it then follows that  $\nu_0(p_{\max}) = \lambda_0(p_{\max})$  and  $\nu_n(p_{\max}) = \lambda_{2n-1}(p_{\max})$  ( $n \geq 1$ ).

Towards the general case, choose a sequence  $(p_n)_{n \geq 1}$  of potentials in  $L^2[0, 1]$  such that  $p = \lim_{n \rightarrow \infty} p_n$  in the norm topology of  $L^2[0, 1]$  and such that, for  $n \geq 1$ ,  $p_n$  has only a finite number of periodic eigenvalues. (Cf. [CK] for an elementary proof concerning the existence of such a sequence). This implies that  $\lim_{n \rightarrow \infty} \lambda_k(p_n) = \lambda_k(p)$  ( $k \geq 0$ ) as the eigenvalues depend continuously on the potential. Denote by  $q_n$  the potential  $(p_n)_{\max}$  in  $\text{Iso}(p_n)$ . Then  $\lambda_{2k}(p_n) = \mu_k(q_n)$  ( $k \geq 1$ ),  $\lambda_0(p_n) = \nu_0(q_n)$  and  $\lambda_{2k-1}(p_n) = \nu_{2k-1}(q_n)$  ( $k \geq 1$ ). Moreover  $\|p_n\|_{L^2[0,1]} = \|q_n\|_{L^2[0,1]}$ . Thus  $(q_n)_{n \geq 1}$  is a sequence, bounded in  $L^2[0, 1]$ ; without loss of any generality we may assume that  $(q_n)_{n \geq 1}$  converges weakly to a potential  $q$  in  $L^2[0, 1]$ . As  $E$  is a closed subspace of  $L^2$ ,  $q$  must be an element of  $E$ . Now use that all eigenvalues  $\lambda_k$  ( $k \geq 0$ ),  $\nu_k$  ( $k \geq 0$ ) and  $\mu_k$  ( $k \geq 1$ ) depend continuously on the potential with respect to the weak topology of  $L^2[0, 1]$  to conclude that  $\lambda_k(q) = \lim_{n \rightarrow \infty} \lambda_k(q_n) = \lim_{n \rightarrow \infty} \lambda_k(p_n) = \lambda_k(p)$  ( $k \geq 0$ ) as well as  $\mu_k(q) = \lambda_{2k}(p)$  ( $k \geq 1$ ),  $\nu_0(q) = \lambda_0(p)$  and  $\nu_k(q) = \lambda_{2k-1}(p)$  ( $k \geq 1$ ). Thus  $p_{\max} := q$  has the desired properties.

REMARK. The proof of the lemma shows that for a given  $q$  in  $L^2[0, 1]$  one has  $\|p\|_{L^2} = \|q\|_{L^2}$  for all potentials  $p$  in  $\text{Iso}(q)$ . For  $q$  in  $C^\infty(\mathbf{R}/\mathbf{Z})$  this is a consequence of results in [MT].

## BIBLIOGRAPHY

- [CK] Y. COLIN DE VERDIÈRE and TH. KAPPELER, *On double eigenvalues of Hill's operator*, to appear in J. of Funct. Anal.
- [GT1] J. GARNETT and E. TRUBOWITZ, *Gaps and bands of one dimensional periodic Schrödinger operators*, Comm. Math. Helv. 59 (1984), p. 258–312.
- [GT2] J. GARNETT and E. TRUBOWITZ, *Gaps and bands of one dimensional periodic Schrödinger operators II*, Comm. Math. Helv. 62 (1987), p. 18–37.
- [MW] W. MAGNUS and W. WINKLER, *Hill's equation*, Interscience, Wiley, New York 1966.
- [MO] V. A. MARCENKO and I. O. OSTROVSKII, *A characterization of the spectrum of Hill's operator*, Math. USSR-Sbornik, 97 (1975), p. 493–554.
- [MM] H. P. MCKEAN and P. VAN MOERBEKE, *The spectrum of Hill's equation*, Invent. Math. 30 (1975), p. 217–274.
- [MT] H. P. MCKEAN and E. TRUBOWITZ, *Hill's operator and hyperelliptic function theory in the presence of infinitely many branch points*, Comm. on Pure and Appl. Math. 29 (1976), 143–226.
- [PT] J. PÖSCHEL and E. TRUBOWITZ, *Inverse Spectral Theory*, Academic Press, 1987.

*Mathematics Department*  
*Brown University*  
*Providence, RI 02912*  
*USA*

Received August 31, 1988.