

Kleine Mitteilungen

Objektyp: **Group**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **25 (1970)**

Heft 1

PDF erstellt am: **12.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*
ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

<http://www.e-periodica.ch>

Kleine Mitteilungen

The Missing Seventh Circle

(To the memory of Dr. A. Aepli, of Zürich)

The problem of Apollonius, to construct circles tangent to three given circles C_1, C_2, C_3 , has given rise to many papers and much discussion. In 1896 MUIRHEAD [2] stated that nobody before STOLL [6], in a paper published in 1873, had ever discussed the possibility of the circles C_i being so specialized that the number of contact circles varies. Stoll remarks that the inspiration for his paper was Prof. Sturm, who pointed out that in the general Apollonius configuration the number of contact circles is either 8, 4 or 0, and that Prof. Sturm encouraged him to investigate all possible cases. Stoll does observe that if the circles C_i are allowed to touch, the number of tangent circles is never 1 or 7. He allows the possibility of a tangent circle being one of the circles C_i , assuming, correctly, that since the problem is an algebraic one, a circle may be considered to touch itself. For example, if the three circles C_i touch each other, the maximum number of contact circles distinct from the C_i is 2, and since each one of the C_i satisfies the conditions of the problem, we have a total of 5 tangent circles. For the Descartes formula connected with this case see AEPPLI [1] and PEDOE [4]. Again, Stoll recognizes that a circle of zero radius (a point-circle) lying on a circle can be considered to touch this circle, so that if the three circles C_i pass through a point O , there are 4 circles of non-zero radius which touch the C_i , and to these must be added the point-circle O , so that there are 5 tangent circles.

But neither Stoll nor Muirhead noticed that it is impossible to specialize the circles C_i so that the number of tangent circles is 7, and the proof of this curious result is the purpose of this paper. It is possible to specialize the C_i so that the number of tangent circles is 0, 1, 2, 3, 4, 5 and 6, and 8 is the maximum finite number of tangent circles, but the mystic number 7 is absent. It was when I was engaged in making up exercises for a forthcoming book (PEDOE [5]) that I noticed that I could not specialize the C_i to produce 7 contact circles, and investigation revealed why this is so. Multiplicity problems are of not too frequent occurrence in geometry, and the Apollonius problem is even more interesting than was at first suspected.

The circle C_0 which is orthogonal to each of the C_i is uniquely defined, unless the C_i belong to a coaxial system (pencil) of circles. In the latter case the only tangent circles are two point-circles, in the case when the coaxial system is of the intersecting type. The circle C_0 plays a very special rôle with regard to the C_i since inversion (transformation by reciprocal radii) in C_0 maps each C_i onto itself, and maps a tangent circle C onto a tangent circle C' . When there are 8 tangent circles (which may be called the general case) these can be split into 4 pairs. We shall call the circles in a pair conjugate circles. (For all this, proved algebraically, see PEDOE [3]). If we wish to specialize the C_i so that there are only 7 tangent circles, the specialization must aim at making a pair of conjugate circles identical, since if two tangent circles which are not conjugate become identical the conjugates also become identical, and the number of tangent circles would reduce to 6, at most.

We therefore specialize the C_i so that a conjugate pair C and C' become the same circle D , say. This means that inversion in C_0 maps the tangent circle D onto itself. If this is the case, D must be orthogonal to C_0 . We therefore find ourselves with three circles C_i , a circle C_0 orthogonal to the C_i , and a circle D which touches the C_i and is also orthogonal to C_0 . We show that this means that two of the C_i touch each other.

Invert with respect to a centre of inversion on C_0 . We obtain three circles C'_i , with diameters which lie along the line C'_0 . These three circles are touched by a circle D' whose diameter also lies along C'_0 . If two circles with diameters along the same line touch at a point not on this line, they have the same centre, and must therefore coincide. If the circles are distinct contact can only take place at an endpoint of a diameter. Since D' has only 2 points of intersection with the line C'_0 , and has to touch each of C'_1, C'_2 and C'_3 at a

point on C'_0 , the three points of contact cannot be distinct. Hence at least two of the circles C'_i intersect C'_0 at the same point. That is, at least two of the circles C'_i touch each other. But if at least two of the circles C'_i touch each other, the number of circles tangent to the three C'_i is readily seen to be 6, at most.

DANIEL PEDOE, University of Minnesota, Minneapolis, USA

REFERENCES

- [1] AEPPLI, *El. Math.* 15, 9–13 (1960).
- [2] MUIRHEAD, *Proc. Edin. Math. Soc.* 14, 135–147 (1896).
- [3] PEDOE, *Circles* (Pergamon Press, 1957).
- [4] PEDOE, *Amer. Math. Monthly* 74, 627–640 (1967).
- [5] PEDOE, *A Course of Geometry for Colleges and Universities* (Cambridge University Press).
- [6] STOLL, *Math. Ann.* 6, 613–632 (1873).

Die Jakobische Achsenkonstruktion einer Ellipse

Anregung zu dieser kurzen Mitteilung ist die Arbeit [1].

Eine Ellipse ε (Mittelpunkt M) sei gegeben durch zwei konjugierte Halbmesser $\overline{MA} = a_1$ und $\overline{MB} = b_1$. Die Richtungen der Hauptachsen lassen sich folgenderweise finden. Es sei g die Gerade durch B parallel zu MA und es seien γ_1 (Mittelpunkt M_1) und γ_2 (Mittelpunkt M_2) die beiden Kreise (Halbmesser a_1) welche g in B berühren. Dann gibt es eine Affinität α_k die γ_k in ε überführt ($k = 1, 2$). Die übliche Konstruktion für das invariante Rechtwinkelpaar von α_k führt nun zu den Richtungen der Hauptachsen von ε . Der Zusammenhang mit der Jakobischen Achsenkonstruktion tritt aus der Figur in aller Klarheit hervor. Setzt man $\sphericalangle AMB = \varphi$ und sind $2a$ und $2b$ die Längen der Hauptachsen ($a > b$), dann gilt in $\triangle MM_1B$:

$$\overline{MM_1}^2 = a_1^2 + b_1^2 - 2a_1b_1\sin\varphi = (a - b)^2;$$

daher: $\overline{MM_1} = a - b$. Analog findet man $\overline{MM_2} = a + b$.

G. R. VELDKAMP, TH. EINDHOVEN

LITERATURVERZEICHNIS

- [1] H. SIEBER, *Über das invariante Rechtwinkelpaar einer schiefen Affinität und dessen Zusammenhang mit der Jakobischen Konstruktion der Achsen einer Ellipse*, *El. Math.* 24, 31–35 (1969).

