

# The impossibility of a tessellation of the plane into equilateral triangles whose sidelengths are mutually different, one of them being minimal

Autor(en): **Scherer, Karl**

Objektyp: **Article**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **38 (1983)**

Heft 1

PDF erstellt am: **08.08.2024**

Persistenter Link: <https://doi.org/10.5169/seals-37174>

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# ELEMENTE DER MATHEMATIK

Revue de mathématiques élémentaires – Rivista di matematica elementare

*Zeitschrift zur Pflege der Mathematik  
und zur Förderung des mathematisch-physikalischen Unterrichts*

El. Math.

Band 38

Nr. 1

Seiten 1–32

Basel, den 10. Januar 1983

## The impossibility of a tessellation of the plane into equilateral triangles whose sidelengths are mutually different, one of them being minimal

**Theorem.** *There is no tessellation of the euclidean plane  $\mathbf{R}^2$  into equilateral triangles whose sidelengths are mutually different, one of them being minimal.*

Proof: Assume that there is such a tessellation of  $\mathbf{R}^2$  into equilateral triangles  $t_i, i \in I$ , where  $I$  is an arbitrary set of indices. We shall eventually see that this assumption leads to a contradiction.

For  $i \in I$ , let  $l_i$  denote the sidelength of  $t_i$ . Let  $l$  be the minimum of the sidelengths. By scaling we can attain  $l = 1$ . Then the area of each triangle is at least

$$\frac{1}{4} \cdot \sqrt{3}.$$

Therefore the triangle can be enumerated and we can assume  $I = \mathbf{N}$  and  $l_1 = 1$ .

For each  $i \in \mathbf{N}$  let  $d_i$  denote the boundary of  $t_i$ . Further we define  $D$  to be the ‘grid’ of the tessellation:  $D = \cup \{d_i, i \in \mathbf{N}\}$ .

If  $n$  and  $\varepsilon$  are positive real numbers, the points

$$P_1 = \left(-\frac{n}{2} - \frac{\varepsilon}{2}, \frac{\varepsilon}{2} \sqrt{3}\right), \quad P_2 = \left(-\frac{n}{2}, 0\right), \quad P_3 = \left(\frac{n}{2} - \frac{\varepsilon}{2}, \frac{\varepsilon}{2} \sqrt{3}\right)$$

$$\text{and } P_4 = \left(\frac{n}{2}, 0\right)$$

define a parallelogram in  $\mathbf{R}^2$  with the sidelengths  $\varepsilon$  and  $n$  and with angles of 60 and 120 degrees. Let  $S(n, \varepsilon)$  denote the union of the three sides  $\overline{P_1 P_2}$ ,  $\overline{P_2 P_3}$  and  $\overline{P_3 P_4}$  (fig. 1).

More generally we denote by  $S(n, \varepsilon)$  any subset of the plane that can be obtained from this special  $S(n, \varepsilon)$  by translations, rotations and reflections. If any such  $S(n, \varepsilon)$  is given, let  $R_n$  denote the interior of the associated parallelogram.

We show that there are a strongly decreasing sequence of positive numbers  $n_j, j \in \mathbf{N}$ , with  $n_j \leq n_{j-1} - 1$  ( $j \geq 2$ ) and for each  $j$  a number  $\varepsilon_j \in (0, \infty)$  as well as a set  $S_{n_j} = S(n_j, \varepsilon_j)$ , contained in  $D$ , such that  $S_{n_j}$  suffices one of the following two properties (or both):




Figure 1




Figure 2

- a) There is a natural number  $k$  depending on  $j$  such that  $n_j = l_k$  and  $R_{l_k} \cap t_k = \emptyset$ .  
 b) The 'base' of  $S_{n_j}$  ( $\overline{P_2P_3}$  in fig. 1) is not a side of a triangle of the tessellation.

The numbers  $n_j$  and the associated sets  $S_{n_j}$  will be constructed by induction on  $j$ . Since  $n_j \leq n_{j-1} - 1$  for  $j \geq 2$ , some of the numbers  $n_j$  must be negative. But we assumed that each  $n_j$  is positive. Thus we get a contradiction and the theorem is proved.

The induction is performed in two steps.

### First step

We show that there is a subset  $S_{n_1}$  of  $D$  for some positive real number  $n = n_1$  which has property a) or property b).

Look at figure 2. Let  $P_1, P_2$  and  $P_3$  be the vertices of  $t_1$ . Since  $t_1$  is the smallest triangle of the tessellation, each neighbor of  $t_1$  (i.e. each triangle that shares a boundary segment of positive measure with  $t_1$ ) must be larger than  $t_1$ . It is obvious that this is only possible if there are only three neighbors of  $t_1$ , say  $t_2, t_4$  and  $t_6$ , and if (up to symmetry) they are arranged like in figure 2.

Hence there exist three triangles in the tessellation that have only a vertex in common with  $t_1$ . We can assume that these are the triangle  $t_3, t_5$  and  $t_7$  and that the triangles  $t_2, t_3, t_4, t_5, t_6$  and  $t_7$  are arranged around  $t_1$  in a clockwise manner.

Let us call  $t_i, i = 2, \dots, 7$ , the *surrounding triangles*. The vertices of the surrounding triangles that are not points of  $t_1$  may be called the *outer corners*. Clearly each surrounding triangle has exactly two outer corners. For each surrounding triangle  $t_k$ , let  $P_{2k}$  and  $P_{2k+1}$  be the outer corners written clockwise (see fig. 3). Since the triangles of the tessellation have pairwise different sizes, we get the inequalities  $P_5 \neq P_6$ ,




Figure 3

$P_9 \neq P_{10}$  and  $P_{13} \neq P_{14}$ .

We now have to distinguish the two cases whether  $t_2$  is larger or smaller than  $t_3$ .

*Case 1.* We assume that  $l_2 > l_3$ .

Consider figure 3. If  $D$  would contain a horizontal straight line whose right endpoint is  $P_7$ , the points  $P_6$  and  $P_7$  would define a set  $S_{l_3} \subset D$  with  $R_{l_3} \cap t_3 = \emptyset$ .

If this is not the case,  $D$  contains a straight line which elongates  $\overline{P_6 P_7}$  over  $P_7$  and therefore a straight line whose left endpoint is  $P_7$ , too.

If  $P_7 \neq P_8$ , the points  $P_2$  and  $P_7$  define a set  $S_{n_1} \subset D$  that has property b). So we may assume  $P_7 = P_8$ .

Now consider all systems of straight lines that may start at  $P_9$  (fig. 4). In the cases i) and ii) of figure 4 there is a set  $S_{l_4} \subset D$  with  $R_{l_4} \cap t_4 = \emptyset$ , defined by  $P_7$  and  $P_9$ . In case iii) there is a set  $S_{l_4} \subset D$  with  $R_{l_4} \cap t_4 = \emptyset$ , defined by  $\overline{P_2 P_{11}}$ ,  $\overline{P_2 P_9}$  and the elongation of  $P_8 P_9$  over  $P_9$ . Therefore we may assume that the lines starting from  $P_9$  look as in figure 4 iv).

This implies  $l_4 > l_5$ .

Two further quite similar argumentations – we now have  $l_4 > l_5$  like we had  $l_2 > l_3$ , and we will get  $l_6 > l_7$  – show that if there is not a set  $S_{n_1} \subset D$  with property a) or b), the conditions  $P_{11} = P_{12}$  and  $P_4 = P_{15}$  must be fulfilled. Moreover, the line systems starting at  $P_5$ ,  $P_9$  and  $P_{13}$  are of type iv) of figure 4 (see fig. 5).

Let  $d_1$ ,  $d_2$  and  $d_3$  denote the (positive) lengths of  $\overline{P_5 P_6}$ ,  $\overline{P_9 P_{10}}$  and  $\overline{P_{13} P_{14}}$ , respectively. Then each  $d_i$ ,  $i = 1, 2, 3$  is a sum of sidelengths of some triangles of the tessellation. Therefore  $d_i > l_1$  for  $i = 1, 2, 3$ . Hence  $d_1 + d_2 + d_3 > 3 l_1$ .

Comparing the sidelength of the seven triangles  $t_1$ ,  $t_2$ ,  $t_3$ ,  $t_4$ ,  $t_5$ ,  $t_6$  and  $t_7$  yields the equations  $l_2 = l_1 + l_7$ ,  $l_2 = l_3 + d_1$ ,  $l_4 = l_1 + l_3$ ,  $l_4 = l_5 + d_2$ ,  $l_6 = l_1 + l_5$ ,  $l_6 = l_7 + d_3$ . They tell us that  $3 l_1 = d_1 + d_2 + d_3$  in contradiction to the last inequality.

Therefore in case 1 there must be a set  $S_{n_1} \subset D$  with property a) or b).

*Case 2.* We assume that  $l_2 < l_3$ .

If there is not a set  $S_{l_2} \subset D$  with  $R_{l_2} \cap t_2 = \emptyset$  defined by  $P_4$  and  $P_5$ , the situation at  $P_4$  can only be as indicated in figure 6. Then the points  $P_1$  and  $P_4$  define a set  $S_{l_2} \subset D$  with  $P_{l_2} \cap t_2 = \emptyset$ .

## Second step

A set  $S_n (= S_{n_j}) \subset D$  may be given that has property a) or b). It will be shown that there is a set  $S_m \subset D$  with  $m \leq n - 1$  which has property a). We then put  $n_{j+1} := m$ .




Figure 4




Figure 5



Figure 6




Figure 7a




Figure 7b




Figure 7c

After a suitable translation, rotation and reflection if necessary we can assume that the given  $S_n$  has the position shown in figure 1. With the notations of figure 1, let  $t_r$  be the triangle of the tessellation which 'stands on the base of  $S_n$  at the left corner', which means that  $t_r$  is defined by the three conditions  $P_2 \in t_r$ ,  $R_n \cap t_r \neq \emptyset$ ,  $\overline{P_2P_3}$  contains a side of  $t_r$  (see fig. 7a). The vertices of  $t_r$  may be  $P_2$ ,  $P_5$  and  $P_6$ , where  $P_5 \in \overline{P_2P_3}$ . Since  $S_n$  has property a) or b),  $P_3$  is different from  $P_5$ .

Let  $t_s$  denote the neighbor of  $t_r$  at the side  $\overline{P_5P_6}$  such that  $P_5 \in t_s$ . We now have to distinguish whether  $l_s$  is greater or less than  $l_r$ .

*Case 1.* If  $l_s > l_r$ , the points  $P_2$  and  $P_6$  define a set  $S_{l_r} \subset D$  with  $R_{l_r} \cap t_r = \emptyset$  (see fig. 7b). Moreover,  $l_r \leq n-1$  because  $S_n \subset D$  and  $l_1 = 1$ . So we can define  $m = l_r$ .

*Case 2.* If  $l_s < l_r$ , let  $P_7$  denote the vertex of  $t_s$  that is not a point of  $t_r$  and let  $P_8$  denote its third vertex (fig. 7c). An examination of all possible line systems starting at  $P_7$  (they are listed in fig. 4) shows that in each case there is a set  $S_{l_s} \subset D$  with  $R_{l_s} \cap t_s = \emptyset$ , either defined by the points  $P_5$  and  $P_7$  or by  $P_7$  and  $P_8$ . Moreover,  $l_s < l_r \leq n-1$ . So we can define  $m = l_s$ .

Hence the induction step is completed, q.e.d.

Karl Scherer, Universität Kaiserslautern

## REFERENCES

- 1 W. Tutte: Squaring the Square. Can. J. Math. 1950, 197-209.
- 2 Ross Honsberger: Ingenuity in Mathematics. Mathematical Association of America 23, Yale University, 1970.

## Quelques considérations concernant le problème de l'aiguille de Buffon dans l'espace euclidien $E_n$

0. Soit  $E_n$  l'espace euclidien à  $n$  dimensions de coordonnées  $x_1, \dots, x_n$ .

La mesure élémentaire cinématique dans  $E_n$ , invariante par rapport au groupe de mouvements euclidiens, est [1]:

$$dK = dP \wedge dO_{n-1} \wedge \dots \wedge dO_1, \quad (1)$$