

Note on packing of 19 equal circles on a sphere

Autor(en): **Tarnai, Tibor**

Objektyp: **Article**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **39 (1984)**

Heft 2

PDF erstellt am: **10.08.2024**

Persistenter Link: <https://doi.org/10.5169/seals-38013>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

ELEMENTE DER MATHEMATIK

Revue de mathématiques élémentaires – Rivista di matematica elementare

*Zeitschrift zur Pflege der Mathematik
und zur Förderung des mathematisch-physikalischen Unterrichts*

El. Math.

Band 39

Nr. 2

Seiten 25–56

Basel, 10. März 1984

Note on Packing of 19 Equal Circles on a Sphere

Consider the problem of determining the largest angular diameter a_n of n equal circles (or spherical caps) which can be packed on the surface of a sphere without overlapping. For $n = 20$ van der Waerden [4] has published an arrangement as conjectured solution with circle diameter $a_{20} = 47^\circ 26'$. The graph of this packing may be seen in a simplified stereographic projection in Fig. 1. The vertices of the graph are the centres of the spherical circles and the edges of the graph are great-circle arcs joining the centres of the touching spherical circles. Goldberg [1] has thought that after removing an appropriate circle from van der Waerden's arrangement the packing of the remaining 19 circles can be improved. Goldberg has come very near to finding a correct improved packing, however, he has probably overlooked something and obtained faulty results. In [2], this mistake has been mentioned but not corrected. Therefore, at present, no packing is known for 19 circles better than that obtained by removing a circle from van der Waerden's packing of 20 circles.

The aim of this paper is to correct Goldberg's results and to present a packing of 19 equal circles on a sphere, in which a_{19} is greater than a_{20} due to van der Waerden.

In fact, two errors are made in [1]: first, to sharpen van der Waerden's result a_{20} to $47^\circ 24' 51''$, though its correct value is $a_{20} = 47^\circ 25' 51.7''$; second, to compute the circle diameter a_{19} from such an arrangement which contains some overlapping circles. This second can be shown, e.g., in the following way.

Goldberg has discovered that by removing the isolated points M and N from the graph in Fig. 1 and reflecting the vertices A, C, E and G, I, K in the great-circle arcs FB, BD, DF and LH, HJ, JL , respectively, a packing with the same circle diameter can be obtained for $n = 18$. It is obvious that a packing for $n = 19$ is obtained with the same circle diameter if this process is only executed on one of the two hemispheres (Fig. 2). Goldberg has considered the graph in Fig. 2 with additional edges GN, IN, KN and obtained that the circle diameter in this arrangement is $a_{19} = 47^\circ 25' 22''$. Since the graph in Fig. 2 is rigid in Danzerian sense [3] the edges GN, IN, KN cannot be added to the graph by moving the graph and preserving all of its edges. The removal of the isolation of the isolated point N with preservation of the




Figure 1




Figure 2

three-fold rotational symmetry of the graph can be done only by removing some of the edges of the graph in Fig. 2 and with decreased edge-lengths. It can be done, e.g., as shown in Fig. 3 where $a_{19} = 47^\circ 25' 20.2''$, or in Fig. 4 where $a_{19} = 47^\circ 25' 21.4''$ and the graph is obtained by moving the graph of Fig. 3. Thus, this kind of process for improvement is unsuccessful.

Goldberg ([1], Fig. 4) has considered in bypassing also a further, less symmetrical arrangement of 19 circles. Now a slight modification of the graph of that arrangement indeed leads to a good packing. The improving process in this case can be the following. Let the isolated point M and the edges $AU, AO, BP, CR, ES, FT, IP, KT$ be removed from the graph of our Fig. 1. Reflect the vertex A with the edges AB, AF in the great-circle arc BF (snap




Figure 3




Figure 4

the pair of edges AB, AF through between the vertices B, F). In this way the graph in Fig. 5 is obtained, which is not rigid in Danzerian sense and so can be moved with a simultaneous increase in the length of the edges. The edge-length, that is, the diameter of the circles can be increased until the distances between points A' and C, A' and E, G and N, I and N, K and N are equal to the increased edge-length in the graph. So, five additional edges appear in the graph, which make the graph rigid in Danzerian sense and prevent any further increase in the edge-length. In this way we obtained a new packing of 19 equal circles on the sphere, at which the angular diameter of the circles is

$$a_{19} = 47^\circ 40' 33.3''.$$

The graph of the new arrangement may be seen in a simplified stereographic projection in Fig. 6. The edge-length of the graph in Fig. 6 was computed by spherical trigonometry and iteration using the fact that the arrangement has a plane of symmetry, but the details are omitted here.




Figure 5




Figure 6

Tibor Tarnai
Hungarian Institute for Building Science, Budapest

REFERENCES

- 1 M. Goldberg: Packing of 19 equal circles on a sphere. *Elemente der Mathematik* 22, 108–110 (1967).
- 2 E. Székely: Sur le problème de Tammes. *Annales Univ. Sci. Budapest, Sect. Math.* 17, 157–175 (1974).
- 3 T. Tarnai and Zs. Gáspár: Improved packing of equal circles on a sphere and rigidity of its graph. *Math. Proc. of the Cambridge Phil. Soc.* 93, 191–218 (1983).
- 4 B. L. van der Waerden: Punkte auf der Kugel. *Drei Zusätze. Math. Annalen* 125, 213–222 (1952).