

Objekttyp: **TableOfContent**

Zeitschrift: **Eclogae Geologicae Helvetiae**

Band (Jahr): **68 (1975)**

Heft 2

PDF erstellt am: **12.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Geology and Paleontology of Soldado Rock, Trinidad (West Indies)

Part 1: Geology and Biostratigraphy¹⁾

By HANS G. KUGLER²⁾ and C. M. BRAMINE CAUDRI³⁾

ABSTRACT

The little islet called Soldado Rock, off the Southwest tip of Trinidad, is but a Paleocene and Eocene olistostrome mélange in Miocene turbidites. Compared with the former interpretation of the stratigraphic sequence of beds (KUGLER 1938), showing a gap between Paleocene and Late Eocene rocks, there is now clear evidence also of the presence of the Lower Middle Eocene (or very high Lower Eocene = Boca de Serpiente Formation), and of the Lower Upper Eocene.

Soldado Rock is the type locality of 11 Smaller Foraminifera, 20 Larger Foraminifera, 54 mullusks, 2 brachiopods, 3 echinoids and 5 ostracods.

In the paleontological part of this monographic study¹⁾ forty-nine species and varieties of Larger Foraminifera from the Paleocene, early Middle Eocene and Upper Eocene of the Soldado Rock are described, ten of them new: *Operculinoides trinitatis* var. *granulata*, *Operculinoides spiralis*, *Neodiscocyclina mauryae*, *Asterocyclus soldadensis*, *Amphistegina undecima*, *Amphistegina pauci-septata*, *Amphistegina grimsdalei*, *Lepidocyclus peruviana* var. *nana*, *Lepidocyclus pustulosa* var. *compacta* and *Lepidocyclus spatiosa*. Special attention has been paid to the stratigraphical significance of *Helicostegina soldadensis* GRIMSDALE [here renamed *Helicosteginopsis soldadensis* (GRIMSDALE)] and to the fauna and general nature of the Boca de Serpiente Formation, which at the moment is known only from this islet, in the form of a remnant slump mass. The recognition of heavy reworking in the upper part of the section has changed older ideas about the vertical range of several of the species described before.

CONTENTS

Introduction	366
A. The Soldado Formation	369
Bibliographic history	369
Type locality and type section	373
Bed 1	373
Bed 2	374

¹⁾ Part 2: "The Larger Foraminifera" will appear towards the end of this year, in Eclogae geol. Helv., Vol. 68, No. 3.

²⁾ Naturhistorisches Museum, Augustinergasse 2, 4051 Basel, Switzerland.

³⁾ Avenue des Alpes 29, 1814 La Tour de Peilz, Switzerland.

Remnant blocks of the Soldado Formation, not directly connected with Beds 1 and 2	378
Dasyclad algae limestone	379
Unidentified fossiliferous limestone	380
The Athecocyclina limestone	380
The Ranikothalia limestone	382
The Neodiscocyclina grimsdalei limestone	384
Reworking of the Paleocene fauna into the younger sediments	385
Occurrence of equivalents of the Soldado Formation	385
Trinidad	386
Other Caribbean islands, Venezuela and elsewhere	386
B. The San Fernando Formation	388
Bed 3	388
Bed 4	393
Bed 5	396
Bed 6	396
Bed 7	397
Bed 8	398
Bed 9	399
Bed 9a ("Asterocyclina marl")	400
Bed 10	404
C. The Boca de Serpiente Formation	411
Bed 11	411
D. Undetermined formation	420
Bed 12	420
E. Miocene beds	420
F. Recapitulation of the sample lists of various beds	422
G. Annotations to the distribution chart	426
H. Conclusions: the geological history of Soldado Rock	427
Acknowledgments	430
References	430

Introduction

The Soldado Rock stands sentinel in the strong tidal currents of the shallow Southern passage, called the Serpent's Mouth, between Trinidad and the South American mainland, about 10 kilometers West of Columbus Bay on the Cedros Peninsula.

The Rock comprises an area of about 6500 square meters. From its Northern point, which carries the trigonometrical signal, and which is 36 meters above sea level, one can see the flat Pedernales area at the Northern end of the Orinoco delta. The coordinates of this point are: Latitude $10^{\circ} 04' 24''$, Longitude $62^{\circ} 00' 56''$. To the South of the highest point there follows a saddle of softer beds and then another, lower, elevation. During the breeding season the intruder is greeted by the shrieks of terns and other guano producing birds. The guano and its resulting phosphate of lime cover the island from the top to the high-water level.

Though Amerindians, and later on fishermen of Trinidad, regularly collected eggs during the breeding season of the terns, one cannot expect anyone to visit the inhospitable Soldado Rock excepting biologists, surveyors and geologists. The first geologist to have visited Soldado Rock known to us was V.C. Veatch, whose collection of fossil shells, together with some stratigraphic details, was described by MAURY in