

Acknowledgments

Objekttyp: **Chapter**

Zeitschrift: **Eclogae Geologicae Helvetiae**

Band (Jahr): **87 (1994)**

Heft 1

PDF erstellt am: **27.06.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

(Fig. 14, 20 and 21). The sharp-based HCS calcarenites were probably formed under very rare, unusually stormy conditions that caused a strong oscillatory flow regime near the bottom during the Middle and Upper Toarcian. Characteristic abrasion surfaces on some benthic foraminiferal tests seem to be due to oscillatory conditions at the sediment/water interface during the deposition of the winnowed beds (Fig. 13). The minimum depth and sedimentation rate of the depositional environment was reached in the upper part of the Toarcian and corresponds roughly to an outer/middle shelf environment, near the major storm wave base (Fig. 21). Moreover in the Erbaense-Meneghinii Zones the succession is strongly condensed – testified by repeated hardgrounds – and represented by sediments only 7.5 m thick, in comparison with the *Tenuicostatum*-*Bifrons* Zones where deposits 60 m thick occur.

9.4 *Tectonics and eustacy*

The microforaminiferal assemblage BC present at the boundary between Corniola and Marne del M. Serrone is lacking in the Valdorbia area where *Glomospirella* disappears in the Upper Domerian, earlier than in the other areas. At the Domerian/Toarcian boundary slumps, mass-flow and calcareous turbidites occur. These sedimentary features can be interpreted as indicative of regional synsedimentary tectonics rather than a eustatic lowstand phase because this sedimentological character seems to be relatively local. Hence, the Domerian regressive stage expressed in the eustatic curve of Hallam (1988) is not evident in this area, probably because of local tectonic activity of M.Catria-Valdorbia area (Fig. 21).

The deepening found in the *Tenuicostatum* Zone can be connected to sea-level rise (Hallam 1967), according to the Jurassic eustatic curve of Hallam (1988), and/or to an increase in the rate of the subsidence (Fig. 21). In fact the degree of reworking reaches its maximum intensity in the Lower Toarcian.

The shallowing trend suggested for the Middle/Late Toarcian fits better into a geological context clearly affected by a regressive phase (Hallam 1988), than into one affected by tectonic activity (Fig. 21). In fact a regressive-shallowing can be considered to be widespread in the central Apennines, as in the Umbria-Marche basin and the Lazio-Abruzzi carbonate platform area (Giannini et al. 1970; Colacicchi & Bigozzi 1992). Mass-flow deposits which occurred in the Lower Aalenian are widely scattered in the Umbria-Marche area (M. Cucco, M. Serrone, Narni-Amelia ridge, M. Martani) and seem to reflect regional sea-floor instability. The cause of these features is still uncertain, although the Aalenian regression is probably the result of Western Tethys tectonics (Hallam 1988).

Acknowledgements

Our best thanks to Prof. F. Venturi who has provided us with his data on ammonite horizons. We are very grateful to Dr. R. Rettori, University of Perugia, for help in microfacies description and to Dr. C. Arias, University of Madrid, for the ostracod classification. We would like also to express our thanks to A. Bartolucci and G. Vinti, University of Perugia, for their technical assistance and to G. Tosti for his help with the photography. Work published with the financial support of M.U.R.S.T. (40%, Prof. R. Colacicchi) and by the Project PB-92-0960 (D.G.I.C.Y.T.- Spain).