

**Zeitschrift:** Eclogae Geologicae Helvetiae

**Band:** 89 (1996)

**Heft:** 1

**Artikel:** Western Préalpes Médiannes Romandes : timing and structure : a review

**Autor:** Mosar, Jon / Stampfli, Gérard M. / Girod, François

**DOI:** <https://doi.org/10.5169/seals-167907>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 15.10.2024

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Western Préalpes Médiannes Romandes: Timing and structure. A review.

JON MOSAR, GÉRARD M. STAMPFLI & FRANÇOIS GIROD<sup>1</sup>

*Key words:* Préalpes Médiannes, klippen, Briançonnais, folds, thrusts, tear faults, décollement, thin-skinned tectonics, synsedimentary tectonics, alpine tectonics, passive margin, active margin, sedimentation, transported metamorphism, internal deformation

### ABSTRACT

Between the original position and their present day location as klippen, the Préalpes Médiannes underwent a complex history of paleotectonics and alpine tectonics. Due to the opening of the Piemont ocean the Briançonnais sedimentation realm of the Préalpes Médiannes evolved as a rim basin of the northern passive margin during Jurassic to Eocene times. Different paleotectonic features (normal faults, synsedimentary growth structures, inversion structures) developed and were active above a basal detachment in evaporitic layers. The tectonic movements were a consequence of thermal events in the crust. Isolated from the Iberic continent at the end of the Late Cretaceous, the Briançonnais exotic terrain was incorporated into the accretionary prism of the closing Piemont ocean and the incipient alpine orogeny during the Lutetian-Bartonian. The Préalpes Médiannes were detached from their homeland during the Bartonian-Priabonian and were transported onto the foreland. The tectonic style is one of a thin-skinned foreland fold and thrust belt. Fault associated fold development above a main décollement, together with internal deformation, represent the Préalpes Médiannes main structural features. The very low-grade metamorphic conditions have their origin in the heat flux induced by tectonic burial by overriding nappes in the accretionary prism. After having been transported on top of the developing Helvetic nappes the Préalpes were emplaced in their present day position in front of the Alpine mountain belt during Oligocene times. Post-emplacment and out of sequence thrusting, possibly younger than Oligocene, is observed and can be related to thrusting in the sedimentary substratum and the basement.

### RESUME

Entre leur position de départ et leur situation actuelle en tant que klippe tectonique les Préalpes Médiannes ont subi une histoire complexe de paléo-tectonique et tectonique alpine. Suite à l'ouverture de l'océan Piemontais, le domaine de sédimentation des Préalpes Médiannes se développe en forme de bassin frangeant le long de la marge passive nord du Briançonnais entre le Jurassique et l'Eocène. Différents types de structures paléotectoniques (failles normales, failles inverses, structures de croissance synsédimentaire) se développent au-dessus d'un décollement basal dans les niveaux évaporitiques à la base des Préalpes Médiannes. Ces mouvements sont les conséquences de réajustements occasionnés par les événements thermiques crustaux. Une fois isolé du continent Ibérique par une faille transformante sénestre N-S durant le Crétacé terminal, le terrain exotique Briançonnais vient s'insérer dans le prisme d'accrétion qui voit le jour suite à la fermeture de l'océan Piemontais dès le Lutétien-Bartonien. Le métamorphisme faible résulte d'un réchauffement du à l'effet d'enfouissement tectonique des Préalpes Médiannes dans le prisme d'accrétion pendant le Bartonien-Priabonien. Elles sont ensuite détachées de leur patrie et mises en place sur les nappes Helvétiques en pleine formation, puis transportées sur l'avant-pays alpin, où elles arrivent pendant l'Oligocène. Le style structural est celui d'une tectonique «pelliculaire» caractéristique d'une chaîne d'avant-pays plissée. La déformation interne des roches

---

<sup>1</sup> Institut de Géologie et Paléontologie, Université de Lausanne, UNIL-BFSH 2, CH-1015 Lausanne


va de paire avec la formation de rampes, plis et plis-failles. Les chevauchements intervenus après la mise en place, probablement post Oligocène, sont associés aux déformations et à la tectonique intervenant dans les sédiments du substratum et de leur socle.

## Introduction

The Préalpes are formed by several allochthonous tectonic klippen resting on the N and NW central European alpine foreland, in front of the Helvetic nappes and the external crystalline basements of the Alps (Schardt 1898; Jeannet 1922; Trümpy 1960; Badoux & Mercanton 1962; Caron 1973; Plancherel 1979; Fig. 1). Several klippen can be recognized from E of Lucerne (Mythen klippe) to Annecy (France) in the W (Annes klippe). The largest are the Préalpes Romandes NE of lake Geneva in Switzerland and the Chablais Préalpes S of Lake Geneva in Switzerland and France. Within the klippen several tectonic units with different paleogeographical origins can be identified. One can differentiate from top to bottom (Fig. 1): [i] the Nappe Supérieure which itself is subdivided into four different units: the Gets nappe, the Simme nappe, the Dranses nappe and the Gurnigel nappe (Caron 1972, 1976; Trümpy 1980); [ii] the Breccia nappe, resting on the trailing part of the Préalpes Médiannes only (Lugeon 1896; Jaccard 1904; Schroeder 1939; Weidmann 1972; Steffen et al. 1993); [iii] the Préalpes Médiannes nappe and [iv] the Nielsen nappe which exists in the Préalpes Romandes only and which today forms the southernmost structural unit (McConnel 1951; Lombard 1971; Matter et al. 1980; Homewood et al. 1984; Ackermann 1986; Caron et al. 1989). At the base of the Préalpes Médiannes and separating them from the Nielsen nappe, is the "Zone Submédiane" structural unit (Weidmann et al. 1976). Elsewhere the sole of the Préalpes is in contact with the underlying units (the Helvetic nappes in the S and the Subalpine Molasse and flysch to the N) by means of the "Ultrahelvetic" (Badoux 1963; Homewood 1977; Jeanbourquin 1991, 1992, Jeanbourquin & Goy-Eggenberger 1991; Jeanbourquin et al. 1992; Lempicka-Münch & Masson 1993) which are characterized by mélanges and cornieules (Rauhacken) associated with anhydrites.

It was Schardt (1884; 1893a; 1898) who first clearly demonstrated that the Préalpes were allochthonous (see also Lugeon 1902). He also recognized and described the fold and thrust style of the Préalpes Médiannes (Favre & Schardt 1887), and concluded that there was a genetic link between thrusts and folds. Historical accounts of the development of ideas and geologic research in the Préalpes are given in Jeannet (1912–1913), Bailey (1935), Masson (1976) and Lemoine (1988); more than 800 references on the Préalpes can be found in Mosar & Borel (1995).

In terms of paleogeography the Préalpes Médiannes formed part of the Subbriançonnais and Briançonnais sedimentation realm (Trümpy 1960; Caron 1972, 1973; Bernoulli et al. 1979; Caron et al. 1980a, 1980b; see general introduction Boillot et al. 1984; Lemoine et al. 1986; Debelmas 1989). Equivalent stratigraphic units have been found both in the Préalpes Médiannes and the Pennine nappes south of the Rhône valley. In these units of the Grand Saint Bernard nappe, the Siviez-Mischabel nappe has been recognized as homeland for the Médiannes Rigides and the Pontis nappe for the Médiannes Plastiques (Schardt 1907; Ellenberger 1952; Baud & Septfontaine 1980; Sartori 1987, 1990; Escher et al. 1993; Sartori & Marthaler 1994). The Préalpes Médiannes nappe was detached from its basement and underwent thin-skinned tectonics during the Alpine



B = Bex; C = Corbeyrier; CC = Col de Chaude; CD = Châtel St. Denis; Ch = Charmey; CM = Col des Mosses; CO = Château d'Oex; Cs = Chamossaire; D = Dorena; DB = Dent de Broc; DC = Dent de Corjon; G = Gruyère; GE = Grande Eau; Gf = Gummfluh; Gp = Gorges du Pissot; Gt = Gastlosen; J = Jaun; L = Leysin; LM = Les Millets; M = Moléson; MD = Mont d'Or; Mo = Montbovon; MP = Mont Pélerin; N = Niremout; O = Ollon; R = Les Rodomonts; Rb = Rübli; RN = Rochers de Naye; Ro = Rossinière; StT = Saint Triphon; TA = Tours d'Ai; VN = Vanil Noir

Fig. 1. Tectonic map of the western Préalpes Romandes and their neighbouring structural units with simplified location-map of the Chablais and Romandes Préalpes (in black) in the Alpine mountain belt. Although the Nappe Supérieure of the Préalpes is made of four distinct tectonic units, only the Gurnigel nappe has been shown with a different pattern. UMM = Lower Marine water Molasse; USM = Lower Fresh water Molasse; OMM = Upper Marine water Molasse; OSM = Upper Fresh water Molasse.

orogeny. The basal décollement occurred in evaporites located at the base of the Middle and Late Triassic. The Zone Submédiane and Niesen nappe paleogeographic homeland is on the northern European margin of the Valais ocean (Stampfli 1993; Stampfli & Marchant in press). The Breccia nappe has its origin on the north Piemontais margin and represents mainly a synrift deposit of that margin. The Nappe Supérieure comes from the Piemontais oceanic domain (Dranses nappe; see also controversy in Hsü 1989) or the south Piemontais active margin (Gets, Simme and Gurnigel nappes).

The Préalpes Médiannes are subdivided into Médiannes Plastiques, forming the frontal (NW) part of the nappe and Médiannes Rigides, forming the trailing (SE) part of the nappe (Fig. 2; Lugeon & Gagnebin 1941). This subdivision is based on the large scale structural geometry (folds in the Médiannes Plastiques and imbricates in the Médiannes Rigides), internal deformation (strain) is on the contrary weak and brittle in the Médiannes Plastiques and stronger and more "plastic" in the Médiannes Rigides (Mosar 1989). In the Préalpes Romandes, a domain with intermediate structural and sedimentological characteristics exists between the Médiannes Rigides and Plastiques: the Gastlosen range (Lugeon & Gagnebin 1941; Baud 1972; Plancherel 1979).


Before reaching their present day location the Préalpes Médiannes experienced successive geologic events (Fig. 3). We will focus hereafter on two major families of structural and tectonic features: [i] those associated with syn-sedimentary deformation and tectonics, such as rift development, sedimentary growth faults, inversion structures etc., [ii] those associated with the alpine orogenesis s.str., starting during Late Eocene, after the Préalpes Médiannes flysch sedimentation.

### **Paleo-tectonics and sedimentology**

Sediments in the Préalpes Médiannes range from Triassic to Paleogene (Eocene) in age (Fig. 4, Badoux 1962a; Plancherel 1979, 1990; Baud & Septfontaine 1980; Trümpy 1980; Baud et al. 1989). They are formed by limestones, dolomites, marls and shales. The only basement relics in the Préalpes themselves are Permo-Carboniferous sediments, thought to belong to the Préalpes Médiannes. They are found in the SW Chablais Préalpes near Tanninges (Septfontaine & Wernli 1972; Wernli & Broennimann 1973). The Préalpes Médiannes basement presently found in the Siviez-Mischabel and Pontis nappes (middle Pennine) is formed by polycyclic rocks as well as monometamorphic rocks (Thélin et al. 1993; Sartori & Marthaler 1994). The polymetamorphic basement is considered Proterozoic to early Paleozoic in age and has suffered ante-alpine metamorphism. The monometamorphic basement is Carboniferous to Early Triassic in age and is probably derived from several distinct basins (Thélin et al. 1993). The basement of both the Siviez-Mischabel and the Pontis nappes formed one single continuous substratum.

---

Fig. 2. Cross-sections through the western Préalpes Médiannes Romandes. (a) The profile in the Sarine valley (A-B, location on Fig. 1) shows the formation of a trough-like structure (very deep syncline) beneath the Sarine syncline and the deepening of the basal décollement by some 2000 m. (b) The section along the Rhône valley (C-D, location see Fig. 1) shows the shallow depth of the basal décollement and top-to-the-W/WNW movements along thrusts surfaces in the frontal imbricates. In both sections backthrusts are remarkable (indicated by -xx-).



Tectonic event table for the Préalpes médianes




Fig. 3. Time table indicating the different tectonic (synsedimentary and alpine) and "metamorphic" events and their most likely development in time. Detailed discussion is provided in the text. Geological time scale by Odin (1994).

Two major sedimentation realms clearly differentiated since the Liassic: to the N-NW, in the Médianes Plastiques, a large basin is marked by an important thickness of sediments (Fig. 4). To the south this subsiding domain turns into a ramp (associated with a continuously active structural high, the Château d'Oche-Corbeyrier zone; Septfontaine 1983) that gives way since the Dogger to a platform and lagoonal environment in the Médianes Rigides (Baud & Septfontaine 1980). The sedimentary record is by no means complete nor continuous and is quite different in the two major domains. Sedimentary gaps are most important during Liassic times and in the Cretaceous-Tertiary period where hardgrounds may develop. The gaps may either reflect periods of non-deposition or of erosion. The reader can find more detailed descriptions and discussions of the various formations of the Préalpes Médianes in:

Baud (1972; 1984) for the Triassic;

Mettraux (1989), Mettraux & Mosar (1989) and Thury (1973) for the Early Jurassic (Lias);

Septfontaine (1983), Furrer & Septfontaine (1977) and Furrer (1979) for the Middle Jurassic (Dogger);




Fig. 4. Stratigraphic table showing the times of deposition and sedimentary gaps (non-deposition and/or erosion) in the Préalpes Médiannes Plastiques and Rigides, with the different formation names (references given in the text). Geological time scale by Odin (1994). The simplified palinspastic reconstruction shows the main paleogeographic regions of the Préalpes Médiannes (modified from: Baud & Septfontaine 1980).



Weiss (1949), Isenschmid (1983), Heinz (1985) and Heinz & Isenschmid (1988) for the Late Jurassic ("Malm");  
Boller (1963) for the Early Cretaceous (Neocomian);  
Caron & Dupasquier (1989) and Python-Dupasquier (1990) for the "Middle" Cretaceous;  
Guillaume (1986) for the Late Cretaceous-Tertiary and Flück (1973) and Caron et al. (1980b; 1989) for the Préalpes Médiannes flyschs.

The Préalpes Médiannes sedimentation realm has been interpreted in recent years in terms of a rim basin; the Sub-briançonnais (Médiannes Plastiques p.p.) and the Briançonnais (Médiannes Rigides and Médiannes Plastiques p.p.) developing in this rim basin (Stampfli & Marthaler 1990). This basin was located to the N-NW of the Briançonnais upper plate rift shoulder which separated it from the Breccia nappe sedimentation domain to the south containing syn-rift deposits. This upper plate shoulder developed as a consequence of the breakup of Pangea which gave birth to the Alpine Tethys oceanic realm (Piemont ocean; Frisch 1979; Tricart 1984; Debelmas 1989; Favre & Stampfli 1992, 1993; Stampfli & Marchant in press). During the formation of the Valais ocean to the north in Late Jurassic-Early Cretaceous, the Briançonnais domain s.l. evolves as a micro-continent forming the eastern termination of the Iberic plate. Once detached from the Iberic plate, it will be incorporated as exotic Briançonnais terrane into the oblique closure of the Piemont ocean with its accretionary prism and mélange development into which the Nappe Supérieure is also included. The subsequent continental collision and alpine orogen will be discussed in the alpine tectonic chapter.

Based on a synthetic subsidence reference curve for the Médiannes Plastiques (Girod 1995) we discuss the different stages of sedimentation and geodynamics (Fig. 5). This curve is based on the most complete and the thickest sediment record in the Préalpes Romandes (Sarine valley; Fig. 2a). The changes in slope of the subsidence curve are compared to geodynamic events occurring in the Tethyan area at a given time. These geodynamic stages are based mainly on thermal evolution except for the final stages of continental collision. Thermal heating due to asthenospheric upwelling, i.e. in a lithospheric fracture zone, will stop any previously ongoing progressive thermal subsidence. This is marked by a "plateau" (or even a slight bulge) in the subsidence curve. The possible oceanisation following this thermal expansion is thought to be expressed on the developing passive continental margins by an initially rapid and then a slower cooling. The subsidence curve will show a steeper slope. The ocean closure and the subsequent collision in an active margin cause a flexural response of the subducting plate in addition to the isostatic re-equilibration. On the subsidence curve this will show as steepening of the slope. General uplift results in a positive slope of the subsidence curve.

The different sedimentary facies and paleotectonic events can thus be integrated into a geodynamic **interpretation** (model) for the plate-tectonic evolution of the Briançonnais s.l. micro-continent. This interpretation is part of a larger scale attempt to re-evaluate the plate tectonics of the Tethyan domain. We chose to integrate data from the Préalpes into this model, aware of its debatable nature.

[i] The platform carbonates that develop during the Triassic period are associated with the thermal subsidence which may be seen as resulting from the opening of the Hallstatt-Meliata ocean (Fig. 5; Stampfli, this volume for references, Stampfli & Marchant in




Fig. 5. Synthetic subsidence curves for the Préalpes Médiannes sedimentation domain. The synthetic curve is based on data from the Médiannes Plastiques basin in the Sarine valley area. The curve corresponds to the maximum tectonic burial curve after decompaction. Circled values refer to the different synsedimentary tectonic periods as described in the text. Geological time scale by Odin (1994).

Table shows the values used for constructing the subsidence curve with the program "Backstripp89" by R. Schegg. Thick. = sediment thickness in meters as observed or estimated. (x) indicate periods of eroded material (1) = 2000m eroded, (2) = 2m eroded, (3) = 0m eroded (no deposit), (4) = 10m eroded, (5) = 40m eroded; Tmin/max = minimum and maximum water depth of formation top; Bmin/max = minimum and maximum water depth of formation bottom. For age attribution and complete names of the different formations see Fig. 4.

press). Deposited in a steadily subsiding, stable shallow water environment, these sediments are the only ones in the Préalpes Médiannes where a consistent signature of the interplay with eustatic variations is recorded (Baud 1984).

[ii] A break in the synthetic subsidence curve (Fig. 5-A) is shown by a “plateau” during the Rhetian-Liassic-Early Dogger period (Fig. 5A-B). This event is most likely associated with isostatic rebound related to the initial stages of rift development that eventually leads to the Piemont ocean opening. In our model the ongoing oceanic opening will be followed by extensive thermal subsidence, mainly during Dogger times.

At the beginning of the oceanic opening, during the Rhetian and Liassic, a series of synsedimentary normal faults develop. This ENE-WSW trending fault-system subdivides the Préalpes Médiannes into several distinct ENE-WSW oriented sedimentation domains. They are associated with a N-S trending fault-system developing important scarps yielding breccias (Mettraux & Mosar 1989). Fast differential subsidence starts in the southern Préalpes Médiannes during the Rhetian, with the associated depocenters subsequently migrating northward. The different small basins that develop are separated by structural highs (Fig. 4), the most remarkable of which is the Château d'Oche-Corbeyrier high (Septfontaine 1983). To the S of this structure develops a deep basin filled by the Sinemurian to Bajocian Heiti Formation (including the Tours d'Aï area) and further S is the Médiannes Rigides platform domain. To the N two shallow highs separate three smaller basins. Further to the north three tectonic imbricates (located today in the frontal Médiannes Plastiques) form three small domains associated with normal faults (Mosar 1994).

During this period we also see the formation of karst in the Liassic sediments (Mettraux 1989; Mettraux & Mosar 1989) and in the underlying Triassic limestones (Genge 1957; Baud & Masson 1975). Breccias of Late Liassic – Dogger age, indicative of strong erosion, form at the basin/platform transition (ramp) (Baud et al. 1989). Onlaps and erosion are common (Mettraux & Mosar 1989).

A generalized and continuous subsidence is well recorded in the Médiannes Plastiques basin (Fig. 5B-C). These sediments are in excess of 1500–1800 m thickness (W of the Vanil Noir in the Sarine Valley, Préalpes Romandes). Important thickness changes occur both perpendicular to the E-W basin axis and along strike. Responsible for these variations are small en échelon synsedimentary listric normal faults. Because of the small size of the basins created by these faults (10 km<sup>2</sup>) we suggest that the faults above which they originate are all linked to a “basal” décollement located in the Triassic evaporites. This detachment above which the Préalpes Médiannes rim basin (and the listric normal faults) evolves is most likely gravity induced (Bally 1981) due to the small northward slope created by thermal expansion and related rift shoulder uplift to the S. According to this concept, the basement-rooted faults which controlled the Permo-Carboniferous troughs did not significantly affect local Préalpes Médiannes paleo-tectonics, unlike proposed so far (Baud & Septfontaine 1980, Sartori 1987, 1990; Mettraux & Mosar 1989; Sartori & Marthaler 1994; Septfontaine 1995). The proposed “model” allows us to explain locally compressional synsedimentary structures such as progressively rotated (up to 100°) sedimentary unconformities (synsedimentary growth folds: Suppe et al. 1992; Septfontaine 1995) developing on a structural high to the N of an inverted listric fault of the Liassic Heiti Formation basin, without invoking reactivation of basement, when the general tectonics are extensional and the general trend is thermal subsidence. Growth structures on faults and folds, fan shaped sedimentary wedges, strong onlaps, slumps and resediments

are but some of the more obvious sedimentation structures due to syndepositional tectonics during this period. Although interpreted as resulting from the combined movement along normal listric faults above a major décollement, the synsedimentary structural high of the Château d'Oche-Corbeyrier zone can be compared with the structural high described in the Schams nappe equivalent of the Briançonnais terrain in E Switzerland (Schmid et al. 1990). The authors describe this zone as resulting from transpressive movements forming a flower structure rooted in the basement. During this stage, synsedimentary tectonics are related to the progressive oceanisation with fault development during the Rhetian and Liassic followed by thermal subsidence and gravity induced basin development above shallow normal faults during the Dogger. Transpressive events could mark the onset of rifting in the Valais ocean in mid to late Jurassic times (Late Oxfordian; Stampfli 1993).

Beginning with the Dogger only one large basin remains to the N of the Château d'Oche-Corbeyrier structural high (Fig. 4: "Cancellophycus basin"), separated by a sedimentary ramp from a lagoon and platform to the S (Médiannes Rigides; Septfontaine 1983). Progressive onlaps of sediments from the N onto this platform (Malm onto Triassic and Late Cretaceous onto Malm) and the formation of cuestas (Septfontaine 1995) show that this domain has constantly suffered from erosion and non-deposition. This is indicative of the proximity to a rift shoulder in the south (S of Médiannes Rigides domain; between Briançonnais and "Ultra-briançonnais" domains). To the S of this rift shoulder the Breccia nappe sedimentation is dominated by breccia deposits from Sinémurian to late Dogger (Jaccard 1904; Schroeder 1939; Gagnebin 1940; Weidmann 1972; Steffen et al. 1993).

[iii] Another important break in the subsidence curve (Fig. 5C) marks the beginning of the "Callovo-Oxfordian phase" of Septfontaine (1983). The "plateau" developing during the final Dogger and Malm and lasting for about 20 Ma indicates a period of little tectonic subsidence (Fig. 5C-D). Malm limestones vary in thickness between 30 and 300 m. Platform carbonates develop to the S in the Médiannes Rigides, condensed facies are observed in the transition zone (sedimentary ramp; Château d'Oche-Corbeyrier zone) with the basinal sediments of the Médiannes Plastiques (Isenschmid 1983; Heinz 1985; Heinz & Isenschmid 1988). During this period thickness changes in the Malm layers could be enhanced by differential compaction and be associated with listric faults (f.i. former Permian basins or Heiti Formation basin; Girod 1995).

The geodynamic change leading to the development of the Malm "plateau" in the subsidence curve may be explained by a thermal influx due to the opening of the Valais ocean. This rifting separates Europe to the N from the Iberian plate to the S, the Briançonnais domain thus becoming a peninsula. The thermal influence of the Valais ocean formation combined with the thermal subsidence associated with the Piemont ocean phase, lasts for most of the Cretaceous until the Campanian (Fig. 5D-E). The continuing basin infill in the Médiannes Plastiques is documented by the 100–150 m thickness of the Early Cretaceous Calcaires plaquetés Formation (Boller 1963). Present in the Médiannes Plastiques basin this formation pinches out towards the S in direction of the Médiannes Rigides platform along a slope close to an ancient coastline. The pelagites and hemipelagites of the Middle Cretaceous Intyamon Formation (Caron & Dupasquier 1989; Python Dupasquier 1990) are followed by the Rote Platte Formation of Late Cretaceous age (Guillaume 1986). The latter is transgressive on the Calcaires plaquetés For-






Fig. 6. Middle to Late Eocene palinspastic map of the Alpine region. The Briançonnais (Br) is becoming an isolated terrain being pushed onto the South-Helvetian (S-H) continental margin. The Helvetic domain (He) represents a rim basin along the European continental margin. The Austro-Alpine plate and its Carpatho-Dinaric continuation is being spliced into several segments or terrains incorporated into active margins. The Austroalpine terrain represents the former passive margin of the Apulian plate, already inverted in early Cretaceous. Subduction is also active under the Iberic plate.

Fig. 7. a) Present day cross-section of the N-E Pacific active margin modified from the Lithoprobe transect of southern Vancouver Island (Fuis & Clowes 1993). The 400 °C gradient is taken from van den Beukel (1990). The position of this gradient is indicated for different ages of oceanic lithosphere entering the accretionary prism (25, 10, 3 and 0 Ma). Widespread amphibolitic metamorphism and related hydrothermalism can be expected in the prism when young oceanic crust is being subducted (i.e. when 400 °C gradient crosses from the mantle to the crustal rocks of the descending slab).

b) Mid to Late Eocene (~45-40 Ma) pre-collisional stage of the Alpine orogen (location of cross-section in Fig. 6). For the upper plate the Vancouver Island transect has been used as an actualistic reference frame in order to compare the thermal behaviour of the subducted terrains. Size geometry and juxtaposition of the alpine elements in this upper plate is therefore schematic. In Eocene time the Briançonnais terrain has been detached from its lithosphere and lower crust substratum and pushed onto the North Valais ocean margin. The Briançonnais was first covered by flysch and then by the accretionary prism of the closing Piemont ocean (Gers-Tsaté prism). The Gurnigel flysch (GURN) is the witness of the obduction of this prism but was not necessarily located where shown in the figure. The lower plate represented by the Helvetic margin is affected mainly by changing flexuration behaviour related to the buoyancy of the lithosphere being subducted.



mation in the N, further S on the Intyamou Formation and in the Médiannes Rigides on the Calcaires massifs Formation of the Malm.

At the very end of this stage follows a period mainly marked by non-deposition and/or erosion that ends only with the Forclettes Formation of the very final Late Cretaceous. Due to the ongoing geodynamic development of the Valais ocean and the Briançonnais peninsula development, the Préalpes Médiannes sedimentation realm is cut from its regular sediment supply and is both starving and drowning. Variations in the subsidence curve steepness indicate changes in thermal subsidence rate as for example around 70 Ma (Fig. 5).

It is during this period that the Briançonnais peninsula is separated to the W from its Corso-Sardinia-Iberia motherland and evolves into an isolated terrain (Maury & Ricou 1983; Ricou & Siddans 1986). At the same time the northern Valaisan margin receives sediments (Niesen nappe flysch) from an uplift located along the southern European margin, possibly related to inversion structures developing in the margin (Figs. 6, 7).

The fact that the Forclettes Formation sediments are deposited during a general period of starvation (period related to an "elevation" by Guillaume 1986) is most likely connected to combined effects of the complex geodynamic situation at that time. The isolation of the Briançonnais terrane may result in some isostatic uplift; the thermal influx caused by the possible subducting towards the S of the still warm Valais ocean may also cause some uplift and a new source of sediments might come from the uplifting of the shear zone between the Briançonnais and Corsica. Eustatic sea level changes as a cause for the sedimentation of the Forclettes Formation may be excluded.

[iv] Following a new 15 Ma sedimentation gap the Briançonnais microcontinent is finally incorporated into the Tsaté accretionary prism (Fig. 6, 7) developing at the expense of the closing Piemont ocean (Marthaler & Stampfli 1989). The load of the thrust units coming from the S causes the subducting plate to flex (Fig. 5E-F). It is during this period, starting in Early Eocene, that the Chenaux Rouges Formation is deposited (pelagic limestones and clays). This formation begins in the Ypresian and grades into the Préalpes Médiannes flysch formation. These are the youngest sediments preserved in the Préalpes Médiannes sedimentation realm. The portion of the synthetic subsidence curves associated with this last stage has a steeply inclined concave shape. The great depth is suggested by the fact that the red clays at the transition between carbonates and flysch were deposited below the CCD.

Except for a N-S and an ENE-WSW trending normal fault system with small displacement, no local tectonic disturbances are known from this period. The small faults (Guillaume 1986; Python Dupasquier 1990) cause local and small changes in depositional thickness. Occasionally breccias along fault scarps may be expected. These breccias become quite important in the Briançonnais domain in France (Chaulieu 1992).

[v] The final stage is represented as a steep positive slope between 40 Ma and 10 Ma (Fig. 5F-G). This portion of the synthetic subsidence curve is based on the eroded sediments of the thrust Nappe Supérieure whose thickness is estimated to about 2000 m (based on studies of very low-grade metamorphism, Mosar 1988a). The Préalpes Médiannes plus the supported Nappe Supérieure and Breccia nappe were subsequently uplifted during continental collision. The uncertainties in timing of the thrusting (events taking place between 40 and 10 Ma) allowed only to construct one single straight segment for the subsidence curve during these post-sedimentation events.

## Alpine tectonics

Alpine tectonics in the Préalpes Médiannes start with the end of sedimentation during Late Eocene (Bartonian) and their complete involvement in the active accretionary prism at the front of the developing Alpine mountain belt. This corresponds to the beginning of the continental collision stage resulting from the final closure of the Piemont ocean. The Valais ocean is at this stage almost closed (hence largely subducted) and the European continental plate starts subducting beneath the accretionary prism. While most of the Briançonnais terrain suffers intense thick-skinned deformation (with basement involved) and high-grade metamorphism during this Meso-alpine phase (Steck & Hunziker 1994), the Préalpes Médiannes are detached from their substratum and suffer thin-skinned tectonics. Although these cover nappes may reach a total of some 10 km no basement is involved, they represent “cover nappes” according to the definition of Escher et al. 1993. This décollement occurs where evaporite series were present near the base of the sedimentary cover (Fig. 4). The areas devoid of this evaporite formation, such as the Barrhorn series (considered as the lateral equivalent of the Médiannes Rigides) stick to their Pennine homeland (Trümpy 1955; Sartori 1990; Sartori & Marthaler 1994). Consequently a thick-skinned tectonic regime develops “basement nappes” (Escher et al. 1993). The differences in distribution of the evaporite series is thought to be controlled by N-S trending syn-sedimentary paleofaults. Thus the Médiannes Rigides and the Médiannes Plastiques were detached from the Siviez-Mischabel nappe basement and from the Pontis nappe, respectively. They have been substituted by a new “cover”: the Mont Fort nappe (Sartori 1990; Escher et al. 1993). The main detachment is thought to occur after the arrival of the Nappe Supérieure onto the Préalpes Médiannes during the Bartonian-Priabonian, coeval or slightly postdating the cessation of sedimentation (Bartonian).


In the following we will discuss the formation of the Préalpes Médiannes fold and thrust belt first from the perspective of large-scale structural development, then from the point of view of internal rock deformation associated with the larger scale structures and finally in a more general environment of alpine tectonics.

### *Structural geology*

Three distinct structural domains can be recognized in the Préalpes Médiannes (Lugeon & Gagnebin 1941; Plancherel 1979; Mosar 1991): the Médiannes Plastiques to the N are formed by a series of large fault-related synclines and anticlines, the Médiannes Rigides to the S by several sets of large imbricates and in between lies a unit with intermediate features, the Gastlosen–Tours d’Aï chain (Fig. 1, 2a, 8).

[i] The Médiannes Plastiques are bordered to the North and the South by the Nappe Supérieure (Gurnigel nappe to the N and Simme nappe in the Corbeyrier–Château d’Oex–Simmenthal syncline). The external, northernmost zone sees the characteristic development of several, mostly three, imbricates. The fault-bend fold style is dominant although some open upright folds, as in the Moléson (Gruyère) area (Mauve 1921) and even tight overturned folds may form (Gagnebin 1922, 1924; Peterhans 1923; Mosar 1991, 1994; Weidmann 1992). The thrust ramps often terminate with one or several splays. The southernmost part of the Médiannes Plastiques, along the N limit of the Corbeyrier–Hongrin–Châteaux-d’Oex syncline, forms the transitional Dt d’Oche – Sex de la Sarse zone from the Médiannes Plastiques to the Médiannes Rigides sedimentation domain.





The basinal Cancelliphyucus Formation series from the Dogger are thinning and by means of a transitional facies (Sommat Formation, Septfontaine 1983) turning into lagoonal facies of the Mytilus Formation; the Early Cretaceous Calcaires plaquetés Formation is missing (Fig. 4).

Folds and thrusts are oriented E-W in the eastern Médiannes Plastiques Romandes and have a more NE-SW, and even N-S, orientation in the western part (Jeannet 1922; Plancherel 1979). The Jaun-lac Noir N-S to NNE-SSW sinistral strike-slip zone is the dominant structural feature separating these two domains (Jeannet 1922; Plancherel 1976, 1979). A similar change in trend of the main structures of the Médiannes Plastiques can also be observed in the Chablais Préalpes.

Fault-propagation and/or detachment type folds seem to prevail, though synclinal and anticlinal breakthrough thrusts are common. Box-fold shaped structures are another frequently observed fold style, such as in the Dent de Broc anticline (Chassé 1924) or in the Rochers de Naye-Vanil Noir synclines (Wengen 1924; Revertera 1926). Folds and associated thrust planes die out laterally, resulting in the lateral transfer of displacement to other "en échelon" arranged fold-thrust structures (Bieri 1925; Nicol 1956; Mettraux & Mosar 1989; Mosar & Borel 1993). These transfer zones are locally associated with tear faults, which are kinematically linked with the fold-associated thrust planes (Mosar & Borel 1992). Displacements are left-lateral along N-S to NNE-SSW oriented tear faults and right-lateral along NW-SE to WNW-ESE oriented faults. Stress directions associated with thrusts and tear faults were calculated using fault/striae analyses and show a general NW-SE orientied compression direction. Fold interference structures, giving way to very tight folds, result from the simultaneous development of neighboring anticlines, as in the Col de Chaude area of the Tinière-Dorena anticline (Plancherel & Weidmann 1972).

Thrust surfaces are generally dipping towards the SE, as are fold axial planes. Only when closely spaced fault-related folds form imbricated stacks, are they overturned and dipping towards the NW such as in the Gantrisch area of the eastern Médiannes Plastiques (Plancherel 1979; Mosar 1991). This type of overturning mechanism is related, in the Médiannes Plastiques, to a thrust and fold development sequence progressively moving towards the N-NW foreland.

---

Fig. 8. Crustal cross-section of the frontal Alps along the N side of the Rhône valley (Fig.1 for location). The section has been constructed based upon existing surface (Badoux et al. 1960; Badoux 1965; Weidmann 1988; Badoux et al. 1990; Weidmann 1992; Mosar 1994) and subsurface data (Escher et al. 1993). (A) data from Sommaruga (pers. comm.) from the S Molasse, work in progress at Neuchâtel University, see also: Gorin et al. 1993; Marchant 1993; (B) seismic data and original interpretation are to be found in Steck et al. in press. The cross-section is not balanced at depth due to lack of information. Thrusts and imbricates beneath the Préalpes are to be considered as possible solutions that may help explain surface structures and a way in which the present day mountain front evolves. Thus the out of sequence thrusts that cut the whole Préalpes nappe pile are thought to be related to these thrusts at depth and the developing basement imbricates. (C) refers to "basement high" that may be explained in different ways: imbrication and duplexes in the sedimentary cover, inversion of a former permo-carboniferous trough or antiform developing above a newly developing basement imbricate whose basal thrust or rather shear zone (D') may connect to the seismic reflectors indicated by (D).

Overtured to recumbent large-scale folds are found in the Chablais Médiannes Plastiques. With its isoclinal geometry and its axial surface dipping at a shallow angle to the S, the Cornettes de Bise syncline is but one of the most spectacular ones (Chamot 1961; Badoux 1962a; Girod 1995).

Although described very early, backthrusts (top-to-the-S movement) appear to be a widespread structural feature (Schardt 1893a; Wengen 1924; Revertera 1926). They appear to develop in two distinct types of situations. Associated with ramps between two flats of a fault-bend fold, or together with breakthrough thrusts near the core of fault-propagation type anticlines thus forming pop-up structures in a crestal position. A second type is linked to anticlines showing important thickness changes between two adjacent limbs. Those changes are thought to be due to former synsedimentary normal faults, that are later inverted to form backthrusts. The most prominent of these backthrusts is in the Tsavas-Millelets anticline along the E slopes of the Sarine valley (Fig. 2a). The Dogger series change thickness from 2000 m on the W limb to 500 m on the E limb.

The general thrust direction is to the N-NW (Mosar 1988b, 1991) except for some thrusts found in the western termination of the Médiannes Plastiques near Montreux (Mosar 1994). This area forms a structural high above the Rhône valley and the Léman lake, with its structures steeply plunging towards the NE into the Sarine valley. From their geometry and their intersection of strata in the hangingwall it appears that some of the N-S trending thrusts indicate a movement towards the W (Fig. 2).

[ii] The Gastlosen-Tours d'Aï structure is located to the S of the transitional zone from the Médiannes Plastiques to the Médiannes Rigides sedimentation domain. In this domain the series of Malm age may rest directly on top of the Liassic Heiti formation or even the Triassic in the Tours d'Aï area (Fig. 4a).

To the N as well as to the S this structure is bordered by the Nappe Supérieure units (Chenevart 1945; Page 1969; Müller & Plancherel 1982). As between the E termination of the Tours d'Aï anticline (Jeannet 1912–1913) and the Laitemaire imbricate (Campana 1943), virtually all of the structure is “buried” underneath the Nappe Supérieure. Some small imbricates, that we propose to group with the Gastlosen-Tours d'Aï structure, can be seen in isolated outcrops N of the main structure in the Corbeyrier-Hongrin-Château d'Oex syncline. Based on stratigraphic criteria the Gastlosen-Tours d'Aï domain has been subdivided into two different zones (Baud 1972): a) the Gastlosen zone between the Laitemaire to the W and the Mittagflue near Boltigen to the E (Médiannes Rigides externes) and b) the Tours d'Aï – Grande Eau valley – Etivaz (Gorges du Pissot) area and the imbricates N and E of Boltigen along the Simme valley in the E Préalpes Médiannes (Médiannes Plastiques internes).

The tectonic imbricates or folds form above thrusts dipping to the S-SE. In the central Gastlosen-Laitemaire portion of the Préalpes Médiannes Romandes we observe a large imbricate sometimes with one or two small splays (Fig. 2). In the eastern part, in the Simmental area this structure is found as far east as Boltigen and is absent further east (Bieri 1946). It has been speculated that its equivalent might be hidden below the Médiannes Rigides south of the Simme valley (Mosar 1991).

In the western part in the Tours d'Aï and the Etivaz (Gorges du Pissot) area, the structure forms a large anticline with a vertical, thinned northern limb. To the S, the Tours d'Aï anticline connects to the Leysin syncline which is cored by the Nappe Supérieure. The Tours d'Aï anticline has a periclinal closure dipping by some 20° to-

wards the NE and disappears beneath the Nappe Supérieure S of lake Hongrin, only to be seen again in the Etivaz window. NE of here the fold “transforms” into several imbricates.

The subdivision of the Tours d’Aï-Gastlosen zone, based on sedimentological criteria is also reflected by the differences in structural geology: large fold (possibly fault-propagation type) in the Tours d’Aï-Etivaz portion and imbricate (probably fault-bend fold type) in the Laitemaire-Gastlosen portion. However, unlike the N-S juxtaposition proposed by paleogeographic criteria, we suggest here that the two zones form a lateral continuity with the original paleogeographical trends oblique to them.

[iii] The trailing part of the nappe, the Médiannes Rigides, is a N/NW-dipping thrust sheet, although locally there are one or two minor imbricate thrusts (Fig. 2a). The imbricates are gently north-dipping in the east (Spillgerten-Niederhorn area; Jaccard 1904; Rabowski 1912, 1920), while in the west (Rübli-Gummfluh) they dip more steeply (Jaccard 1907; Dubey 1962; Dousse 1965; Lonfat 1965). The major thrust sheets form fault-bend like folds that are cut by a large backthrust through the layers in the hangingwall of their ramps. Because the fault-bend fold structure has been tilted to the NW (its present day position), the backthrust and associated splays suffered a rotation and hence appear as normal faults with the southern compartment downthrown by 200–400 m (Jaccard 1904, 1907; Mosar 1991). The backthrusts cut not only through the Médiannes Rigides but also through the overlying Breccia nappe and Nappe Supérieure. Thus these two nappes have been emplaced on the Médiannes Rigides before the fault-bend fold and the associated backthrust developed (Mosar 1991). In the Mont d’Or and the Grande Eau, the Médiannes Rigides are reduced to slices consisting of series of Triassic age only (Fig. 1). Moreover, these slices (embedded in the Zone Submédiane) are overturned and dipping to the S-SE. A similar slice in the Rhône valley, the Saint Triphon klippe, is in an upside down position and completely detached from the rest of the Préalpes Médiannes. Because of its isolated position the connection of the Saint Triphon klippe with the Médiannes Rigides has remained speculative and several solutions, from a lateral or frontal portion of the inverted Médiannes Rigides folded imbricate (Weidmann et al. 1976; Escher et al. 1993), to a detached slab torn off during Late Tertiary to Quaternary times and emplaced independently of the structural evolution of the Préalpes Médiannes (Badoux 1962b) have been proposed.

The different Médiannes Rigides imbricates are mostly surrounded by terrains belonging to the Zone Submédiane, except where covered by the Breccia nappe. Between the Gummfluh-Rübli imbricates and, to the N and W, of the Mont d’Or inverted imbricate the Zone Submédiane (called locally “Petite fenêtre mitoyenne” as opposed to the “Grande fenêtre mitoyenne” for the major part of the Zone Submédiane, Lombard 1975; Weidmann et al. 1976) is in contact with the Nappe Supérieure (Fig. 1). Between the Grande Eau valley and the Hongrin lake this contact has a N-S direction (it juxtaposes Zone Submédiane to both the Nappe Supérieure in its N portion and to the Préalpes Médiannes in its S section) suggesting thrusting of the Zone Submédiane onto the Nappe Supérieure and the Préalpes Médiannes, probably in a westerly direction.

Especially in the Gummfluh-Rübli imbricate the large faults and thrusts are numerous and form a number of small slices. Next to inverse and normal displacements, strike-parallel movements, along more or less NE-SW oriented fault zones, appear to be more important than previously thought.

### *Internal deformation*

The internal deformation of limestones from the Préalpes Médiannes was measured using the distortion of pellets as well as twinning of sparitic calcite (Mosar 1989). The strain intensity in the frontal Médiannes Plastiques is very low and dominated by transgranular deformation mechanisms, mainly pressure solution, with weak development of calcite twins. The weak slaty cleavage is parallel to the fold axial surface (Mosar 1989, 1991). This deformation corresponds to an "early" shortening parallel to bedding emphasized by the development of tectonic stylolites and an early rock cleavage. Due to the shearing induced by the arrival of the Nappe Supérieure on top of the Médiannes Plastiques, small scale metric folds developed near the basal thrust in the Late Cretaceous marls and limestones (Mosar & Borel 1993). These folds are aligned on a great circle (on the stereonet projection) subparallel to the thrust surface and indicate a transport direction towards the NW.

The strain increases going into the more internal Médiannes Rigides where intragranular deformation mechanisms, such as twinning (numerous large, curved and twinned twins) and dynamic recrystallisation, prevail. The most intense strain is associated with large scale shearing due to thrusting on the basal décollements of the Préalpes Médiannes nappe and of the overlying Breccia nappe. The calculated finite extension is parallel to the general NW transport direction in the vicinity of the thrust planes. This is corroborated by the flat lying rock cleavage, mainly in the Late Cretaceous rocks at the top of the Médiannes Rigides (Mosar 1989, 1991). Elsewhere in the Médiannes Rigides nappe the extension direction is oriented at a high angle to the thrust planes, reflecting early shortening similar to that recognized in the Médiannes Plastiques.

In the Gummfluh klippe a mylonitic layer has been described at the base of the imbrication above the underlying Zone Submédiane and Niesen Nappe (Baud & Masson 1976; Cosca et al. 1992). Intense folding (cm to dm scale folds in Triassic limestones) has not been studied yet, but appears to be associated with intense thrust-induced shearing at the basal décollement of the Médiannes Rigides.

A strong cm to m thick cataclasis and locally a crenulation cleavage developed along some of the larger strike-slip faults as well as in the ramp-associated backthrust in the Médiannes Rigides or along thrust surfaces (Mosar 1988b; 1989; 1991; 1993).

Veins with calcite filling are ubiquitous, but only rarely are regular and well developed systems recognized, even more rarely two systems with cross-cutting relationship enabling relative chronological dating. One such example in the Dogger formation from the Médiannes Plastiques seems to have formed before and during folding (Mosar 1988b).

Anhydrites and gypsum from the Zone Submédiane have suffered strong deformation during the different tectonic events. Field observations and measurements from well log data have shown the existence of two directions of small scale fold axes: one N-S the second NE-SW (Zahner & Mosar 1993). They may be explained in terms of two superposed tectonic events; one associated with NW directed thrusting of the Préalpes Médiannes, the other possibly related to W directed movements most likely after the Préalpes were emplaced in their present day position. The actual timing of these events remains speculative due to the uncertainty of the structural relationship between the Préalpes Médiannes and the Zone Submédiane.

Because of their controversial origin and their ubiquity in the Préalpes, cornieule

(also called *cargneules* or *Rauhwaacke* (Weidmann 1971)) deserve a brief review. They are chemical breccias with honeycomb aspect due to preferred dissolution of dolomite clasts with respect to the calcite cement. Mainly three possible origins for this type of breccia have been proposed: i- a sedimentary Triassic or Tertiary origin (Schardt 1884; Brunnschwiler 1948), ii- a chemical genesis by dissolution and de-dolomitization during the Late Tertiary and the Quaternary (Brückner 1941; Jeanbourquin 1988), iii- a tectonic origin (Leine 1971; Masson 1972). The latter hypothesis proposes a mechanism of hydraulic fracturing in Triassic dolostones, which creates a high pressured fluid mush weakening and lubricating thrust surfaces (Masson 1972; Masson et al. 1980).

Recent studies in the Alps (Müller 1982, Jeanbourquin 1986, 1988; Amieux & Jeanbourquin 1989; Jeanbourquin & Lualdi 1994, Schaad 1995) clearly demonstrate that the *cornieules* are chemical or collapsed breccia formations evolving either directly or indirectly from evaporite sequences during Tertiary or Quaternary times. These transformations testify to high amounts of fluids circulating in these sheared Triassic sequences during the telogenesis.

The fact that most occurrences of *cornieule* are associated with or result from the transformation of Triassic dolostones is readily explained because major décollement levels are precisely located in evaporite layers (base of Middle and Upper Triassic in the Préalpes Médiannes). The deformed evaporites, and the accompanying fault gouge (breccias and cataclasites) resulting from the important heterogeneous deformation imposed by thrusting, make these sequences prone to “*cornieulisation*”. Thus, in the end several processes may have operated within the very same material. This renders their comprehension difficult.

### *Tectonics at the scale of the Alps*

In order to discuss the position and tectonics of the Préalpes in their larger present-day tectonic setting we constructed a cross-section extending from the autochthonous Molasse to the N, through the western termination of the Préalpes, to the external basement massifs of the Aiguilles Rouges (Fig. 8). This cross-section on a crustal scale is based on surface data along the profile as well as subsurface data from seismic investigations (petroleum prospection under the thrust Molasse and NFP20 survey below the Helvetic). No subsurface data are available below the Préalpes in this area. Classical cross-sections through this portion of the Alps (Escher et al. 1993) rely on projection of information into the section datum plane from distances up to 25 km. Except for the 10–12 km projection of the seismic data under the Helvetic from the Chablais Préalpes, the construction of Fig. 8 did not use long distance projections. The cross-section shows a rather flat sole of the Préalpes Médiannes close to sea-level (Mosar 1991; Vollmayr 1992). The folds and thrusts shown at depth beneath the Préalpes in the parautochthonous sedimentary cover (Jurassic and Molasse) are speculative and meant to highlight a kinematic process that can be related to structures observed in the Préalpes cover nappes and the thrust Molasse.

Pebbles in the Subalpine Molasse of the Mont Pélerin area find their origin in the Nappe Supérieure (Trümpy & Bersier 1954). These sediments of Chattian age are deposited on arrival of the Préalpes on the alpine foreland. Folding and thrusting of the Subalpine Molasse units thus can be considered post-Préalpes Médiannes and hence post-Nappe Supérieure emplacement.

Inside the Préalpes Médiannes several thrusts clearly hint at late post-emplacment movements. Thus the Tours d'Aï fault-propagation fold has developed a breakthrough thrust that brings it on top of the northerly adjacent isoclinal Corbeyrier syncline which is cored with Nappe Supérieure sediments. At the base of this syncline another late thrust can possibly be traced all the way along the Châteaux d'Oex syncline up to a position W of the Laitemaire imbricate. The transition from the Médiannes Rigides to the Médiannes Plastiques is mostly hidden under the Nappe Supérieure cover (Simme and Sarine valleys). A thrust juxtaposes the overturned Grande Eau Médiannes Rigides imbricate to the Leysin syncline of the Médiannes Plastiques in the Col des Mosses – Grande Eau area (Baud 1972). To the N of the overturned Mont d'Or slice the transition is by means of the Zone Submédiane, assimilated here to a broad and diffuse shear zone. The movement along these thrusts or shear zones (Fig. 8) is thought to be post-emplacment (post-Oligocene) and may possibly be linked at depth to the imbrications developing in association with basement thrusts such as the Infra Aiguilles-Rouges slice (Caron et al. 1989; Escher et al. 1993).

Along the base of the Médiannes Rigides, at the contact with the underlying Niesen Nappe, we always observe the Zone Submédiane. The presence of at least two fold axial directions in the evaporites possibly indicates a multiphase deformation history for this zone, one of which may be associated with the post-emplacment movements. NE-SW trending fold axes may have developed during the fold and thrust development and the emplacements of the Préalpes Médiannes, while a N-S trending set of folds may be linked to later west-directed movements. The N-S oriented contact of the Zone Submédiane directly on top of the Tours d'Aï periclinal closure (Zone Submédiane against Nappe Supérieure) may be linked to late thrusting thus further substantiating the existence of such west-directed movements. These W-directed movements, probably post-Oligocene in age, are not without recalling the large W and SW directed Oligocene to Quaternary movements in the Central and SW (French) Alps (Steck 1990; Mancktelow 1992; Seward & Mancktelow 1994; Steck & Hunziker 1994).

Further evidence for post-emplacment thrusting is found in the western termination of the Médiannes Plastiques E of Montreux (Fig. 1, 2b). Thrusts with a top to the W movement have been identified in this area. They are related to the structural high developing in the western termination of the Médiannes Plastiques Romandes. Indeed between this area and the central Sarine valley syncline the base of the Médiannes Plastiques plunges to the E-NE and drops from an altitude around +500 m to -1200 m (Fig. 2). This structural high appears to be related to a doming in the basement as indicated by seismic profiles (Fig. 8; Marchant 1993). This high is especially noticeable in the Chablais Préalpes Médiannes and we think is at the origin of the westward thrusting and the development of a structural high in the western termination of the Médiannes Plastiques Romandes. This structure is no longer visible further E than the Montreux area. The origin of the structural doming in the basement remains controversial. Thus an inverted Permo-Carboniferous trough has been invoked (Gorin et al. 1993). Alternatively it could represent the frontal part of a newly developing basement thrust whose detachment horizon could correspond to a shear zone possibly represented by a set of reflectors in the crust under the Infra Aiguilles-Rouges nappe (Escher et al. 1993; Steck et al. in press). Similar basement imbricates are described under the Alpine foreland SE of the Chablais Préalpes beneath the Annes and Sulens klippen (Guellec et al. 1990). A further possibility could imply the

development of a duplex structure in the sedimentary cover. Only further detailed seismic investigation may help resolve this problem.

Post-emplacment thrusting is also recognized in the neighbouring structural units (Fig. 1, 8):

- thus the Gurnigel nappe is found north of the Médiannes Plastiques which indicates that when the Nappe Supérieure was emplaced on top of the Médiannes Plastiques this transport went beyond the N edge of the Préalpes Médiannes nappe. The fact that the Gurnigel nappe is overturned in its S portion and overridden by the Médiannes Plastiques indicates thrusting after the Préalpes were emplaced in front of the Helvetic nappes.
- inside the Gurnigel nappe lenses of Molasse and Ultrahelvetic rocks are located along a thrust plane that brought them to the surface from underneath thus indicating, in our opinion, a late post-emplacment thrusting.
- a similar feature of a breaching thrust surface along with Molasse lenses transported from underneath the Préalpes nappe stack, is found at the tectonic contact between Gurnigel nappe and the Ultrahelvetic unit at the front of the Préalpes, thus indicating that this tectonic contact is a post-emplacment thrust.
- the outermost N Préalpes are in contact with the Subalpine Molasse by means of a belt of “wildflysch” and Lower Marine Molasse. The northern thrust bounding this zone cuts the existing thrusts in the Subalpine Molasse indicating yet another post-emplacment thrusting.

The orientation of the three latter thrust surfaces is NNE-SSW to N-S hinting to a possible westward movement similarly to the late movements and thrusts described in the Préalpes Médiannes.

Evidence for late thrusting postdating the development of folds and thrusts in the Molasse substratum, is found in the E Chablais area. Here Molasse bedding and structures do not have the same orientation as those in the overlying Médiannes Plastiques by which they appear to be cut.

Work in progress in the Chablais Préalpes Médiannes shows the importance of large strike-slip faults cutting the existing fold and thrust structures. This system of conjugate faults trending N-S and WNW-ESE thus postdates the main folding and thrusting event and corresponds to a late post-emplacment event.

### *Structural development and chronology*

The alpine “deformation” in the Préalpes Médiannes starts with the final closure of the Valais and the Piemont oceans. Relevant information about the timing of this event come from the depositional age of the youngest flysch sediments of the different Préalpes nappes (Caron et al. 1980a, 1989; Winkler 1984; Wildi 1985; Homewood & Lateltin 1988). Indeed, these deep marine sediments of detrital and orogenic character, are directly related to the mountain building process of the Préalpes and Alps (Studer 1827; Schardt 1893b; Tercier 1948; Homewood & Lateltin 1988; Caron et al. 1989). The Préalpes Médiannes flyschs range in age from Middle to Late Eocene (Lutetian-Bartonian). In the Nappe Supérieure the oldest flyschs are of Turonian to Maastrichtian age, while the youngest ones, found in the Gurnigel nappe, are Maastrichtian to Bartonian in age. In



the Niesen nappe the oldest flyschs are Maastrichtian in age and the youngest ones Lutetian. The most recent orogenic deposits of the Préalpes are the Ultrahelvetic flyschs with their Lutetian to Bartonian age.

We thus suggest a tectonic convergence that first reaches the Nappe Supérieure sedimentation domain (with the Gurnigel nappe being involved later than the rest of the Nappe Supérieure units). The Préalpes Médiannes are second, while the Niesen nappe and finally the Ultrahelvetic units register the youngest sedimentary development of the Préalpes.

In the Médiannes Rigides, the Nappe Supérieure is emplaced on top of the Breccia nappe which itself rests on top of the Médiannes Rigides. Because a large backthrust associated with the Médiannes Rigides ramp fold formation cuts these three units, their emplacement has to be prior to the fold and thrust development in the trailing Préalpes Médiannes. Since the Nappe Supérieure is folded with the Médiannes Plastiques, its emplacement had to predate the fold and thrust development in this portion of the Préalpes Médiannes. The arrival of the Nappe Supérieure and the Breccia nappe on top of the Médiannes Plastiques caused strong shearing at the contact thus developing a strong deformation with a finite extension parallel to the transport direction. The very northern portions of the Nappe Supérieure (mainly the Gurnigel nappe) have traveled further than the present day northern limit of the Préalpes Médiannes, since we find them resting on top of Ultrahelvetic beyond the northern and northwestern edge of the Médiannes Plastiques.

We thus can propose an emplacement sequence as follows: Nappe Supérieure on Breccia nappe and Médiannes Rigides (possibly also on Médiannes Plastiques); Breccia nappe (carrying the N. Supérieure) on Médiannes Rigides and Nappe Supérieure on Médiannes Plastiques.

It is only after this emplacement during Bartonian times that the Préalpes Médiannes detach from their homeland and undergo thin-skinned tectonics in the shallower part of the accretionary prism active at the front of the growing alpine orogeny. The undetached homeland cover, such as the Siviez-Mischabel or the Pontis nappes sedimentary series and the basement are incorporated in the deeper part of the forming mountain belt and suffer high-grade metamorphism and strong deformation. The basement is subducted and underplated to form the eclogitic root zone of the orogen (Marchant & Stampfli in press).

The emplacement on top of the Ultrahelvetic units, the Helvetic nappes and the Niesen nappe development (Steck 1984) are so far poorly understood. Investigation in the Ultrahelvetic mélanges (Jeanbourquin et al. 1992; Jeanbourquin 1994), however, reveal evidence for "early" (Tertiary) deformation developed in the accretionary prism below the overriding Pennine nappes.

In the Préalpes Médiannes layer-parallel shortening indicates a general compression and deformation just prior to folding and thrusting. The thrust-related fold development appears to be foreland migrating (Mosar 1991).

Imbrication and backthrusting in the Médiannes Rigides develops before the intermediate Gastlosen-Tours d'Aï zone forms. It is in this latter zone that the basal décollement steps up from the basal Middle Triassic to the base of the Late Triassic. In the Médiannes Plastiques overturned folds and fold axial surfaces with a S-SE dip are found south of normal and upright folds. This is another indication of a foreland (N-NW) propagating

sequence of fold and thrust development. Internal deformation is associated to the general fold and thrust development (Mosar 1989).

The progressive steepening and overturning of the Médiannes Rigides from E to W is linked to the differences in development of the underlying Helvetic nappes. Thus in the W part, the present day position of the Médiannes Rigides imbricates has been interpreted as belonging to the frontal and overturned limb of a large scale fold involving the Helvetic, Ultrahelvetic, Niesen and Médiannes Rigides nappes. This fold refolds intra-Préalpes thrusts and develops above a thrust surface rooted in the basement (Caron et al. 1989; Escher et al. 1993). The structure is thought to develop upon or after the arrival of the Préalpes in their present position. It will later develop a breakthrough juxtaposing the Médiannes Rigides of the Grande Eau area with the S Médiannes Plastiques.

The final emplacement in front of the Helvetic nappes and the external basement massifs is associated with the formation of several thrusts cutting the Préalpes Médiannes (see prior discussion). We propose that these thrusts root in the present day Préalpes substratum, that is the Mesozoic and Tertiary cover where they may be linked to thrust surfaces going all the way down to the crystalline basement (see also: Caron et al. 1989; Escher et al. 1993). Structural features such as large conjugate strike-slip systems (f.i. in the Chablais Préalpes Médiannes) are records of this late tectonic event. It is also during this period that we believe the W-directed movements occur.

### **Metamorphism s.l.**

Numerous studies (see references in Kübler et al. 1979; Mosar 1988a; Zahner & Mosar 1993) showed that the Préalpes underwent low-grade metamorphism ranging from diagenetic to epizonal (300–400 °C) conditions. These studies, covering most of the Préalpes units with a varying density, applied a variety of techniques such as illite crystallinity, the evolution of the clay mineral parageneses (Mosar 1988a; Jaboyedoff & Thélin in press), maturation of different types of organic material (Kübler et al. 1979; Burkhard & Kalkreuth 1989; Borel 1991), fluid inclusions in quartz crystals (Stalder & Touray 1970; Mullis 1979) and oxygen isotopes (Cosca et al. 1992).

This metamorphism varies from diagenesis in the north (i.e. in the Médiannes Plastiques) to epizone in the south (i.e. in the Médiannes Rigides). This zonation also exists from top to bottom in the southern part of the Préalpes nappe stack (Masson et al. 1980; Baud 1984; Mosar 1988a). The uppermost Nappe Supérieure and the upper part of the Breccia nappe are diagenetic, and metamorphic grade increases towards the lower part of the Breccia nappe. In contrast the diagenetic stage is preserved in the upper part of the underlying Niesen nappe (Dortmann 1982; Mosar 1988a). Thus metamorphism in the Préalpes Médiannes, Breccia nappe and Nappe Supérieure is transported. Discontinuous metamorphic gradients also occur between the Niesen nappe and the underlying Ultrahelvetic units, and between the Ultrahelvetic units and the underlying Helvetic nappes (Burkhard 1988; Burkhard & Kalkreuth 1989; Goy-Eggenberger & Kübler 1990) as well as amongst different units of the Ultrahelvetic.

Since the frontal parts of the Préalpes Médiannes have not evolved beyond diagenesis, the very low-grade metamorphism of this portion of the Préalpes Médiannes begins with the deposition and compaction of the sediments. The main metamorphism, especially in the trailing Préalpes Médiannes, the Breccia nappe and the Nappe Supérieure, however,

occurred during thrusting of the overlying nappes and subsequent burial since Late Eocene (Bartonian) times. It would require the Médiannes Rigides to be buried to a depth of 10 km during some stage of the Alpine orogeny to explain the epizonal temperatures above 300 °C (gradient of ~30 °C/km, Mosar 1988a; Jaboyedoff & Thélin in press).

Because this burial occurred in the geodynamic context of an accretionary prism a lower geothermal gradient may be expected for the Préalpes, at least during the beginning of their involvement in the accretionary prism (Fig. 7). Thus the Préalpes Médiannes might find their way as cold material along the subduction zone and reach greater depths than normal.

The burial metamorphism of the Préalpes Médiannes by the superposition of the Nappe Supérieure and the Breccia nappe is corroborated by syn-metamorphic internal deformation mechanisms associated with the arrival of the latter nappes (Mosar 1989, 1991).

The peak metamorphism is attained before the arrival of the Préalpes Médiannes onto the Niesen, Ultrahelvetic and Helvetic domains, as shown by the discontinuous metamorphic gradients between these units.

Recent studies have shown two events that appear to be related to hydrothermal fluid circulation, but whose interpretation is still open to discussion. The first study shows an unusually high concentration of barium, present under the form of barite scattered through the rock, in Cretaceous sediments of the Préalpes Médiannes (Meisser, Musée géologique, Lausanne, work in progress). This type of mineralisation is thought to have a geothermal, syn-depositional origin. A cupriferous mineralisation associated with fossils in Late Cretaceous rocks is thought to develop in hydrothermal environments in the 100 °C range during sedimentation (Meisser 1992). Other mineralisations in veins (Meisser 1992) and in country rock seem to be associated with low temperature hydrothermal circulation possibly during the alpine orogenesis (Zingg 1984; Oberhänsli et al. 1985).

The second study also includes Rb/Sr, K/Ar and  $^{40/39}\text{Ar}$  dating of white micas from Triassic and Middle Jurassic limestones that yield ages of ~60 to 80 Ma in the trailing part of the Médiannes Rigides in the western Préalpes Médiannes (Masson et al. 1980; the Gummfluh imbricate: Huon et al. 1988; Cosca et al. 1992; the Amselgrat imbricate: De Coulon 1990). The “mylonitic rocks” described at the base of this imbrication, in the vicinity of the basal thrust (Baud & Masson 1976; Cosca et al. 1992), are thought to be heated up to 280 °C, as shown by oxygen isotope measurements. The ages are interpreted by the authors as strong evidence for a Cretaceous tectonic and/or thermal event that was probably related to the initial stages of closure of the Alpine Tethys. The synchronous heating of the basal layer is thus explained by the authors of that study, by hydrothermal fluid circulation during thrusting towards the S-SE in a transpressive continental collision environment in Late Cretaceous/Paleocene times. The conspicuous coincidence of these events with the sedimentation gaps during the Paleocene and Late Cretaceous as well as the rather continuous sedimentation during the Late Maastrichtian (Forclettes Formation) has still to be explained. The temperature range given by oxygen isotopes is similar to the one from the Eocene anchizonal to epizonal burial metamorphism (T ~300–350 °C) and would thus explain why the older temperatures have not been reset by the younger burial metamorphism.

## Discussion

The chronology and structural interpretation of the Préalpes Médiannes given here, though based on firmly established observational data, have been the subject to several controversies:

[i] Synsedimentary tectonics have so far always been linked to faults rooted in the underlying basement. They thus reflect active basement tectonics; either normal faults, inversions along former normal faults or transpressional movements resulting in flower structures. Most authors tried to explain synsedimentary structures in terms of local or regional structural highs and lows (basins) related to extensional or compressional tectonic regimes. Alternatively we propose that the whole of the Préalpes Médiannes sedimentation realm develops above an evaporite décollement and that the structures in the hanging-wall are the result of large-scale sliding outward with respect to an uplifting rift-shoulder. The developing structures are the result of the interplay of faults (normal and inverted) in an overall extensional passive margin tectonic regime.

[ii] Since the Bartonian the development of the very low-grade metamorphism is attributed mainly to burial by overriding tectonic nappes during the involvement of the Préalpes Médiannes in the accretionary prism after the closure of the Piemont ocean (Fig. 7). Radiometric ages around 80–60 Ma and associated isotopic temperatures around 280 °C have so far been explained by hot fluid circulation along thrust planes active during the Late Cretaceous/Paleocene. We do not think however that these observations are to be linked to a Cretaceous orogeny in the Préalpes. We rather propose to integrate these data, together with the information on hydrothermalism, in the geodynamic context of an accretionary prism s.l. (Fig. 6, 7). The Briançonnais micro-continent underwent two major tectonic events: first it is breaking off the Iberic plate in Late Cretaceous and subsequently evolves as an exotic terrain detached from its lithospheric substratum; secondly it is integrated into the accretionary prism between the convergent Adriatic and European plates during Late Eocene (Fig. 6, 7). We propose that the 80–60Ma year radiometric ages and the associated hydrothermal fluid circulation are linked to the “inclusion” of the Briançonnais exotic terrain between the accretionary prism of the closing Valais ocean to the N and the closing Piemont ocean to the S. This thermal event might also be linked to the delamination of the Briançonnais from its lithospheric substratum and the subsequent heat influx. Further investigation especially on the microtectonics of the thrust zone at the base of the Médiannes Rigides will help to obtain more detailed information on the geodynamic context.

[iii] The dynamics of the Préalpes emplacement have long thought to be dominated by gravity gliding (Schardt 1898; Lugeon 1943; Campana 1963; Debelmas & Kerckhove 1973). Thus models like the “diverticulation” (Lugeon 1943) have been proposed in order to explain the superposition and imbrication of the different tectonic units such as the Ultrahelvetic units. But since the 1970's it appeared that gravity gliding could not be the sole mechanism active in the Préalpes (Lemoine 1973). It appears now that the Préalpes evolved in a classic mountain building environment of plate collision (Mosar 1991; Stampfli 1993) such as described for example by tapered or non-tapered wedge models (Davis et al. 1983; Willett et al. 1993; Beaumont et al. 1994; Beaumont & Quinlan 1994).

[iv] Two major end-member models explaining the fold and thrust development in the Préalpes Médiannes have been proposed to date: in a first model, folds and thrusts

may result from post- to syn-emplacement tectonics after the Préalpes Médiannes were transported over the Helvetic nappes (Plancherel 1979); in a second model their development may be explained by “early” tectonics before their arrival on top of the Helvetic nappes (Mosar 1991). In the first model alpine structures are induced by strike-slip faults rooted in the autochthonous basement. These faults would affect the whole tectonic nappe pile of the present day klippen. In the second possibility the main deformation is considered to develop before, together or shortly after the detachment of the Préalpes Médiannes from their homeland. Thus beginning with their involvement in the accretionary prism of the closing Piemont ocean, the Préalpes Médiannes, the Breccia nappe and the Nappe Supérieure are superposed by thrusting. The subsequent metamorphism is associated with the main fold and thrust development. It is the impingement of the Adriatic plate on the Briançonnais and European plate as well as the isostatic rebound (flexural effect) following the detachment of the deeper portions of the subducted Valaisan and Piemont crust that caused the necessary change in mountain belt dynamics. This in turn alters the critically tapered wedge shape of the accretionary prism. In order to recover a stable configuration, shallow depth thrusting in the frontal part of the dynamic wedge (including the Préalpes thrust sheets) would explain why the Préalpes were not underplated (Marchant & Stampfli in press).

We strongly favour this latter interpretation. The transport on top of the Helvetic nappes was not a passive one (meaning without reactivation of former thrust and development of new thrusts and folds), neither is the emplacement in front of those nappes and the external massifs. Several tectonic events and structures can be attributed to those periods with the passage on top of the Helvetics remaining the more uncertain.

Topographic and geomorphologic criteria may help prove very recent basement uplift due to the development of a new basement imbricate underneath the Préalpes (Fig. 8). Thus there is a step in mean elevation between the autochthonous Molasse and the thrust Molasse/Préalpes area. This “plateau” is located, in our opinion, above the potential basement imbricate developing at depth. Detailed analyses of topography and of geomorphologic criteria such as valley incision and pass height, may help us understand this process and the timing of the uplift (tectonically induced).

[v] Strike-slip faults s.l. (possibly tear faults) are an important structural feature in the Préalpes and especially in the Préalpes Médiannes (Plancherel 1976, 1979). Typically, the western edges of the Chablais and Romandes Préalpes Médiannes are formed by such fault zones; but also within the Préalpes Médiannes important vertical fault zones may develop such as the Jaun-Lac noir zone (see chap. “Structural geology”). Their importance and role has been variably appreciated and incorporated into the tectonic evolution of the Préalpes Médiannes. Plancherel (1979) links the motion of N/NNE-S/SSE oriented, left-lateral strike-slip faults to the fold and thrust development of the Préalpes Médiannes. These faults would root in the basement and were active after the emplacement of the Préalpes in front of the Helvetic nappes. They are also involved in the deformation of the underlying subalpine and autochthonous Molasse (see also: Pavoni 1987; Sambeth & Pavoni 1988).

Similar ideas of large scale strike-slip deformation have been applied to the Briançonnais and Sub-briançonnais domains of the western French Alps (Maury & Ricou 1983; Ricou 1984; Ricou & Siddans 1986). For these authors those strike-slip faults are, however, not only active in Oligocene to present times, but already existed in the

Briançonnais s.l. paleogeographic realm (active since Late Cretaceous) where they played an important role in structuring the sedimentation domains (see also discussion herein and: Sartori 1990).

Investigations in the Préalpes Médiannes Romandes also show that some of the major strike-slip faults are in fact tear faults (that do not root in the present day substratum) associated with the fold and thrust development (Mosar & Borel 1992; Mosar 1994). It thus appears that the strike-slip faults s.l. of the Préalpes Médiannes probably originated in the paleogeographic realm (true strike-slip faults rooted in the basement). These structures have subsequently been reactivated as tear faults during the main fold and thrust development after the Bartonian. No conclusive evidence, however, shows that they are still active in the NW-SE oriented compressional stress regime of the alpine foreland (Deichmann 1990).

[vi] Although considered to be derived from one paleogeographic sedimentation domain, the connection between the Médiannes Plastiques – Médiannes Rigides is nowhere clear in the Préalpes. Especially the connection between the Médiannes Rigides and the intermediate Tours d'Aï -Gastlosen structure is always buried under the overlying Nappe Supérieure. This led to the hypothesis that they might have been separated already during Cretaceous times. The continuous sedimentation and the lack of tectonic markers (for a potential tectonic separation during the Cretaceous), however, leads us to discard this possibility. The present day discontinuity between the two tectonic domains results from their involvement in folding of the Helvetic nappes and thrusting after their emplacement.

## Conclusions

From data and discussions presented, keeping in mind that the structural evolution appears to be continuous and progressive, we propose to differentiate four main stages in the tectonic history of the Préalpes Médiannes, each of which saw the development of different and specific structural features.

[i] A passive margin period (mainly Jurassic) during which we see the formation of normal faults, subsidence-induced basins above extensional faults associated with a basal décollement in evaporites. NE-SW trending basins are separated by structural highs resulting from local tectonics. Inversion structures are sealed by synsedimentary growth features. These structures formed in a rim basin bordered to the S by a rift shoulder issued from the Piemont oceanisation. The different types of structures observed during the sedimentation history can be related to thermal events linked to the development of the different Alpine oceanic domains, mainly the opening of the Piemont ocean but also the development of the Valais domain.

[ii] An active margin period (starting in Middle to Late Eocene: Lutetian-Bartonian) during which the Briançonnais flysch sedimentation marks the location of the Préalpes Médiannes at the front of the Alpine orogeny. Finally, during Bartonian to Priabonian the now exotic terrain is incorporated into the accretionary prism of the closing Piemont ocean.

Continental collision during Late Eocene to present sees the development of the main structural features of the Préalpes Médiannes fold-and-thrust belt. The Breccia nappe and the Nappe Supérieure are thrust on top of the Médiannes Rigides and Plas-

tiques and the metamorphism develops by tectonic burial. Thrust movements are recorded by internal deformation along the nappe contacts. It is after this emplacement that the Préalpes Médiannes are detached from their substratum to undergo thin skinned tectonics in the brittle domain. Internal rock deformation registers pre- and synfolding and -thrusting deformation. The fault-related folds (fault-propagation folds, fault-bend folds), box-shaped folds, imbricates, imbricated fans and en échelon structures associated with tear faults develop in sequence, progressively migrating towards the foreland. Backthrusts and breakthrough thrusts are frequent.

[iii] The arrival of the Préalpes on the European continent, that is on top of the Ultrahelvetic units during final Eocene – beginning Oligocene and on the Helvetic units during Oligocene (probably Rupelian) times. By then the Préalpes Médiannes have achieved most of their deformation and structural development. The transport on top of the deforming Ultrahelvetic and Helvetic nappes caused renewed large-scale deformation in the Préalpes Médiannes nappe. Thus the Médiannes Rigides, at least their western portion, and the basal décollement are involved in the folding of the frontal portion of those nappes, which confers them their present day steeply inclined or overturned position.

The arrival of the Préalpes on the Molasse basin (in subaerial conditions?) is recorded by the deposition of the conglomerates of the Mont Pélerin of Lower Chattian age. These sediments largely result from the erosion of the Nappe Supérieure.

[iv] A final period of thrusting, probably Chattian to younger in age, occurs in the Préalpes at the front of the still active mountain belt. Thus out of sequence thrusting occurs inside the Préalpes Médiannes (such as between the Médiannes Rigides and the Médiannes Plastiques). These thrusts are associated with the development of the external crystalline massifs of the Alps. Similar thrusts are also active in the Subalpine Molasse and we propose they are also linked to the development of imbrications in the basement and in the allochthonous sedimentary substratum. Westward directed thrusting occurs in the Préalpes Médiannes (frontal imbricate, Zone Submédiane) and is linked with a basement high developing underneath the western termination of the Préalpes Romandes and extending into the Chablais Préalpes. It may not be excluded that these movements towards the W are related to the large scale W to SW directed displacements in the central and southwestern Alps. Since these movements are of Oligocene to Quaternary age, hence also contemporaneous to the Jura folding, the latest tectonic “activity” in the Préalpes may be of the same age.

Unlike most of the Pennine units the Préalpes Médiannes thus succeeded in escaping an untimely reshaping (overheating and awful deformation) in the accretionary prism due to changes in the plate tectonic regime and the subsequent changes in thrust dynamics. Since then they enjoyed the unbearable lightness of thin-skinned tectonics and have been surfing the frontal “wave” of the alpine orogeny.

### Acknowledgments

We are thankful to A. Baud, G. Borel, M. Cosca, P. Jeanbourquin, R. Marchant, N. Meisser and M. Septfontaine for their many discussions and suggestions. The “Backstripp89” software program for subsidence curves was kindly provided by R. Schegg at University of Geneva. We appreciate the help of A. Sommaruga of Neuchâtel University who provided the seismic data and depth conversions for the S Molasse and Flysch area. A. Escher and A. Steck are kindly thanked for providing the seismic lines and interpretation from the Chablais area. We also would like to thank T. Baudin, H. Funk, S. Schmid and R. Trümpy for reviewing and helping improve this paper as for helpful suggestions. This work has partially been financed by the Swiss FNSRS (requests n° 21-29011.90, 20-33450. 92).

## REFERENCES

- ACKERMANN, A. 1986: Le flysch de la nappe du Niesen. *Eclogae geol. Helv.* 79, 641–684.
- AMIEUX, P. & JEANBOURQUIN, P. 1989: Cathodoluminescence et origine diagénétique tardive des cargneules du Massif des Aiguilles Rouges (Valais, Suisse). *Bull. Soc. géol. France* 8, 123–132.
- BADOUX, H. 1962a: Géologie des Préalpes valaisannes. *Matér. Carte géol. Suisse n.s.* 113, 1–86.
- 1962b: La géologie des collines de Saint-Triphon. *Bull. Soc. vaud. Sci. nat.* 68, 35–48.
- 1963: Les unités ultrahelvétiques de la Zone des Cols. *Eclogae geol. Helv.* 56, 1–13.
- 1965: Montreux, Feuille n°47, notice. *Comm. géol. Suisse, Basel.*
- BADOUX, H., CHESSEX, R., JEANNET, A., LUGEON, M. & RIVIER, F. 1960: Monthey, Feuille n°37, notice. *Comm. géol. Suisse, Basel.*
- BADOUX, H., GABUS, J.H. & MERCANTON, C.H. 1990: Les Diablerets, Feuille n°88, notice. *Serv. hydrol. géol. national.*
- BADOUX, H. & MERCANTON, C.H. 1962: Essai sur l'évolution tectonique des Préalpes médianes du Chablais. *Eclogae geol. Helv.* 55, 135–188.
- BAILEY, E.B. 1935: *Tectonic essays, mainly Alpine.* Clarendon Press, Oxford.
- BALLY, A.W. 1981: Thoughts on tectonics of folded belts. In: *Thrust and nappe tectonics* (Ed. by McClay, K.). *Geol. Soc. London*, 13–32.
- BAUD, A. 1972: Observations et hypothèses sur la géologie de la partie radicale des Préalpes médianes. *Eclogae geol. Helv.* 65, 43–55.
- 1984: Histoire naturelle des calcaires de Saint-Triphon (Anisien, Préalpes et Alpes occidentales). PhD thesis, Univ. Lausanne.
- BAUD, A., HEINZ, R. & SEPTFONTAINE, M. 1989: Compte rendu de l'excursion de la Société Géologique Suisse dans les Préalpes, du 2 au 4 octobre 1988. *Eclogae geol. Helv.* 82, 359–377.
- BAUD, A. & MASSON, H. 1975: Preuves d'une tectonique de distension dans le domaine briançonnais: failles conjuguées et paléokarst à Saint-Triphon (Préalpes médianes, Suisse). *Eclogae geol. Helv.* 68, 131–145.
- 1976: Déformation ductile et bréchification le long du plan de chevauchement de l'écaïlle de la Gummfluh (Préalpes médianes rigides, Suisse)-Résumé. *Eclogae geol. Helv.* 69, 471–472.
- BAUD, A. & SEPTFONTAINE, M. 1980: Présentation d'un profil palinspastique de la nappe des Préalpes médianes en Suisse occidentale. *Eclogae geol. Helv.* 73, 651–660.
- BEAUMONT, C., FULLSACK, P. & HAMILTON, J. 1994: Styles of crustal deformation in compressional orogens caused by subduction of the underlying lithosphere. *Tectonophysics* 232, 119–132.
- BEAUMONT, C. & QUINLAN, G. 1994: A geodynamic framework for interpreting crustal-scale seismic-reflectivity patterns in compressional orogens. *Geophys. J. Int.* 116, 754–783.
- BERNOULLI, D., CARON, C., HOMEWOOD, P., KALIN, O. & STUIJVENBERG, v.J. 1979: Evolution of continental margins of the Alps. *Schweiz. mineral. petrogr. Mitt.* 59, 165–170.
- BEUKEL, J.V.D. 1990: Breakup of young oceanic lithosphere in the upper part of a subduction zone: implication for the emplacement of ophiolites. *Tectonics* 9, 825–844.
- BIERI, P. 1925: *Der Bau der Klippendecke zwischen Gantrisch und Simmental (Berner Oberland).* *Iber. phil. Fak. II, Univ. Bern* 5, 89–109.
- 1946: Über die Ausbreitung der Simmendecke in den östlichen Préalpes médianes. *Eclogae geol. Helv.* 39, 25–34.
- BOILLOT, G., MONTADERT, L., LEMOINE, M. & BIJU-DUVAL, B. 1984: Les marges continentales actuelles et fossiles autour de la France. Masson, Paris.
- BOLLER, K. 1963: Stratigraphische und Mikropaläontologische Untersuchungen im Neocom der Klippendecke (östlich der Rhône). *Eclogae geol. Helv.* 56, 15–102.
- BOREL, G. 1991: *Etudes géologiques et minéralogiques de la région du Widdersgrind (Préalpes romandes).* Licence, Neuchâtel.
- BRÜCKNER, W. 1941: Über die Entstehung der Rauwacken und Zellendolomite. *Eclogae geol. Helv.* 34, 117–134.
- BRUNNSCHWILLER, R.O. 1948: Beiträge zur Kenntnis der Helvetischen Trias östlich des Klausenpass. PhD thesis, Univ. Zürich.
- BURKHARD, M. 1988: L'Helvétique de la bordure occidentale du Massif de l'Aar (évolution tectonique et métamorphique). *Eclogae geol. Helv.* 81, 63–114.
- BURKHARD, M. & KALKREUTH, W. 1989: Coalification in the northern Wildhorn nappe and adjacent units, western Switzerland. Implications for tectonic burial histories. *Int. J. Coal Geol.* 11, 47–64.



- CAMPANA, B. 1943: Géologie des nappes préalpines au NE de Château-d'Oex. Matér. Carte géol. Suisse n.s. 82, 1–64.
- 1963: The emplacement of the Prealpine and Helvetic nappes as submarine gliding of plastic masses. In: Syntaphral tectonics and diagenesis (Symp. ed. by ELLISTON, J. & CAREY, S.W.). Hobart, University of Tasmania. E, E1–E6.
- CARON, C. 1972: La nappe supérieure des Préalpes: subdivisions et principaux caractères du sommet de l'édifice préalpin. *Eclogae geol. Helv.* 65, 47–73.
- 1973: Survol géologique des Alpes occidentales. *Bull. Soc. fribourg. Sci. nat.* 62, 73–81.
- CARON, C., HOMEWOOD, P. & STUIJVENBERG, J. 1980a: Les flyschs Préalpins. In: *Sédimentation détritique, documents complémentaires*, 1–23.
- 1980b: Flysch and Molasse of Western and Central Switzerland. In: *Geology of Switzerland, a guide-book, Part B, Geological Excursions* (Ed. by Trümpy, R.). Wepf and Co., Basel, 274–278.
- CARON, C., HOMEWOOD, P. & WILDI, W. 1989: The Original Swiss flysch: A Reappraisal of the Type Deposits in the Swiss Prealps. *Earth-Sci. Rev.* 26, 1–45.
- CARON, M. 1976: Révision des foraminifères planctoniques décrits dans la région du Montsalvens (Préalpes fribourgeoises). *Eclogae geol. Helv.* 69, 327–333.
- CARON, M. & DUPASQUIER, C. 1989: Litho- et biostratigraphie des dépôts du "Crétacé moyen" dans les Préalpes Médiannes. *Geobios, mém. spéc.* 11, 49–58.
- CHAMOT, G. 1961: La Nappe des Préalpes médianes entre les Cornettes de Bise et le col du Corbier (Chablais français). *Bull. Lab. géol. minéral. geophys. Univ. Lausanne* 131, 1–86.
- CHASSÉ, V.T. 1924: Beitrag zur Geologie der Dent de Broc und ihrer Umgebung. PhD thesis, Univ. Zürich.
- CHAULIEU, P. 1992: La sédimentation détritique sur la marge nord téthysienne. Témoin de l'évolution géodynamique des Alpes Occidentales. PhD thesis, Univ. Grenoble.
- CHENEVART, C. 1945: Les nappes des Préalpes Médiannes et de la Simme dans la région de la Hochmatt. *Mém. Soc. fribourg. Sci. nat.* 12, 1–213.
- COSCA, M.A., HUNZIKER, J.C., HUON, S. & MASSON, H. 1992: Radiometric age constraints on mineral growth, metamorphism, and tectonism of the Gummfluh klippe, Briançonnais domain of the Préalpes, Switzerland. *Contr. Mineral. Petrol.* 112, 439–449.
- DAVIS, D.M., SUPPE, J. & DAHLEN, F.A. 1983: Mechanics of fold and thrust belts and accretionary wedges. *J. geophys. Res.* B 88, 1153–1172.
- DE COULON, S. 1990: Géologie et minéralogie de l'écaille de l'Amselgrat. Diplôme, Lausanne.
- DEBELMAS, J. 1989: On some key features of the evolution of the Western Alps. In: *Tectonic evolution of the Tethyan region* (Ed. by SENGÖR, A.M.C.). Kluwer Academic Press, 23–42.
- DEBELMAS, J. & KERCKHOVE, C. 1973: Large gravity nappes in the French-Italian and French-Swiss Alps. In: *Gravity and Tectonics* (Ed. by DE JONG, K.A. & SCHOLTEN, R.). J. Wiley and Sons, 189–200.
- DEICHMANN, N. 1990: Seismizität der Nordschweiz, 1980–1988, und Auswertung der Erdbebenserien von Günsberg, Läuelfingen und Zeglingen. *Cédra, Rapport technique* 90–46, 52.
- DORTMANN, T. 1982: Ein Querprofil durch die Niesendecke. Sedimente, Tektonik, Metamorphose und ihre Bedeutung für die Gebirgsbildung. Diplom, Bonn.
- DOUSSE, B. 1965: Géologie des Rochers de Château d'Oex. Matér. Carte géol. Suisse, n.s. 119, 1–77.
- DUBEY, R. 1962: Géologie de la partie occidentale des Rochers de Château d'Oex (Préalpes romandes). PhD thesis, Univ. Fribourg.
- ELLENBERGER, F. 1952: Sur l'extension des faciès Briançonnais en Suisse, dans les Préalpes médianes et les Pennides. *Eclogae geol. Helv.* 45, 285–286.
- ESCHER, A., MASSON, H. & STECK, A. 1993: Nappe geometry in the Western Swiss Alps. *J. struct. Geol.* 15, 501–509.
- FAVRE, E. & SCHARDT, H. 1887: Description géologique des Préalpes du canton de Vaud et du Chablais jusqu'à la Dranse et de la chaîne des Dents du Midi formant la partie NW de la feuille XVII. Matér. Carte géol. Suisse 22, 1–636.
- FAVRE, P. & STAMPFLI, G. 1992: From rifting to passive margin: the Red Sea, the central Atlantic and the Alpine Tethys as examples. *Tectonophysics* 215, 69–97.
- FLÜCK, W. 1973: Die Flysche der prealpinen Decken im Simmental und Saanenland. Matér. Carte géol. Suisse n.s. 146, 1–88.
- FRISCH, W. 1979: Tectonic progradation and plate tectonic evolution of the Alps. *Tectonophysics* 60, 121–139.
- FUIS, G.S. & CLOWES, R.M. 1993: Comparison of deep structure along three transects of the western North American continental margin. *Tectonics* 12, 1420–1435.

- FURRER, U. 1979: Stratigraphie des Doggers der östlichen Préalpes (Bern). *Eclogae geol. Helv.* 72, 623–672.
- FURRER, U. & SEPTFONTAINE, M. 1977: Nouvelles données biostratigraphiques dans le Dogger à faciès briançonnais des Préalpes médianes romandes (Suisse). *Eclogae geol. Helv.* 70, 717–737.
- GAGNEBIN, E. 1922: Préalpes entre Montreux et le Moléson et du Mont Pélerin, carte spéciale n°99. *Comm. géol. Suisse*, Basel.
- 1924: Description géologique des Préalpes bordières entre Montreux et Semsales. *Mém. Soc. vaud. Sci. nat.* 2, 1–72.
- 1940: Le front de la nappe de la Brèche et les plis des Préalpes médianes entre la Haute Pointe et le Roc d'Enfer (Haute-Savoie). *Bull. Soc. vaud. Sci. nat.* 61, 1–22.
- GENGE, E. 1957: Ein Beitrag zur Stratigraphie der S Klippendecke im Gebiet Spillgerten – Seehorn. *Eclogae geol. Helv.* 51, 151–211.
- GIROD, F. 1995: Cornettes de Bise. Diplôme, Lausanne.
- GORIN, G.E., SINGER, C. & AMBERGER, G. 1993: Structural configuration of the western Swiss Molasse Basin as defined by reflection seismic data. *Eclogae geol. Helv.* 86, 693–716.
- GOY-EGGENBERGER, D. & KÜBLER, B. 1990: Résultats préliminaires d'un essai de zonéographie métamorphique à travers les formations calcaires de la nappe de Morcles. *Schweiz. mineral. petrogr. Mitt.* 70, 83–88.
- GUELLEC, S., MUGNIER, J.L., TARDY, M. & ROURE, F. 1990: Neogene evolution of the western Alpine foreland in the light of ECORS data and balanced cross sections. In: *Deep structure of the Alps* (Ed. by ROURE, F., HEITZMANN, P. & POLINO, R.). *Mém. Soc. géol. Suisse* 1, 165–184.
- GUILLAUME, M. 1986: Révision stratigraphique des Couches Rouges de la nappe des Préalpes Médiannes Romandes. PhD thesis, Univ. Fribourg, n° 910.
- HEINZ, R. & ISENSCHMID, C. 1988: Mikrofazielle und stratigrafische Untersuchungen im Massifkalk (Malm) der Préalpes Médiannes. *Eclogae geol. Helv.* 81, 1–62.
- HEINZ, R.A. 1985: Mikrofazielle Untersuchungen des Massifkalks (Malm) der Préalpes Médiannes im Querschnitt Moléson Ruebli/Gummfluh. PhD thesis, Univ. Bern, n° 209.
- HOMWOOD, P. 1977: Ultrahelvetic and North-Penninic flysch of the Prealps: a general account. *Eclogae geol. Helv.* 70, 627–641.
- HOMWOOD, P., ACKERMANN, T., ANTOINE, P. & BARBIER, R. 1984: Sur l'origine de la nappe du Niesen et la limite entre les zones ultrahelvétiques et valaisanne. *C.R. Séances Acad. Sci. (Paris), sér. II* 299, 1055–1059.
- HOMWOOD, P. & LATELTIN, O. 1988: Classic Swiss clastics. Flysch and Molasse. *The Alpine connection. Geodynamica Acta* 2, 1–11.
- HSÜ, K.J. 1989: Time and place in Alpine orogenesis – the Fermor lecture. In: *Alpine Tectonics* (Ed. by COWARD, M.P., DIETRICH, D. & PARK, R.G.). *Geol. Soc. (London) Spec. Publ.* 45, 421–443.
- HUON, S., KÜBLER, B. & HUNZIKER, J. 1988: Identifications de mélanges de micas blancs par diffraction des rayons X: application à des séries carbonatées faiblement métamorphiques. *Schweiz. mineral. petrogr. Mitt.* 68, 185–202.
- ISENSCHMID, C. 1983: Der Malm im Mittelabschnitt der Préalpes médianes zwischen Thuner- und Genfersee. PhD thesis, Univ. Bern.
- JABOYEDOFF, M. & THÉLIN, P. in press: New data on low metamorphism in the Briançonnais domain of the Préalpes, Western Switzerland. *Europ. J. Mineral.*
- JACCARD, F. 1904: La région de la Brèche de la Hornfluh (Préalpes Bernoises). *Bull. Lab. Géol. Univ. Lausanne* 5, 1–205.
- 1907: La région Rübli-Gummfluh (Préalpes médianes, Suisse). *Bull. Soc. vaud. Sci. nat.* 43, 407–548.
- JEANBOURQUIN, P. 1986: Les Cornieules polymictes des Préalpes internes et de l'autochtone helvétique en Suisse romande. PhD thesis, Univ. Lausanne.
- 1988: Nouvelles observations sur les cornieules en Suisse occidentale. *Eclogae geol. Helv.* 81, 511–538.
- 1991: Ultrahelvétique et mélanges sur le dos des nappes helvétiques des Diablerets et du Wildhorn. *Bull. Soc. fribourg. Sci. nat.* 80, 76–104.
- 1992: Les mélanges suprahelvétiques dans le synclinal de Thônes (Massifs subalpins, Haute-Savoie, France): une nouvelle hypothèse de travail. *Arch. Sci. phys. nat. (Genève)* 45, 109–126.
- 1994: Early deformation of Ultrahelvetic mélanges in the Helvetic nappes (Western Swiss Alps). *J. struct. Geol.* 16, 1367–1383.
- JEANBOURQUIN, P. & GOY-EGGENBERGER, D. 1991: Mélanges suprahelvétiques: sédimentation et tectonique au front de la nappe de Morcles (Vaud, Suisse). *Géol. alp. (Grenoble)* 67, 43–62.
- JEANBOURQUIN, P. & LUALDI, A. 1994: Telogenesis and thrust-related fluid flow, the example of the Rocca Grande Rauhawacke (Ligurian Alps, Italy). *Ann. Tect.* 13, 70–84.

- JEANBOURQUIN, P., KINDLER, P. & DALL'AGNOLO, S. 1992: Les mélanges des Préalpes internes entre Arve et Rhône (Alpes occidentales franco-suisse). *Eclogae geol. Helv.* 85, 59–83.
- JEANNET, A. 1912–1913: Monographie géologique des Tours d'Ai. 1. Stratigraphie de la nappe rhétique, du Trias et du Lias des Préalpes médianes et de la zone interne. *Matér. Carte géol. Suisse n.s.* 32, 1–466.
- 1922: Das romanische Deckengebirge, Préalpes und Klippen. In: *Geologie der Schweiz* (Ed. by HEIM, ALB.). II/2, Tauchnitz, Leipzig, 589–676.
- KÜBLER, B., PITTION, J.-L., HEROUX, Y., CHAROLAIS, J. & WEIDMANN, M. 1979: Sur le pouvoir réflecteur de la vitrinite dans quelques roches du Jura, de la Molasse et des nappes préalpines, helvétiques et penniques. *Eclogae geol. Helv.* 72, 347–373.
- LEINE, L. 1971: Rauhacken und ihre Entstehung. *Geol. Rdsch.* 60, 448–524.
- LEMOINE, M. 1973: About gravity gliding tectonics in the western Alps. In: *Gravity and Tectonics* (Ed. by DE JONG, K.A. & SCHOLTEN, R.). J. Wiley and Sons, 201–216.
- 1988: Des nappes embryonnaires aux blocs basculés: évolution des idées et modèles sur l'histoire mésozoïque des Alpes occidentales. *Bull. Soc. géol. France*, 8ème série 4, 787–797.
- LEMOINE, M., BAS, T., ARNAUD-VANNEAU, A., ARNAUD, H., DUMONT, T., GIDON, M., BOURBON, M., GRACIANSKY, P.C., RUDKIEWICZ, J.L., MÉGARD-GALLI, J. & TRICART, P. 1986: The continental margin of the Mesozoic Tethys in the Western Alps. *Marine Petroleum Geol.* 3, 179–199.
- LEMPICKA-MÜNCH, A. & MASSON, H. 1993: Dépôts d'âge tertiaire de la nappe ultrahelvétique du Sex Mort entre la Simme et Adelboden (Préalpes). *Bull. Soc. vaud. Sci. nat.* 82, 369–389.
- LOMBARD, A. 1971: La nappe du Niesen et son flysch. *Matér. Carte géol. Suisse n.s.* 141, 1–251.
- 1974: Les Mosses, Feuille n°64, Notice explicative. *Comm. géol. Suisse*, Basel.
- LONFAT, F. 1965: Géologie de la partie centrale des Rochers de Château-d'Oex (Rübli-Gummfluh). *Matér. Carte géol. Suisse n.s.* 120, 1–86.
- LUGEON, M. 1896: La région de la Brèche du Chablais. *Bull. Serv. Carte géol. France* 7, 337–646.
- 1902: Les grandes nappes de recouvrement des Alpes du Chablais et de la Suisse. *Bull. Soc. géol. France* 4, 723–825.
- 1943: Une nouvelle hypothèse tectonique: la diverticulation. *Bull. Soc. vaud. Sci. nat.* 62, 301–303.
- LUGEON, M. & GAGNEBIN, E. 1941: Observations et vues nouvelles sur la géologie des Préalpes romandes. *Bull. Lab. Géol. Univ. Lausanne* 72, 1–90.
- MANCKTELOW, N.S. 1992: Neogene lateral extension during convergence in the Central Alps: evidence from interrelated faulting and backfolding around the Simplonpass (Switzerland). *Tectonophysics* 215, 295–317.
- MARCHANT, R. 1993: The Underground of the Western Alps. *Mem. Geol. (Lausanne)* 15.
- MARCHANT, R.H. & STAMPFLI, G.M. in press: Crustal and lithospheric structures of the Western Alps: geodynamic significance. In: *Deep structure of Switzerland – Results from NFP 20* (Ed. by LEHNER, P., HEITZMAN, P., FREI, W., HORSTMAYER, H., MUELLER, S., PFIFFNER, A. & STECK, A.). Birkhäuser AG, Basel.
- MARTHALER, M. & STAMPFLI, G. 1989: Les schistes lustrés à ophiolithes de la nappe du Tsaté: un ancien prisme d'accrétion issu de la marge active apulienne. *Schweiz. mineral. petrogr. Mitt.* 69, 211–216.
- MASSON, H. 1972: Sur l'origine de la cornieule par fracturation hydraulique. *Eclogae geol. Helv.* 65, 27–41.
- 1976: Un siècle de géologie dans des Préalpes: de la découverte des nappes à la recherche de leur dynamique. *Eclogae geol. Helv.* 69, 527–575.
- MASSON, H., BAUD, A., ESCHER, A., GABUS, J. & MARTHALER, M. 1980: Compte rendu de l'excursion de la Société Géologique Suisse: coupe Préalpes-Helvétique-Pennique en Suisse occidentale. *Eclogae geol. Helv.* 73, 331–349.
- MATTER, A., HOMEWOOD, P., CARON, C., RIGASSI, D., VAN STUIJVENBERG, J., WEIDMANN, M. & WINKLER, W. 1980: Flysch and Molasse of Western and Central Switzerland. In: *Geology of Switzerland* (Ed. by TRÜMPY, R.) Part B: geological excursions, excursion IV, Wepf, Basel, 261–293.
- MAURY, P. & RICOU, L.E. 1983: Le décrochement subbriançonnais: une nouvelle interprétation de la limite interne-externe des Alpes franco-italiennes. *Rev. Géol. dyn. Géogr. phys.* 24, 3–22.
- MAUVE, C.C. 1921: Geologische Untersuchungen im Molésongebiet. *Eclogae geol. Helv.* 16, 374–455.
- MCCONNELL, R.B. 1951: La nappe du Niesen et ses abords entre les Ormonts et la Sarine. *Matér. Carte géol. Suisse n.s.* 95, 1–94.
- MEISSER, N. 1992: Le cuivre natif et l'anilite de Vouvy, VS. *Cristallier Suisse* 9, 371–376.
- METTRAUX, M. 1989: Sédimentologie, paléotectonique et paléocéanographie des Préalpes Médianes (Suisse Romande) du Rhétien au Toarcien. PhD thesis, Univ. Fribourg, n° 947.
- METTRAUX, M. & MOSAR, J. 1989: Tectonique alpine et paléotectonique liasique dans les Préalpes Médianes en rive gauche du Rhône. *Eclogae geol. Helv.* 82, 517–540.

- MOSAR, J. 1988a: Métamorphisme transporté dans les Préalpes. *Schweiz. mineral. petrogr. Mitt.* 68, 77–94.
- 1988b: Structures, déformation et métamorphisme dans les Préalpes Romandes (Suisse). PhD thesis, Univ. Neuchâtel.
  - 1989: Déformation interne dans les Préalpes Médiannes (Suisse). *Eclogae geol. Helv.* 82, 765–793.
  - 1991: Géologie structurale dans les Préalpes médianes (Suisse). *Eclogae geol. Helv.* 84, 689–725.
  - 1994: Géologie structurale à l'est de Montreux (Préalpes médianes plastiques, Suisse). *Eclogae geol. Helv.* 87, 11–32.
- MOSAR, J. & BOREL, G. 1992: Paleostress from the Préalpes médianes (Switzerland). *Ann. Tect.* 6, 115–133.
- 1993: Le pli de Malatraix: fiction ou réalité. *Bull. Soc. vaud. Sci. nat.* 82, 319–343.
  - 1995: Préalpes geology: A non-exhaustive reference list. Internet: at Department of Earth Sciences, Lausanne Univ.: <http://www-sst.unil.ch>
- MÜLLER, W.H. 1982: Zur Entstehung der Rauhawacke. *Eclogae geol. Helv.* 75, 481–494.
- MÜLLER, I. & PLANCHEREL, R. 1982: Contribution à l'étude de l'hydrogéologie karstique du massif du Vanil Noir et de la chaîne des Gastlosen (Préalpes fribourgeoises, Suisse). *Bull. Soc. fribourg. Sci. nat.* 71, 102–132.
- MULLIS, J. 1979: The system methane–water as a geologi thermometer and barometer from the external part of the Central Alps. *Bull. Minéral.* 102, 526–536.
- NICOL, G. 1956: Geologie der östlichen Stockhorn-Kette (Berner Oberland). *Mitt. natf. Ges. Bern, N.F.* 13, 153–243.
- OBERHÄNSLI, R., HOFMANN, B. & GRUNER, U. 1985: Ein Massivsulfidvorkommen in der Trias der Präalpen. *Schweiz. mineral. petrogr. Mitt.* 65, 95–110.
- ODIN, G.S. 1994: Geological time-scale (1994). *C.R. Acad. Sci. (Paris), série II* 318, 59–71.
- PAGE, C. 1969: Observations géologiques sur les Préalpes au NW des Gastlosen orientales. *Bull. Soc. fribourg. Sci. nat.* 58, 83–177.
- PAVONI, N. 1987: Zur Seismotektonik der Nordschweiz. *Eclogae geol. Helv.* 80, 461–472.
- PETERHANS, E. 1923: Sur la tectonique des Préalpes entre Meillerie et Saint-Gingolph (Haute-Savoie). *Bull. Soc. géol. France* 4. série 23, 51–56.
- PLANCHEREL, R. 1976: Essai d'interprétation de la dislocation transversale de Bellegarde-Lac Noir (Préalpes médianes Fribourgeoises). *Eclogae geol. Helv.* 65, 461–469.
- 1979: Aspects de la déformation en grand dans les Préalpes médianes plastiques entre Rhône et Aar. *Eclogae geol. Helv.* 72, 145–214.
  - 1990: Les Préalpes du Chablais – Présentation générale. In: Suisse Lémanique pays de Genève et Chablais. Guides Géologiques Régionaux (Ed. by CHAROLAIS, J. & BADOUX, H.). Masson, 183–190.
- PLANCHEREL, R. & WEIDMANN, M. 1972: La zone anticlinale complexe de la Tinière (Préalpes médianes vaudaises). *Eclogae geol. Helv.* 65, 75–91.
- PYTHON DUPASQUIER, C. 1990: La Formation de l'Intyamou (Crétacé moyen) des Préalpes médianes romandes. PhD thesis, Univ. Fribourg, No. 978.
- RABOWSKI, F. 1912: Simmenthal et Diemtighal, carte spéciale n°69. *Comm. géol. Suisse.*
- 1920: Les Préalpes entre le Simmental et le Diemtighal. *Matér. Carte Géol. Suisse n.s.* 35.
- REVERTERA, G.C. 1926: Auszug aus der geologischen Monographie der Kette des Monts Cray (SW-Abschnitt der Vanil Noir-Kette) (Préalpes médianes). PhD thesis, Univ. Fribourg.
- RICOU, L.E. 1984: Les Alpes occidentales: chaîne de décrochement. *Bull. Soc. géol. France* 7/26, 861–874.
- RICOU, L.E. & SIDDANS, A.W.B. 1986: Collision tectonics in the Western Alps. *Spec. Publ. geol. Soc. London* 19, 229–244.
- SAMBETH, U. & PAVONI, N. 1988: A seismotectonic investigation in the Geneva Basin, southern Jura mountains. *Eclogae geol. Helv.* 81, 433–440.
- SARTORI, M. 1987: Blocs basculés briançonnais en relation avec leur socle originel dans la nappe de Siviez-Mischabel (Valais, Suisse). *C.R. Acad. Sci. (Paris), série II* 305, 999–1005.
- SARTORI, M. 1990: L'unité du Barrhorn (Zone Pennique, Valais, Suisse). *Mem. Geol. (Lausanne)* 6, 1–156.
- SARTORI, M. & MARTHALER, M. 1994: Exemples de relations socle-couverture dans les nappes penniques du Val d'Hérens. *Schweiz. mineral. petrogr. Mitt.* 74, 503–509.
- SCHAAD, W. 1995: Die Entstehung von Rauhawacken durch die Verkarstung von Gips. *Eclogae geol. Helv.* 88, 59–90.
- SCHARDT, H. 1884: Etudes géologiques sur le Pays d'Enhaut vaudois. *Bull. Soc. vaud. Sci. nat.* 20, 1–183.
- 1893a: Coup d'oeil sur la structure géologique des environs de Montreux. *Bull. Soc. vaud. Sci. nat.* 29, 241–255.

- 1893b: Sur l'origine des Préalpes romandes. *Arch. Sci. phys. nat.* (Genève) 3, 570–583.
  - 1898: Les régions exotiques du versant Nord des Alpes Suisses. Préalpes du Chablais et du Stockhorn et les Klippes. *Bull. Soc. vaud. Sci. nat.* 34, 113–219.
  - 1907: Les vues modernes sur la tectonique et l'origine de la chaîne des Alpes. *Arch. Sci. phys. nat.* (Genève) 4, 356–385 / 483–496.
- SCHMID, S.M., RÜCK, P. & SCHREURS, G. 1990: The significance of the Schams nappes for the reconstruction of the paleotectonic and orogenic evolution of the Penninic zone along the NFP-20 east traverse (Grisons, eastern Switzerland). In: *Deep Structure of the Alps* (Ed. by ROURE, F., HEITZMANN, P. & POLINO, R.). *Mém. Soc. géol. Suisse* 1, 263–287.
- SCHROEDER, W.J. 1939: La Brèche du Chablais entre Giffre et Drance et les roches éruptives des Gets. *Arch. Sci. phys. nat.* (Genève) 21, 1–138.
- SEPTFONTAINE, M. 1983: Le Dogger des Préalpes médianes suisses et françaises (Stratigraphie, évolution paléogéographique et paléotectonique). *Mém. Soc. helv. Sci. nat.* 97, 1–121.
- 1995: Large-scale progressive unconformities in Jurassic strata of the Préalpes S of lake Geneva: interpretation as synsedimentary inversion structures. *Eclogae geol. Helv.* in press.
- SEPTFONTAINE, M. & WERNLI, R. 1972: Contribution à la géologie des environs de Tanninges (Haute-Savoie). *Arch. Sci. phys. nat.* (Genève) 25, 113–128.
- SEWARD, D. & MANCKTELOW, N.S. 1994: Neogene kinematics of the central and western Alps: Evidence from fission-track dating. *Geology* 22, 803–806.
- STALDER, H.A. & TOURAY, J.C. 1970: Fensterquarze mit Methan-Einschlüssen aus dem westlichen Teil der Schweizer Kalkalpen. *Schweiz. mineral. petrogr. Mitt.* 50, 109–130.
- STAMPFLI, G.M. 1993: Le Briançonnais, terrain exotique dans les Alpes? *Eclogae geol. Helv.* 86, 1–45.
- STAMPFLI, G.M. & MARCHANT, R.H. in press: Geodynamic evolution of the Tethyan margins of the Western Alps. In: *Deep structure of Switzerland – Results from NFP 20* (Ed. by LEHNER, P., HEITZMAN, P., FREI, W., HORSTMAYER, H., MUELLER, S., PFIFFNER, A. & STECK, A.). Birkhäuser AG, Basel.
- STAMPFLI, G.M. & MARTHALER, M. 1990: Divergent and convergent margins in the North-Western alps, confrontation to actualistic models. *Geodin. Acta* 4, 159–184.
- STECK, A. 1984: Structures de déformations tertiaires dans les Alpes centrales (transversale Aar-Simplon-Ossola). *Eclogae geol. Helv.* 77, 55–100.
- 1990: Une carte des zones de cisaillement ductile des Alpes Centrales. *Eclogae geol. Helv.* 83, 603–627.
- STECK, A. & HUNZIKER, J. 1994: The tertiary structural and thermal evolution of the central alps – compressional and extensional structures in an orogenic belt. *Tectonophysics* 238, 229–254.
- STECK, A., EPARD, J.-L., ESCHER, A., LEHNER, P., MARCHANT, R. & MASSON, H. in press: Geological interpretation of the seismic profiles through western Switzerland: Rawil (W1), Val d'Anniviers (W2), Matternal (W3), Zmutt-Zermatt-Findelen (W4) and Val de Bagnes (W5). In: *Deep structure of the Swiss Alps - Results from NFP 20* (Ed. by LEHNER, P., HEITZMAN, P., FREI, W., HORSTMAYER, H., MUELLER, S., PFIFFNER, A. & STECK, A.). Birkhäuser AG., Basel.
- STEFFEN, D., JAQUES, C., NYDEGGER, T., PETROONS, D. & WILDI, W. 1993: La Brèche du Chablais à son extrémité occidentale (Hte-Savoie, France): Sédimentologie, éléments stratigraphiques et interprétation paléogéographique. *Eclogae geol. Helv.* 86, 543–568.
- STUDER, B. 1827: Remarques géognostiques sur quelques parties de la chaîne septentrionale des Alpes. *Ann. Sci. nat. Paris* 11, 1–47.
- SUPPE, J., CHOU, G.T. & HOOK, S.C. 1992: Rates of folding and faulting determined from growth strata. In: *Thrust Tectonics* (Ed. by McCLAY, K.R.). Chapman & Hall, 105–121.
- TERCIER, J. 1948: Le Flysch dans la sédimentation alpine. *Eclogae geol. Helv.* 40, 163–168.
- THÉLIN, P., SARTORI, M., GOUFFON, Y. & CHESSEX, R. 1993: The Pre-Alpine basement of the Briançonnais (Wallis, Switzerland). In: *Pre-Mesozoic geology in the Alps* (Ed. by RAUMER, v.J.F. & NEUBAUER, F.). Springer, Heidelberg, 297–315.
- THURY, M. 1973: Der Lias der östlichen Préalpes médianes zwischen Boltigen und Spiez. PhD thesis, Univ. Bern.
- TRICART, P. 1984: From passive margin to continental collision: a tectonic scenario for the Western Alps. *Amer. J. Sci.* 284, 97–120.
- TRÜMPY, R. 1955: Remarques sur la corrélation des unités penniques externes entre la Savoie et le Valais et sur l'origine des nappes préalpines. *Bull. Soc. géol. France* 6, 217–231.
- 1960: Paleotectonic evolution of the Central and Western Alps. *Bull. geol. Soc. Amer.* 71, 843–908.
  - 1980: *Geology of Switzerland: a guide-book.* (Ed. by TRÜMPY, R.). Wepf and Co., Basel.

- TRÜMPY, R. & BERSIER, A. 1954: Les éléments des conglomérats oligocènes du Mont-Pélerin. *Pétrographie, statistique, origine*. *Eclogae geol. Helv.* 47, 119–166.
- VOLLMAYR, T. 1992: Strukturelle Ergebnisse der Kohlenwasserstoff-Exploration im Gebiet von Thun, Schweiz. *Eclogae geol. Helv.* 85, 531–539.
- WEIDMANN, M. 1971: Cagneule ou Cornieule? *Eclogae geol. Helv.* 64, 47–51.
- 1972: Le front de la Brèche du Chablais dans le secteur de Saint Jean d'Aulph (Haute-Savoie). Quelques remarques générales sur la nappe de la Brèche. *Géol. alp. (Grenoble)* 48, 229–248.
  - 1988: Lausanne, Feuille n°85, notice. Service hydrol. géol. national.
  - 1992: Châtel-Saint-Denis, Feuille n°92, notice. Service hydrol. géol. national.
- WEIDMANN, M., HOMEWOOD, P., CARON, C. & BAUD, A. 1976: Réhabilitation de la Zone Submédiane des Préalpes. *Eclogae geol. Helv.* 69, 265–277.
- WEISS, H. 1949: Stratigraphie und Mikrofauna des Klippenmalm. PhD thesis, Univ. Zürich.
- WENGEN, W. 1924: Geologische Untersuchungen im Gebiet des Vanil Noir (Gruyère). PhD thesis, ETH Zürich.
- WERNLI, R. & BRÖNNIMANN, P. 1973: Le Verrucano et le Trias de Tanninges (Haute-Savoie, France) Lithologie et micropaléontologie. *Arch. Sci. phys. nat. (Genève)* 26, 79–92.
- WILDI, W. 1985: Heavy mineral distribution and dispersal pattern in penninic and ligurian flysch basins (Alps, northern Apennines). *G. Geol. Bologna, ser. 3*, 47, 77–99.
- WILLETT, S., BEAUMONT, C. & FULLSACK, P. 1993: Mechanical model for the tectonics of doubly vergent compressional orogens. *Geology* 21, 371–374.
- WINKLER, W. 1984: Palaeooccurrents and petrography of the Gurnigel-Schlieren Flysch: a basin analysis. *Sediment Geol.* 40, 169–189.
- ZAHNER, P. & MOSAR, J. 1993: Résultats des recherches effectuées sur le site potentiel du Bois de la Glaive (Commune d'Ollon, VD). *Cédra Rapport Technique* 93, 1–54.
- ZINGG, M.A. 1984: Die Tertiäre Pb-Zn-Cu-Sulfidvererzung im Sackgraben, Engstligental (Berner Oberland). *Schweiz. mineral. petrogr. Mitt.* 64, 215–225.

Manuscript received May 7, 1995

Revision accepted August 9, 1995

While in press the following two papers, relevant to the Préalpes Geology, were published and could not be referenced in our paper:

- BOREL, G. 1996: Préalpes médianes: courbes de subsidence et implications géodynamiques. *Bull. Soc. vaud. Sci. nat.* 83, 293–315.
- SOMMARUGA, A. 1995: Tectonics of the Central Jura and the Molasse Basin. New insights from the interpretation of seismic reflection data. *Bull. Soc. neuchât. Sci. Nat.* 118, 95–108.

