Zeitschrift:	Entomologica Basiliensia
Herausgeber:	Naturhistorisches Museum Basel, Entomologische Sammlungen
Band:	11 (1986)
Artikel:	Comparison of diving beetles (Coleoptera, Dytiscidae) collected from above and from below a beaver (Castor) dam
Autor:	Swenson, G.
DOI:	https://doi.org/10.5169/seals-980553

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 11.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Entomologica Basiliensia	11	309-312	1986	ISSN 0253-2484	
--------------------------	----	---------	------	----------------	--

Comparison of Diving Beetles (Coleoptera, Dytiscidae) Collected from Above and from Below a Beaver (Castor) Dam

by G. Swenson

Abstract: Collections of dytiscid water beetles were made in a flooded *Carex lacustris* stand situated behind an active beaver (Castor) dam. These were compared to beetles collected in a drier area situated below the same beaver dam and dominated by *Carex rostrata*. Much of the collection in the site above the dam was made in the shade of alder (*Alnus rugosa*). The ecologically different sites, despite geographical proximity, yielded different collections.

Key words: Coleoptera Dytiscidae – *Hydroporus paugus* – beaver – *Carex* – Habitat comparison.

This study continues the work on regional water beetle populations in the Finger Lakes region. Previous studies focused on wide-ranging collections in this area. The next step appeared to be studying the collecting sites more closely and comparing different environments. Two sites were found in Michigan Hollow, where I had collected previously. They seemed to be appropriate sites for comparative study because, even though they were near each other, they had different ecological features.

The two sites are located near the west side of Michigan Hollow Road about 5 km south of Ithaca, New York. One of the collection sites was a deep water site dominated by *Alnus rugosa* and *Carex lacustris*. The water table was always 3–15 cm above the surface because the site was behind a beaver dam. Much of the collection in this site was made in the shade of alder (*Alnus rugosa*).

Collections were also made in a dry site dominated by *Carex* rostrata and *Leersia oryzoides*, where the water table was always below the surface except for small scattered pools varying in depth from 10–40 cm and for a discontinuous network of small rivulets draining the alder thicket. These little streams were about 15–40 cm wide and 20–30 cm deep. It was from these pools and streams that the collections were made. No distinct difference in the species of beetles collected from the two different sources was notable.

G. Swenson

The collections were made primarily with a D-frame aquatic net, sporadically over ten years and weekly during the summer of 1983 in the *C. lacustris* site and only in the summer and early fall of 1983 and the spring and summer of 1984 for the *C. rostrata* site.

The species collected are listed in Table 1 according to those found in one or the other of the two sites and to those found in both sites.

Reference to Blackwelder's Red List and to Larson has indicated that most of the species are widely distributed in Northeastern America, BLACKWELDER (1973), LARSON (1975), (1983 personal communication).

A.	С.	rostrata	site

B. C. lacustris site

C. both sites

Hydroporus melsheimeri Fall Hydroporus notabilis LeConte Hydroporus signatus Mannerheim * Agabus subfuscatus Sharp Agabus semivittatus LeConte Agabus seriatus (Say) Agabus ambiguus (Say) * Agabus erythropterus (Say) Agabus erichsoni (Gemminger & Harold) Agabus leptapsis (LeConte) Copelatus glyphicus Say Dytiscus fasciventris Say

Laccophilus maculosus Say Hygrotus nubilus LeConte Agabus anthracinus Mannerheim * Ilybius angustior (Gyllenhall) Ilybius pleuriticus LeConte Coptotomus interrogatus Fabricius Dytiscus verticalis Say Hydaticus modestus Sharp Acilius sylvanus Hilsenhoff

Hygrotus sayi J. Balfour-Browne Hygrotus laccophilinus (LeConte) Hygrotus impressopunctatus Schaller Hydroporus undulatus Say Hydroporus paugus Fall * Hydroporus dentellus Fall Hydroporus niger Say ' Hydroporus striola Gyllenhall * Agabus semipunctatus Kirby Agabus phaeopterus (Kirby) Agabus gagates Aubé * Ilybius biguttulus Germar * Colymbetes sculptilis Harris Rhantus binotatus Harris Acilius mediatus (Say) Graphoderus fascicollis Harris

Tab. 1: Species collected listed according to site of collection. Asterisk after a species name indicates that species was collected in large numbers.

However, in the Finger Lakes, of the 37 species listed in table 1, I have previously reported collecting 28 species, Swenson (1979, 1982). Of the remaining 9 species only 3 have not been collected by me at any other site. They are a few specimens of *Agabus leptapsis* and one female *Agabus erythropterus* from the *C. rostrata* site and *Ilybius pleuriticus* from the *C. lacustris* site. Further, *Hydroporus paugus*, which previously I have collected one or two at a time and not often from various sites, has appeared in great numbers in the *C. rostrata* site. Many collected from that site were teneral. These tenerals ranged in appearance in the collections from June 2^{nd} to August 5^{th} but most of them were collected in July. The *H. paugus* generally were collected from both the pools and the rivulets in the *C. rostrata* site.

Conclusions

First of all, this study suggests that site comparisons should be both intensive in collection and extensive in years of continuing study. Unless one can collect and identify larvae and teneral adults it is difficult to be sure whether a given species is a transient or a breeding resident. Also, different additional sets of compared sites should be studied to see if consistent patterns of species occupancy can be determined.

It was of interest to note that the number of individuals in each of the various species collected was low in the *C. lacustris* site, especially as compared to the *C. rostrata* site when a given species appeared in both sites. Could it be that the open area is a more natural flyway for migrants than the shaded *C. lacustris* site is? Again, identification of larvae and tenerals, when found, would reduce uncertainty as to which species are (regular?) transients and which are breeding residents.

Several puzzles emerged. Acilius semisulcatus, which I have found almost everywhere else I have collected was missing in the collections in these two sites. Agabus leptapsis appeared in small numbers in the *C. rostrata* site in 1983 and appeared not at all in 1984. Agabus semivittatus and Agabus seriatus, which I usually find in lotic conditions, appeared in the *C. rostrata* site, one or two of the few collected appearing in brown-water pools.

The appearance in great number of *Hydroporus signatus* and *Agabus ambiguus* exclusively in the *C. rostrata* site and of *Agabus anthracinus* exclusively in the *C. lacustris* site suggests a favored resi-

dency. However, no tenerals were noted for these three species, suggesting further study is necessary.

In spite of low collection numbers for many of the species identified in this study and of other problems discussed above, there does appear to be a shadow of difference in the species profiles for these two adjacent but ecologically different sites. The level of significant difference is yet to be discovered for these two different environments.

References

- BLACKWELDER, R. E. (1973): Checklist of the Dytiscidae of Canada, United States, Mexico, Central American and the West Indies. Biol. Res. Inst. of Amer. Inc.
- LARSON, D. J. (1975): Dytiscidae of Alberta. Quaest. Ent. 11: 245-498.

SWENSON, G. (1979): Water beetle records from the vicinity of Ithaca, New York (Coleoptera: Dytiscidae) Col. Bull. 33 (4): 477–479.

SWENSON, G. (1982): Water beetle records from the vicinity of Ithaca, New York (Coleoptera: Dytiscidae and Hydrophilidae) Col. Bull. 36 (2): 350–351.

Author's address: Prof. Dr George Swenson Ithaca College Department of Biology Ithaca, New York, 14850 USA