

§ 3. Properties of the Exponents.

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **2 (1956)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **01.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

The converse is also true, namely that any crystallographic reflection group over R corresponds to some compact semi-simple Lie Group.

It can be shown that T may be covered by a Euclidean space R^r in such a way that the *singular elements* of T (i.e. those whose normalisers are of dimension strictly greater than r) map into hyperplanes of R^r , and further, if the identity of G maps into the origin O of R^r , then those planes passing through O are precisely the hyperplanes of reflection of the Weyl group W . The whole set of hyperplanes form a configuration known as the *diagram* of the Lie Group G and it has the property that reflection in any one of the planes leaves the diagram, as a whole invariant.

Now let W be *any* reflection group over R expressed in orthogonal form, then W may be considered as operating on some sphere S^{r-1} whose centre is at O . The hyperplanes of reflection divide the surface of the sphere into spherical polytopes and it has been shown [5; p. 190] that each of these is necessarily a simplex or a direct product of simplexes. Further, the r hyperplanes that cut the sphere in the faces of one of these polytopes form a *fundamental set* in that the corresponding reflections generate the group. Furthermore, the volume bounded by these hyperplanes forms a fundamental region for W . A property of this fundamental set is given in (vi) of § 3.

Considering again the diagram of the Lie Group G , pick out a fundamental set of hyperplanes through O , defining a fundamental region of the Weyl group. Then the part of the diagram of G that lies within this fundamental region is called a *Weyl chamber*. The Weyl chamber of the group G_2 (the group of automorphisms of the Cayley matrix algebra) is illustrated in (iv) of § 3, where, for the present, the numerals are to be ignored.

§ 3. PROPERTIES OF THE EXPONENTS.

(i) The ring of polynomial invariants of an r dimensional reflection group W over k is the ring $k[I_1, I_2, \dots, I_r]$ where I_i is a polynomial invariant of degree $(m_i + 1)$. The I_i are uniquely determined by this property and are called the *basic invariants* of the group W . The m_i are called the *exponents* of W .

This theorem was proved by CHEVALLEY for k of characteristic zero [4] and a partial converse (for $k = \mathbb{C}$) by TODD [9; p. 282].

From this result we can draw several conclusions. Firstly it implies a formal identity in power series in t :

$$\prod_{i=1}^r \frac{1}{(1 - t^{m_i+1})} = \sum_{j=0}^{\infty} a_j t^j = \frac{1}{\omega} \sum_{S \in W} \frac{1}{|1 - tS|}$$

where ω is the order of W , and a_j is the number of linearly independent polynomial invariants of W of degree j (which, by the above result, is the number of monomials in the I_i of total degree j , and hence is the coefficient of t^j in the expansion of the product on the left). The identity on the right is the classical result of MOLIEREN [3; p. 300]. Considering only the first and last terms of this identity, after a small amount of manipulation we deduce

$$(a) \quad \prod_{i=1}^r (1 + m_i) = \omega$$

$$(b) \quad \sum m_i = b_1 = \text{number of reflections in } W.$$

(ii) More generally there is an identity

$$\prod_{i=1}^r (1 + m_i t) = \sum_{j=1}^r b_j t^j$$

where b_j is the number of transformations in W that leave pointwise invariant a linear subspace of V^r of exactly $(r - j)$ dimensions. Putting $t = 1$ we obtain (a) above, and (b) merely states the equality of coefficients of t on both sides. No general proof is known, but this result has been verified for all reflection groups over \mathbb{C} .

(iii) Writing $H^*(X, \mathbb{R})$ for the cohomology ring of a space X with coefficients in \mathbb{R} , S^n for the n -sphere, and $\Lambda(x_1, x_2, \dots, x_r)$ for the graded exterior algebra on r generators, then for any compact semi-simple Lie group G , we have

$$\begin{aligned} H^*(G, \mathbb{R}) &= H^*(S^{2m_1+1} \times S^{2m_2+1} \times \dots \times S^{2m_r+1}, \mathbb{R}), \\ &= \Lambda(x_1, x_2, \dots, x_r), \quad \partial^0 x_i = 2m_i + 1. \end{aligned}$$

This is the celebrated HOPF theorem [7] except for the number and dimensions of the spheres (or of the generators of Λ). The relation between the dimensions and the exponents of W is due to CHEVALLEY [4]. An independent proof has been given by A. BOREL [1].

(iv) Consider the Weyl chamber of a connected semi-simple Lie group G , whose vertex is O . Then each simplex Δ_r of the chamber is to be labelled with an integer equal to the number of intersections with the hyperplanes of a line joining O to an interior point of Δ_r . It is trivially verified that such an integer is uniquely defined, i.e. is independent of the interior point chosen. Then we have


$$\prod_{i=0}^r \frac{1}{(1 - t^{m_i})} = \sum_{i=0}^{\infty} c_i t^i$$

where c_i is the number of simplexes in the Weyl chamber labelled with the integer i .

By way of example the Weyl chamber of G_2 is illustrated below. Here,

$$\begin{aligned} c_0 &= c_1 = c_2 = c_3 = c_4 = 1, \\ c_5 &= c_6 = c_7 = c_8 = c_9 = 2, \text{ etc.} \end{aligned}$$

The exponents are $m_1 = 1$ and $m_2 = 5$.



The statement of this result is only meaningful if $k = \mathbb{R}$ and W is crystallographic, and then it can be proved as follows:

From (iii) by the theory of spectral sequences

$$H^*(\Omega_G, R) = R[u_1, u_2, \dots, u_r], \quad \partial^0 u_i = 2m_i$$

where the right side is the graded ring of polynomials over R and Ω_G is the space of loops on G . Hence if we write $P(\Omega_G, t)$ (the Poincaré polynomial of Ω_G in t) for $\sum_{i=0}^{\infty} t^i \dim H^i(\Omega_G, R)$ we have

$$P(\Omega_G, t) = \prod_{i=1}^r \frac{1}{(1 - t^{2m_i})}$$

BOTT, by the use of Morse theory [2] proved that

$$\sum_{i=0}^{\infty} c_i t^{2i} = P(\Omega_G, t)$$

from which the given result follows immediately.

(v) The Jacobian of the basic invariants I_i ,

$$J = \frac{\partial (I_1, I_2, \dots, I_r)}{\partial (x_1, x_2, \dots, x_r)}$$

factorises into $\sum m_i$ linear forms, which, when equated to zero, give the hyperplanes of reflection of W , and each hyperplane is repeated $p - 1$ times where p is the order of the corresponding reflection.

Where all the reflections are of order 2, a very simple argument proves this result in a more rigorous manner than that of RACHA [6; p. 775]. For (b) of (i) implies that the degree of J is equal to the number of reflections, and the fact that J changes sign when operated on by a reflection in W implies that the equation of each hyperplane is a factor of J . This proves the result. More generally it can be proved over any field of zero characteristic for reflections of any order.

An interesting conjecture extends this result. In (ii) we defined b_j as the number of transformations in the group that leave a linear subspace of $n - j$ dimensions invariant. The set of all these linear subspaces forms a reducible algebraic variety of dimension $n - j$ and of degree b_j , and it is conjectured that this is given by equating to zero all the $(n - j + 1)$ -rowed

minors of the functional matrix $\partial I_i / \partial x_j$. This conjecture has neither been proved or verified.

(vi) This final property is the only one that holds for *irreducible* groups only. Suppose that r reflections in W serve to generate W (this is always the case for $k = R$, but not for $k = C$). Then it is possible to pick this set of generating reflections so that their product has characteristic roots

$$\exp\left(\frac{2\pi i m_j}{h}\right), j = 1, 2, \dots, r; h = \max(m_j) + 1$$

When $k = R$ it suffices to choose the reflections as those of a fundamental set and then take their product in any order [6; p. 765]. In this case also h has geometric significance as the number of sides of the PETRIE polygon [5; p. 223]. For $k = C$ no general rule for the selection of the correct set of reflections has been given.

This result has been verified for $k = C$, and general proofs are known for $k = R$, $r = 2, 3$ [6; p. 772].

REFERENCES

1. BOREL, Sur la cohomologie des espaces fibres principaux et des espaces homogènes de groupes de Lie compacts. *Annals of Math.*, 57 (1953), pp. 115-207.
2. BOTT, On torsion in Lie groups. *Proc. Nat. Academy of Sciences U.S.A.*, 40 (1954), pp. 586-588.
3. BURNSIDE, *Theory of Groups* (Cambridge, 1911).
4. CHEVALLEY, Invariants of Finite Groups generated by Reflexions. *American J. of Math.*, 77 (1955), pp. 778-782.
5. COXETER, *Regular Polytopes* (London, 1948, New York, 1949).
6. ——— The product of the generators of a group generated by reflections. *Duke Math. J.*, 18 (1951), pp. 765-782.
7. HOPF, Über die Topologie der Gruppen-Mannigfaltigkeiten und ihre Verallgemeinerungen. *Annals of Math.* (2), 42 (1941), pp. 22-52.
8. SHEPHARD, Unitary groups generated by reflections. *Can. J. of Math.*, 5 (1953), pp. 364-383.
9. ——— and TODD, Finite unitary reflection groups. *Can. J. of Math.*, 6 (1954), pp. 274-304.
10. STIEFEL, Über eine Beziehung zwischen geschlossenen Lie'schen Gruppen und diskontinuierlichen Bewegungsgruppen, etc. *Comm. Math. Helv.*, 14 (1941), pp. 350-380.