

Zeitschrift: L'Enseignement Mathématique
Band: 4 (1958)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON CERTAIN ARITHMETICAL FUNCTIONS RELATED TO A NON-LINEAR PARTIAL DIFFERENTIAL EQUATION
Kapitel: 2. Umbral Relations.
Autor: Basoco, M. A.
DOI: <https://doi.org/10.5169/seals-34625>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 18.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$(11) \quad U_k = (2^{2k} - 1) V_k = (-1)^k (2^{2k} - 1) \frac{B_k}{4k}.$$

As is well known, the double series occurring in (1), (4), (5), (6) are absolutely convergent for $k > 1$; for $k = 1$, the convergence is conditional. However, as has been shown by HURWITZ [2] in the case of (1), if the summation is first carried out with respect to m and then with respect to n , the resulting sum agrees with (2) with $k = 1$. For this case ($k = 1$) similar conditions hold for (4), (5) and (6). These matters are of relevance in studying certain modular transformations of these functions to be discussed later.

2. UMBRAL RELATIONS.

The functions defined in what precedes arise in a natural manner as a consequence of the well-known fact that the Jacobi theta functions are solutions of the partial differential equation

$$(12) \quad \frac{\partial^2 z}{\partial s^2} = 2 \frac{\partial z}{\partial t}, \quad z = \theta_r(\nu, \tau), \quad (r = 1, 2, 3, 4),$$

with $s = 2\pi\nu$ and $-t = 2\pi i\tau$, and, what appears to be less well-known, that the functions $u = \ln\theta_r(\nu, \tau)$ satisfy the non-linear equation:

$$(13) \quad \frac{\partial^2 u}{\partial s^2} = 2 \frac{\partial u}{\partial t} - \left(\frac{\partial u}{\partial s}\right)^2.$$

Here, the notation for the theta function is that used in TANNERY-MOLK's treatise [4].

The arithmetical consequence of (13) can best be obtained through the use of the infinite product representation of $\theta_r(\nu, \tau)$. It is found that the calculations needed are greatly facilitated and the results obtained very simply expressed in a symbolic form through an application of the umbral calculus of BLISSARD and LUCAS [5]. It is not feasible to give details for all cases and we merely indicate briefly the nature of the calculations for the case $r = 4$. Thus, since,

$$(14) \quad \theta_4(\nu, \tau) = Q_0 \prod_{n=1}^{\infty} (1 - q^{2n-1} e^{2\pi i \nu}) (1 - q^{2n-1} e^{-2\pi i \nu}),$$

$$Q_0 = \prod_{n=1}^{\infty} (1 - q^{2n}), \quad q = e^{\pi i \tau},$$

and taking into account the change in variables from (ν, τ) to (s, t) it is found that if $u(s, t) = \ln \theta_4(\nu, \tau)$, then

$$(15) \quad \frac{\partial u}{\partial s} = 2 \sum_{n=1}^{\infty} \frac{q^n}{1 - q^{2n}} \sin ns, \quad q = e^{-t/2},$$

and

$$(16) \quad \frac{\partial^2 u}{\partial s^2} = 2 \sum_{n=1}^{\infty} \frac{nq^n}{1 - q^{2n}} \cos ns;$$

moreover,

$$(17) \quad \frac{\partial u}{\partial t} = \sum_{n=1}^{\infty} \frac{nq^n}{1 - q^{2n}} - 2 \frac{\partial}{\partial t} \left\{ \sum_{n=1}^{\infty} \frac{q^n}{1 - q^{2n}} \frac{\cos ns}{n} \right\}$$

provided $\text{Ret} \pm 2\text{Im} s > 0$ in (15), (16) and (17).

Now, in (15) replace $\sin ns$ by its power series development and interchange the order of summation to obtain

$$(15)_1 \quad \frac{\partial u}{\partial s} = 2 \sum_{k=1}^{\infty} (-1)^{k-1} \frac{s^{2k-1}}{(2k-1)!} \Psi_{2k-1}(t).$$

Hence if Ψ is the umbra of the sequence $\{\Psi_{2k-1}(t)\}$ we may write symbolically:

$$(18) \quad \frac{\partial u}{\partial s} \cong 2 \sin \Psi s, \quad \frac{\partial^2 u}{\partial s^2} \cong 2 \Psi \cos \Psi s.$$

Similarly, a more extended calculation shows that

$$(19) \quad \frac{\partial u}{\partial t} \cong \Psi^{(1)} + 2 \frac{\partial}{\partial t} \frac{1 - \cos \Psi s}{\Psi}.$$

In (18) and (19), in order to pass from symbolic equality to actual equality, the functions $\sin \Psi s$, $\cos \Psi s$ and $(1 - \cos \Psi s)/\Psi$ are to be expanded in powers of s and then the exponents in the

powers of Ψ are lowered into subscripts; thus $\Psi^{(1)}$ would then be written $\Psi_1(t)$.

If (18) and (19) are substituted in (13), there results the following umbral identity:

$$(20) \quad \Psi (1 - \cos \Psi_s) + 2 \frac{\partial}{\partial t} \left(\frac{1 - \cos \Psi_s}{\Psi} \right) = 2 \sin \Psi_s * \sin \Psi_s ,$$

where the asterisk (*) indicates umbral multiplication.

For the cases $r = 2, 3$, rather extensive calculations show that umbral identities of the same form exist. We may therefore state the following result which is implied by the non-linear equation (13).

Theorem 1: "Let Ψ, X, Φ be respectively the umbrae of the sequences $\{\Psi_{2k-1}(t)\}$, $\{X_{2k-1}(t)\}$, and $\{\Phi_{2k-1}(t)\}$. If γ is one of these umbrae, then the following umbral identity holds:

$$(21) \quad \gamma (1 - \cos \gamma_s) + 2 \frac{\partial}{\partial t} \left(\frac{1 - \cos \gamma_s}{\gamma} \right) = 2 \sin \gamma_s * \sin \gamma_s .$$

3. RECURRENCES.

It is clear that (20) implies a recurrence relation for the functions $\Psi_j(t)$, and indeed, Theorem 1 yields the following.

Theorem 2: "Let $\gamma_j(t)$ be $\Psi_j(t)$, $X_j(t)$ or $\Phi_j(t)$; then the following recurrence holds:

$$(22) \quad \frac{d}{dt} \gamma_{2n-1}(t) + \frac{1}{2} \gamma_{2n+1}(t) = \sum_{k=0}^{n-1} \binom{2n}{2k+1} \gamma_{2k+1}(t) \gamma_{2n-2k-1}(t) ,$$

and hence $\gamma_{2n+1}(t)$ is a polynomial in $\gamma_1(t)$ and its derivatives up to order n ."

This result, in turn, implies the following

Theorem 3: "Let $\rho_{2k-1}(n)$ be either of the arithmetical functions $\beta_{2k-1}(n)$ or $\xi_{2k-1}(n)$ defined by (8) and (10) respectively; then $\rho_{2k-1}(n)$ satisfies a recurrence relation of the form:

$$(23) \quad \rho_{2k+1}(n) - n \rho_{2k-1}(n) = 2 \sum_{s=0}^{k-1} \sum_{j=1}^{n-1} \binom{2k}{2s+1} \rho_{2s+1}(j) \rho_{2k-2s-1}(n-j) ,$$