

REMARKS ON A THEOREM OF A. WINTNER

Autor(en): **Wong, James S. W.**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **13 (1967)**

Heft 1: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **12.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-41531>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

REMARKS ON A THEOREM OF A. WINTNER *

by JAMES S. W. WONG

Consider the following general second order differential equation

$$(1) \quad x'' + f(t, x) = 0, \quad t \geq 0,$$

where $f(t, x)$ is sufficiently smooth to guarantee existence of solutions. We are here concerned with the following result of Wintner [2] concerning the non-existence of L^2 solutions of the linear equation:

$$(2) \quad x'' + a(t)x = 0,$$

where $a(t)$ is a continuous function of t over the non-negative real axis $t \geq 0$.

THEOREM A (Wintner). *If*

$$(3) \quad \int_0^{\infty} t^3 |a(t)|^2 dt < \infty,$$

then (2) cannot have any non-trivial L^2 solutions.

Slightly modifying Wintner's original proof, Suyemoto and Waltman [1] obtain the following extension:

THEOREM B (Suyemoto and Waltman). *If (3) holds, then the differential equation*

$$(4) \quad x'' + a(t)x^p = 0,$$

where $p \geq 1$, cannot have any non-trivial L^{2p} solution.

The purpose of this note is to present an extension of Wintner's result for the more general equation (1). Although the basic idea of proof remains essentially the same as that of Wintner, we may by sharpening his argument, widen the applicability of Wintner's Theorem to a larger class of equations.

We assume throughout that $f(t, x)$ in addition satisfies

$$(5) \quad |f(t, x)| \leq a(t)\varphi(x),$$

*) This work was partially supported by a Summer Institute Fellowship held by the author during 1966 Summer Institute at Edmonton, and partially by the University of Alberta General Research Fund No. 644.

$\varphi(x)$ is a continuous function of x and $a(t)$ satisfies (3). Let $L\Phi$ denote the collection of all functions $x(t)$ on $[0, \infty)$, satisfying

$$(6) \quad \int_0^{\infty} |\varphi(x(t))|^2 dt < \infty.$$

THEOREM. *If the function $\varphi(u)$ satisfies*

$$(7) \quad \varphi(u) \neq 0 \quad \text{if} \quad u \neq 0,$$

$$(8) \quad \lim_{u \rightarrow \infty} \varphi(u) \geq \alpha > 0,$$

and for each L^2 solution $x(t)$ of (1), we have

$$(9) \quad \lim_{t \rightarrow \infty} \frac{\int_t^{\infty} |x(s)|^2 ds}{\int_t^{\infty} |\varphi(x(s))|^2 ds} > 0,$$

then equation (1) has no non-trivial $L\Phi$ solutions.

LEMMA. *Let $x(t)$ be a non-trivial $L\Phi$ solution of (1). Then $x(t)$ and its derivative $x'(t)$ tend to zero as $t \rightarrow \infty$.*

Proof. Integrating (1), we obtain

$$(10) \quad x'(t) - x'(s) = - \int_s^t f(\tau, x(\tau)) d\tau.$$

Using (3) and (5), we may estimate

$$(11) \quad |x'(t) - x'(s)|^2 \leq \left(\int_s^t |a(s)\varphi(x(s))|^2 ds \right)^2 \\ \leq \left(\int_s^t |a(s)|^2 ds \right) \left(\int_s^t |\varphi(x(s))|^2 ds \right).$$

Under the present hypothesis, the right hand side of (11) tends to zero as $t, s \rightarrow \infty$. Thus $\lim_{t \rightarrow \infty} x'(t)$ exists, say λ . If $\lambda \neq 0$, then $|x(t)| \rightarrow \infty$ as $t \rightarrow \infty$,

which on account of (8) contradicts the assumption (6). Since $x'(\infty) = 0$, we may now write for any $t_0 \geq 0$

$$\left(\int_{t_0}^{\infty} |x'(t)| dt \right)^2 \leq \left(\int_{t_0}^{\infty} \left| \int_t^{\infty} a(s)\varphi(x(s)) ds \right| dt \right)^2$$

$$(12) \quad \begin{aligned} &\leq \left(\int_{t_0}^{\infty} |s a(s) \varphi(x(s))| ds \right)^2 \\ &\leq \left(\int_{t_0}^{\infty} |s a(s)|^2 ds \right) \left(\int_{t_0}^{\infty} |\varphi(x(s))|^2 ds \right) < \infty, \end{aligned}$$

from which it follows that $x(t)$ is absolutely continuous, and in particular $\lim_{t \rightarrow \infty} x(t)$ exists, say μ . In case $\mu \neq 0$, we readily see that (7) contradicts the hypothesis (6). The proof is complete.

Proof of the theorem. From the fact that $|x(t)| \leq \int_t^{\infty} |x'(t)| dt$, we may estimate by (12) in the following manner.

$$(13) \quad \begin{aligned} \int_t^{\infty} |x(t)|^2 dt &\leq \int_t^{\infty} \left[\int_s^{\infty} |\tau a(\tau) \varphi(x(\tau))| d\tau \right]^2 ds \\ &\leq \int_t^{\infty} \left[\left(\int_s^{\infty} |\tau a(\tau)|^2 d\tau \right) \left(\int_s^{\infty} |\varphi(x(\tau))|^2 d\tau \right) \right] ds \\ &\leq \left(\int_t^{\infty} |\varphi(x(\tau))|^2 d\tau \right) \int_t^{\infty} \tau^3 |a(\tau)|^2 d\tau. \end{aligned}$$

Since $x(t)$ is non-trivial, we may divide $\int_t^{\infty} |\varphi(x(\tau))|^2 d\tau$ on both sides and take $\lim \sup$ which yields a contradiction on account of (9).

COROLLARY 1. If in place of (9), we have for some positive constant k_1

$$(14) \quad |\varphi(u)| \leq k_1 |u|^p \quad p \geq 1,$$

in some neighborhood of zero, say $\{u : |u| < \varepsilon_1, \varepsilon_1 > 0\}$, then the conclusion of the theorem remains valid.

Proof. Let $x(t)$ be a non-trivial $L\Phi$ solution of (1). From the above Lemma, we infer that $x(t) \rightarrow 0$ as $t \rightarrow \infty$. Thus, for sufficiently large t , we have on account of (14) that

$$(15) \quad |\varphi(x(t))| \leq k_1 |x(t)|^p \leq k_1 |x(t)|.$$

Substituting (15) in (9), we easily arrive at a contradiction.

COROLLARY 2. Let $\varphi(u)$ satisfy (7), (8), (14) and in addition

$$(16) \quad |\varphi(u)| \leq k_2 |u|^p$$

for some positive constant k_2 in some neighborhood of infinity say, $\{u : |u| > \frac{1}{\varepsilon_2}, \varepsilon_2 > 0\}$, where p is the same fixed number as given in (14).

Then (1) has no non-trivial L^{2p} solutions.

Proof. Let $x(t)$ be a non-trivial L^{2p} solution. Denote $A = \{t : |x(t)| < \varepsilon_1\}$, $B = \{t : |x(t)| > \frac{1}{\varepsilon_2}\}$ and $C = [0, \infty) - A - B$. Observe from (14) and (16) that

$$\begin{aligned} \int_0^{\infty} |\varphi(x(t))|^2 dt &= \int_A + \int_B + \int_C \\ &\leq \int_A k_1 |x(t)|^{2p} dt + \frac{1}{\varepsilon_2} \mu(C) + \int_B k_2 |x(t)|^{2p} dt \\ &< \infty, \end{aligned}$$

where $\mu(C)$ denotes the ordinary Lebesgue measure of C . Hence $x(t) \in L\Phi$. Following the proof of the main theorem, we again arrive at a contradiction, proving that (1) has no non-trivial L^{2p} solutions.

REMARK 1. Corollary 1 reduces to the result of Suyemoto and Waltman by taking $\varphi(u) = u^p$, $p \geq 1$ which reduces to the result of Wintner when $p = 1$.

REMARK 2. Let $\varphi(u) = 2u + \sin u$. It is easy to see that $\varphi(u)$ satisfies all (7), (8), (14) and (16). Hence, from Corollary 2, we may conclude that equation (1) has no square integrable solutions.

REFERENCES

- [1] SUYEMOTO, L. and P. WALTMAN, Extension of a theorem of A. Wintner. *Proc. Amer. Math. Soc.*, 14 (1963), 970-971.
- [2] WINTNER, A., A criterion for the non-existence of L^2 -solutions of a nonoscillatory differential equation. *J. London Math. Soc.*, 25 (1950), 347-351.

(Recu le 9 décembre 1966.)

University of Alberta
Edmonton, Alberta.

Added in Proof. For a closely related paper, see J. Burlak, "On the non-existence of L^2 solutions of a class of non-linear differential equations", *Proc. Edin. Math. Soc.*, 14(1965), 257-268, which contains several similar results.