

Zeitschrift: L'Enseignement Mathématique
Band: 14 (1968)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FLATNESS AND PRIVILEGE
Kapitel: §5. Geometrie applications of the flatness criterions
Autor: Douady, A.
DOI: <https://doi.org/10.5169/seals-42343>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 16.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Proof: $\text{Tor}_1^A(k, F) = \text{Tor}_1^B(\bar{B}, F)$, by prop. 7.

§ 5. *Geometric applications of the flatness criterions*

A) *Flatness for finite morphisms*

Proposition 1: Let $\pi: X \rightarrow S$ be a finite morphism (i.e. proper with finite fibres) of analytic spaces. Then $\pi_* (\mathcal{O}_X)$ is a coherent analytic sheaf over S . The following conditions are equivalent:

- (a) π is flat (i.e. for every $x \in X$, $\mathcal{O}_{X,x}$ is a flat $\mathcal{O}_{S,s}$ -module, $s = \pi(x)$).
- (b) For every s , $(\pi_* \mathcal{O}_X)_s$ is a flat $\mathcal{O}_{S,s}$ -module.
- (c) $\pi_* \mathcal{O}_X$ is a locally free sheaf.

Proof: Because π is finite $\pi_* (\mathcal{O}_X)_s = \bigoplus_{x \in \pi^{-1}(s)} \mathcal{O}_{X,x}$, thus the only point to prove is (b) \Rightarrow (c).

Now if $\mathcal{O}_{X,x}$ is a flat $\mathcal{O}_{S,s}$ -module, then (by theorem 2) $\mathcal{O}_{X,x}$ is free, and a coherent sheaf whose fibers are free is a locally free sheaf.

Proposition 2: Let S be a reduced analytic space and \mathcal{E} a coherent \mathcal{O}_S -module. Let $E(s)$ be the finite dimensional vector space (over \mathbb{C}) $\mathcal{E}_s \otimes_{\mathcal{O}_{S,s}} \mathbb{C}_s$. \mathcal{E} is a locally free $\mathcal{O}_{S,s}$ -module if and only if $\dim_{\mathbb{C}} E(s)$ is locally constant.

Proof: If \mathcal{E} is locally free, then $\dim_{\mathbb{C}} E(s)$ is locally constant. Suppose now that $\dim_{\mathbb{C}} E(s)$ is locally constant in an open set $U \subset S$, and that $\mathcal{O}_U^p \xrightarrow{d} \mathcal{O}_U^q \rightarrow \mathcal{E}_U \rightarrow 0$ is exact. d is determined by a $p \times q$ matrix of analytic functions on U , so it gives a morphism $\mathbb{C}_U^p \xrightarrow{d} \mathbb{C}_U^q$ of trivial vector bundles over U .

From the exact sequence $\mathcal{O}_s^p \xrightarrow{d_s} \mathcal{O}_s^q \rightarrow \mathcal{E}_s \rightarrow 0$, we get (by making tensor-products with \mathbb{C}_s) the exact sequence:

$$\mathbb{C}_s^p \xrightarrow{d(s)} \mathbb{C}_s^q \rightarrow E(s) \rightarrow 0,$$

which shows that d has constant rank in U . Thus $\text{Ker } d$ and $\text{Im } d$ are vector bundles, and we can write

$$\mathbb{C}_U^p = F_1 \oplus G_1, \quad \mathbb{C}_U^q = F_0 \oplus G_0,$$

$$d : \begin{cases} F_1 \rightarrow 0 \\ G_1 \simeq F_0 \end{cases}$$

Now $\mathcal{E} \simeq$ the sheaf of analytic sections of G_0 , therefore \mathcal{E} is locally free.

Definition 1: Let $\pi : X \rightarrow S$ be a finite morphism of analytic spaces, and $s \in S$. For each $x \in X(s) = \pi^{-1}(s)$, $\mathcal{O}_{X(s),x} = \mathbf{C} \otimes_{\mathcal{O}_{S,s}} \mathcal{O}_{X,x}$ is finite dimensional vectorspace over \mathbf{C} . Denote its dimension by $v(x)$. Then the degree $v(s)$ of s is defined by $v(s) = \sum_{x \in X(s)} v(x)$.

Theorem 1: Let $\pi : X \rightarrow S$ be a finite morphism of analytic space and let S be a reduced space. Then X is flat over S if and only if $v(s)$ is locally constant function of s .

$$\begin{aligned} \text{Proof: } v(s) &= \sum_{x \in X(s)} \dim_{\mathbf{C}} \mathcal{O}_{X(s),x} = \dim_{\mathbf{C}} \left(\bigoplus_{x \in X(s)} \mathcal{O}_{X(s),x} \right) \\ &= \dim_{\mathbf{C}} \left(\bigoplus_{x \in X(s)} (\mathbf{C} \otimes_{\mathcal{O}_{S,s}} \mathcal{O}_{X,x}) \right) \\ &= \dim_{\mathbf{C}} \mathbf{C} \otimes_{\mathcal{O}_{S,s}} \pi_* (\mathcal{O}_X)_s = \dim_{\mathbf{C}} E(s). \end{aligned}$$

The theorem follows from propositions 1 and 2.

Examples of flat morphisms

Example 1: If $\pi : X \rightarrow S$ is a local isomorphism near x , then π is flat at x .

Example 2: Consider § 2, Ex. 1. Here $v(x) = 1$.

Examples of non-flat morphisms

Examples 1: If $X \subset S$ is a closed subspace, not open, $v(s)$ is not locally constant.

Example 2: Let X be a subspace of \mathbf{C}^4 defined by the ideal intersection of (x_3, x_4) and $(x_1 - x_1, x_4 - x_2)$ (which is equal to the product ideal) and let π be the projection onto the (x_1, x_2) -plane \mathbf{C}^2 . Then X is a union of two 2-planes in \mathbf{C}^4 , whose intersection is (0) . When $s \neq 0$, $X(s)$ consists of two simple points, so $v(s) = 2$. $X(0)$ is given by the ideal $(x_1, x_2, x_3^2, x_3x_4, x_4^2)$, thus $v(0) = 3$.

Example 3: Let $S = \{(u, v, w) \in \mathbf{C}^3 \mid v^2 = uw\}$ and $\pi : \mathbf{C}^2 \rightarrow S$ be the map $(x, y) \rightarrow (x^2, xy, y^2)$. This map identifies S with the quotient of \mathbf{C}^2 by the equivalence relation identifying (x, y) with $(-x, -y)$. However, π is not flat, since for $s \in S$, $v(s) = 2$ if $s \neq 0$ and $v(s) = 3$ if $s = 0$.

B) *Projection of a product of analytic spaces*

Theorem 2: Let S and X be analytic spaces. If $\pi : S \times X \rightarrow S$ is the projection morphism, then π is flat, i.e. $\mathcal{O}_{S \times X, (s,x)}$ is a flat $\mathcal{O}_{S,s}$ module for every $(s, x) \in S \times X$.

To prove this theorem we need first some homological algebra. Then we shall show it in the particular case, when S is a manifold, and finally in the general case.

(a) *Koszul complex*

Let A be a ring, M an A -module and h_1, \dots, h_n homomorphisms $M \rightarrow M$, which commute with each other, i.e. $h_i h_j = h_j h_i$ for every i, j .

If $1 \leq k \leq n$, set $Q_k = M/h_1(M) + \dots + h_k(M)$, and $Q_0 = M$, thus, in particular, $Q_n = Q = M / \sum_{i=1}^n h_i(M)$. Every h_k induces a map $\tilde{h}_k : Q_{k-1} \rightarrow Q_{k-1}$.

Definition 2: The sequence (h_1, \dots, h_n) is called regular if each of the mappings \tilde{h}_k ($1 \leq k \leq n$) is injective.

The Koszul complex of the module M and of the mappings h_k ($1 \leq k \leq n$) $K. = K. [M; h_1, \dots, h_n]$ is defined in the following way:

$$K_i = \wedge^{n+i} A^n \otimes M \simeq M^{\binom{n}{i}}, \quad 0 \leq i \leq n.$$

We define the homomorphisms $d_i : K_i \rightarrow K_{i-1}$ ($i > 0$) by $\lambda \otimes x \rightarrow \sum_i (e_i \wedge \lambda) \otimes \otimes h_i(x)$, where (e_i) is the natural base of A^n . We also define $\varepsilon : K_0 \rightarrow Q$ as the natural map $: K_0 = M \rightarrow M / \sum_{i=1}^n h_i(M) = Q$. Using the fact that h_1, \dots, h_n commute with each other, it is easy to verify that

$$(d_{i-1} \circ d_i)(\lambda \otimes x) = \sum_{i,j} (e_j \wedge e_i \wedge \lambda) \otimes h_j(h_i(x)) = 0;$$

also $\varepsilon d_1 = 0$. Thus $K.$ is really a complex.

Theorem 3 (Poincaré-Koszul).

If (h_1, \dots, h_n) is a regular sequence, then

$$H_i(K.) = \begin{cases} Q & \text{if } i = 0 \\ \cdot & \cdot \\ 0 & \text{if } i > 0 \end{cases}$$

If $h_i \in A$, it defines the map: $A \xrightarrow{h_i I} A$, which we denote also by h_i . We say that (h_1, \dots, h_n) is a regular sequence of elements if $(h_1 I, \dots, h_n I)$ is a regular sequence.

Corollary. If (h_1, \dots, h_n) is a regular sequence of elements, then the Koszul complex $K. = K. [A; h_1, \dots, h_n] = \{ \wedge^{n-1} A^n \simeq A^{\binom{n}{i}} \}$ is a free resolution of $Q = A/(h_i)$ ((h_i) is the ideal generated by h_1, \dots, h_n).

Example: If $A = \mathbf{C} \{x_1, \dots, x_n\}$; $h_i = x_i$, then $Q_k = A/(x_1, \dots, x_k) = \mathbf{C} \{x_{k+1}, \dots, x_n\}$ and $Q = Q_n = \mathbf{C}$. The complex $K. = K. [A; x_1, \dots, x_n]$ is a free resolution of \mathbf{C} .

(b) *Proof of theorem 2, when S is a complex manifold*

In this case we can take $\mathcal{O}_{S,S} = \mathbf{C} \{t_1, \dots, t_m\} = A$ and if $\mathcal{O}_{X,x} = \mathbf{C} \{x_1, \dots, x_n\}/(f_1, \dots, f_p)$, then

$$\mathcal{O}_{S \times X, (s,x)} = \mathbf{C} \{t_1, \dots, t_m, x_1, \dots, x_n\}/(f_1, \dots, f_p) = B.$$

B is an A -module in a natural way.

By the corollary of the Poincaré-Koszul theorem $K. = K. [A; t_1, \dots, t_m]$ in a free resolution of \mathbf{C} . We want to compute the modules $\text{Tor}_i^A(\mathbf{C}, B) = H_i(K. \otimes B)$ ($i > 0$).

It's easily seen, that we can consider the complex $K. \otimes B$ as a Koszul

complex $K'. = K. [B; t_1, \dots, t_m]$ (where $t_i : B \xrightarrow{t_i I} B$). But now the sequence (t_1, \dots, t_m) is regular, thus by the Poincaré-Koszul theorem $H_i[K'] = 0$ if $i > 0$.

In particular: $\text{Tor}_1^A(\mathbf{C}, B) = H_1[K. \otimes B] = H_1[K'] = 0$. By the second flatness criterion B is A -flat.

(c) *The general case*

The question being local, we can suppose that $S \subset W \subset \mathbf{C}^n$, where W is open, and S an analytic subspace of W . Let S be defined by g_1, \dots, g_r . Then $S \times X \subset W \times X$ and $\mathcal{O}_S = \mathcal{O}_W/(g_1, \dots, g_r)$. On the other hand $\mathcal{O}_{S \times X} = \mathcal{O}_{W \times X}/(g_1, \dots, g_r) = \mathcal{O}_S \otimes_{\mathcal{O}_W} \mathcal{O}_{W \times X}$. The last equality follows from

the fact, that if $\pi : X \rightarrow S$ is a morphism, and $S' \subset S$ a subspace, $X' = \pi^{-1}(S')$,

$$\text{then } \mathcal{O}_{X'} = \mathcal{O}_{S'} \otimes_{\mathcal{O}_S} \mathcal{O}_X.$$

Remark : This a particular case of the following proposition: if π and π' are two morphisms of which at least one is finite, then

$$\begin{array}{ccc} X & & Y \\ \pi \searrow & & \swarrow \pi' \\ & S & \end{array} \quad \mathcal{O}_{X \times_S Y} = \mathcal{O}_X \otimes_{\mathcal{O}_S} \mathcal{O}_Y.$$

We have proved that $\mathcal{O}_{W \times X}$ is \mathcal{O}_W -flat, so by scalar extension $\mathcal{O}_{S \times X}$ is \mathcal{O}_S flat.

Corollary : If X and S are two manifolds and $\pi : X \rightarrow S$ is a submersion, then π is flat.

III. PRIVILEGED POLYCYLINDERS

§ 1. Banach vector bundles over an analytic space

Let E be a Banach space and X an analytic space. We denote then by E_X the trivial bundle $X \times E$ over X .

To define bundle morphisms, we first define the sheaf $\mathcal{H}_X(E)$ of germs of analytic morphisms from X to E . If $U \subset \mathbb{C}^n$ is open, then the set $\mathcal{H}(U, E)$ of analytic morphisms from U into E consists of all functions $g : U \rightarrow E$ having at every point $x \in U$ a converging power series expansion.

Let now X' be a local model for X , i.e. X' is the support of the quotient sheaf \mathcal{O}_U/J , where $U \subset \mathbb{C}^n$ is open and J is a coherent sheaf of ideals of \mathcal{O}_U , then $\mathcal{H}_{X'}(E)$ is the sheaf associated to the presheaf $V \rightarrow \mathcal{H}(V, E)/J_V \cdot \mathcal{H}(V, E)$ ($V \subset U$, V -open).

Remark : If X' is reduced, the sections of $\mathcal{H}_{X'}(E)$ are just the functions from X' to E which are locally induced by analytic functions on open sets in U .

The sheaf $\mathcal{H}_X(E)$ is constructed with help of the local models X' of X , i.e. $\mathcal{H}_X(E)|_{X'} = \mathcal{H}_{X'}(E)$, for every local model X' .

Definition 1 : The set of *analytic morphisms* from an analytic space X into a Banach space E is the set $\mathcal{H}(X; E)$ of sections of the sheaf $\mathcal{H}_X(E)$.

Let $\mathcal{L}(E, F)$ be the Banach space of all continuous linear mappings from the Banach space E into the Banach space F .

Definition 2 : An *analytic vector bundle morphism* from E_X into F_X is an analytic morphism from X into $\mathcal{L}(E, F)$.