

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 22 (1976)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SUPERDIAGONALIZATION AND REAL NORMAL OPERATORS
Autor: Geissinger, Ladnor
DOI: <https://doi.org/10.5169/seals-48179>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 23.01.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

SUPERDIAGONALIZATION AND REAL NORMAL OPERATORS

by Ladnor GEISSINGER

We shall give short direct proofs of all the usual simultaneous superdiagonalization and diagonalization theorems found in linear algebra textbooks, as well as determine the structure of real normal operators. The proofs are all variations on the technique of superdiagonalization, a method well-known to be very useful not just for elementary linear algebra but also in the theory of finite dimensional Lie (and also associative) algebras [Flanders].

Henceforth E will be a finite dimensional vector space over a field K and S will be a set of commuting operators on E . A subspace W of E is S -stable if $AW \subseteq W$ for all A in S . Our main tool is the following strong commutative version of Schur's lemma.

PROPOSITION 1. *If $\dim_K E > 1$, then either there is a proper S -stable subspace or the subalgebra S' generated by S over the scalars KI is a field and $\dim_K S' = \dim_K E$.*

Proof. Suppose first that some operator A in S (or S') has a minimal polynomial which is not irreducible over K ; so $m_A = pq$ where p and q are not constants. Since $p(A) \neq 0$ there is a vector v such that $w = p(A)v \neq 0$. Then $q(A)w = 0$ but $q(A) \neq 0$ so the kernel W of $q(A)$ is a proper subspace. Moreover, W is S -stable since if B is in S then $q(A)B(W) = Bq(A)(W) = 0$. Of course if all the operators in S are scalars (i.e. $S \subseteq KI$) every K -subspace is S -stable. Thus the only remaining case is when for each non-scalar A in S , m_A is irreducible over K of degree bigger than 1. Then the subalgebra K' generated by one such A over KI is a field and E is a vector space over K' of dimension less than $\dim_K E$ (in fact, $(\dim_{K'} E)(\dim_K K') = \dim_K E$). Moreover, the operators in S which are not in K' are K' -linear since they commute with A . Thus the proof can be completed by induction on the dimension of E .

We really only need the following special cases of Proposition 1 which result from successive restriction of S to S -stable subspaces.

COROLLARY. *If each of the operators in S has all of its characteristic roots in K , then a minimal S -stable subspace has dimension 1 over K . If $K = \mathbf{R}$, then a minimal S -stable subspace has dimension 1 or 2 over \mathbf{R} .*

Note that the first statement uses only the first half of the proof of Proposition 1. Also, when $K = \mathbf{R}$, the second half of the proof may be more transparent if E is considered to be a vector space over \mathbf{C} by the isomorphism $s + ti \rightarrow sI + t \frac{A - aI}{b}$ where $m_A(X) = (X - a)^2 + b^2$ and then the proof is completed by appeal to the result already proved for \mathbf{C} in the first statement.

A chain of subspaces $0 = W_0 < W_1 < W_2 < \dots < W_r = E$ will be called a *flag*, or a *fan* if $\dim W_k/W_{k-1} = 1$ for all k , and a basis x_1, \dots, x_n will be called a *flag (fan) basis* if $x_1, x_2, \dots, x_{d(k)}$ is a basis for W_k for each k , where $d(k) = \dim W_k$. A flag is S -stable if each W_k is S -stable.

PROPOSITION 2. *If each operator in S has all of its characteristic roots in K , then there is an S -stable fan. If $K = \mathbf{R}$ there is an S -stable flag $0 = W_0 < W_1 < \dots < W_r = E$ in which $\dim W_k/W_{k-1}$ is 1 or 2 for each k .*

Proof. Apply Proposition 1 or its Corollary to get an S -stable subspace W ; then apply it again to S restricted to W and/or to the operators induced by S on the quotient space E/W .

Note that in the first case, if x_1, \dots, x_n is a fan basis, then the matrix of each A in S relative to this basis is *superdiagonal* (upper triangular). In the second case, if x_1, \dots, x_n is a flag basis then the matrix of each A in S relative to this basis is *nearly superdiagonal*, that is, there are 1×1 and 2×2 blocks down the diagonal and zeros below. Translating these statements entirely into the language of matrices gives the following result.

COROLLARY. *Let S be a commuting set of matrices over K . If each matrix in S has all its characteristic roots in K , there is an invertible matrix P with entries in K such that $P^{-1}AP$ is superdiagonal for all A in S . If $K = \mathbf{R}$, there is an invertible real matrix P such that $P^{-1}AP$ is nearly superdiagonal for all A in S .*

Recall that a polynomial with coefficients in K is *separable* if it factors into a product of distinct irreducible factors over K (and over every field extension of K if characteristic $(K) > 0$), that is, has no multiple roots in

any field extension of K . Call an operator *separable* if its minimal polynomial is separable. If the minimal polynomial of a separable operator A in S factors as $m_A = qp$ over K , there are polynomials r and s such that $rp + sq = 1$. Then $I = r(A)p(A) + s(A)q(A)$ so E is the direct sum of $W = \text{Ker } q(A)$ and $V = \text{Ker } p(A)$.

As in the proof of Proposition 1 it follows that W and V are both S -stable. This gives a strengthened form of Proposition 1.

PROPOSITION 3. *If $\dim_K E > 1$ and if all the operators in S are separable, then either there are proper supplementary S -stable subspaces or the subalgebra S' generated by S over the scalars KI is a field (separable over K) and $\dim_K S' = \dim_K E$.*

COROLLARY. *If each operator in S is separable and has all of its characteristic roots in K , then E is the direct sum of 1 dimensional S -stable subspaces. If $K = \mathbf{R}$, then E is the direct sum of 1 or 2 dimensional S -stable subspaces, provided every operator in S is separable.*

In the first case, if $E = W_1 \oplus W_2 \oplus \dots \oplus W_n$ with the W_i being S -stable and if x_i spans W_i , then each x_i is a common eigenvector for all A in S (or S') and the matrix of each A relative to this basis is diagonal. In the second case, $E = W_1 \oplus \dots \oplus W_k \oplus V_1 \oplus \dots \oplus V_r$ where $\dim W_i = 1$ and $\dim V_i = 2$ and the W_i and V_j are minimal S -stable subspaces. If x_i spans W_i and y_j, z_j span V_j , then the matrix of each A relative to the basis $x_1, \dots, x_k, y_1, z_1, y_2, z_2, \dots, y_r, z_r$ is block diagonal with k eigenvalues followed by r 2×2 -blocks down the diagonal. This gives the usual simultaneous diagonalization theorem for commuting (diagonalizable) matrices, as well as the criterion for diagonalizability of a single operator.

Next suppose K is \mathbf{C} or \mathbf{R} and E has an inner product \langle, \rangle , and S is a commuting set of normal operators on E . Recall that A is *normal* if it commutes with its adjoint A^* . Now for a normal A , if $m_A = pq$ and if $W = \text{Ker } q(A)$ then W is A^* -stable and so the orthogonal complement W^1 is A -stable since $\langle W, AW^1 \rangle = \langle A^*W, W^1 \rangle = 0$. But $p(A)(W^1) \subseteq \text{Ker } q(A) = W$ so $W^1 \subseteq \text{Ker } p(A)$ and hence l.c.m. $(p, q) = m_A$, from which we conclude that p and q are relatively prime and $W^1 = \text{Ker } p(A)$. Thus W and W^1 are S -stable, and every normal operator is separable. A *root space* for S is a maximal proper S -stable subspace W such that each A in S acts as a scalar on W , that is, W is the intersection of a collection of eigenspaces, one for each A in S .

PROPOSITION 4. *If each operator in S is normal and has all of its characteristic roots in K , then E is the orthogonal direct sum of root spaces for S . If $K = \mathbf{R}$ and each operator in S is normal, then E is the orthogonal direct sum of root spaces for S and S -stable subspaces V on which each A in S acts as a scalar or has irreducible (quadratic) minimal polynomial.*

Upon choosing an orthonormal basis for each root space, the first statement of Proposition 4 yields the simultaneous unitary diagonalization of commuting normal operators (or matrices) when $K = \mathbf{C}$. To see that it also yields the simultaneous real orthogonal diagonalization of commuting real symmetric operators (or matrices), it is only necessary to note that in this case all the characteristic roots are real.

LEMMA. *All characteristic roots of a real symmetric operator are real.*

Proof. If A is symmetric and $p(X) = (X-a)^2 + b$ is a factor of the minimal polynomial of A there is a vector $v \neq 0$ such that $p(A)(v) = 0$. Then $0 \leq \langle (A-aI)(v), (A-aI)(v) \rangle = \langle (A-aI)^2(v), v \rangle = (-b) \langle v, v \rangle$ so $b \leq 0$ and p has real roots.

For real normal operators we need only consider a subspace V of the kind in the second statement of Proposition 4. If A is normal with minimal polynomial $(X-a)^2 + b^2 = p(X)$ on V , there is by the Corollary of Proposition 1 a minimal 2-dimensional subspace W of V stable under the commuting operators $\frac{A + A^*}{2}$ (symmetric) and $\frac{A - A^*}{2}$ (skewsymmetric).

Relative to an orthonormal basis y_1, y_2 of W , the matrix of $\frac{A + A^*}{2}$ must

be $\begin{pmatrix} a' & 0 \\ 0 & a' \end{pmatrix}$, the matrix of $\frac{A - A^*}{2}$ must be $\begin{pmatrix} 0 & -b' \\ b' & 0 \end{pmatrix}$, and so the matrix of A

must be $\begin{pmatrix} a' & -b' \\ b' & a' \end{pmatrix}$. Then $p(X) = X^2 - 2a'X + (a')^2 + (b')^2$ so that

$a = a'$ and $\pm b = b'$. Since W^1 is also stable under the symmetric operator $\frac{A + A^*}{2}$ and the skew-symmetric operator $\frac{A - A^*}{2}$, both W and W^1 are

A -stable. It follows that V is the orthogonal sum of such subspaces W , that

$\frac{A + A^*}{2} = aI$ on V , that $\left(\frac{A - A^*}{2}\right)^2 = -b^2I$ on V , and that $P = (a^2 + b^2)^{-\frac{1}{2}}A$

is orthogonal on V . Since $P^* = P^{-1}$ is a polynomial in P , A^* is a polynomial in A and hence for every A -stable subspace W of V , W is A^* -stable and W^1 is A -stable. In particular, the Corollary of Proposition 1 gives a

2-dimensional S -stable subspace W_1 of V , and by the preceding argument W_1^\perp is also S -stable. Thus V is the orthogonal sum of 2-dimensional S -stable subspaces W_1, \dots, W_r . Let y_i, z_i be an orthonormal basis for W_i , then relative to the basis $y_1, z_1, \dots, y_r, z_r$ the matrix of each A in S is block diagonal and the blocks are positive multiples of 2×2 rotation matrices. Thus we have determined the structure of real normal operators.

Finally, return to Proposition 2 and suppose that y_1, \dots, y_n is a fan (flag) basis for an S -stable fan (flag) and that E has an inner product. The Gram-Schmidt process applied to this basis yields an orthonormal fan (flag) basis z_1, \dots, z_n for the same S -stable fan (flag). Thus the matrix P in the Corollary can be taken to be unitary or real orthogonal. Moreover, if S contains A and A^* for some A (hence A is normal) we get directly that the matrices of A and A^* relative to the basis z_1, \dots, z_n are both (nearly) super-diagonal, and since one is the adjoint of the other, they are both diagonal (block diagonal with blocks at most 2×2 in size). This observation could be used to give another proof for Proposition 4 and the structure of real normal operators. In either case, the argument can be simplified a bit if S is assumed to be $*$ -closed.

REFERENCES

- FLANDERS, H. Methods of Proof in Linear Algebra. *American Mathematical Monthly*, January 1956, pp. 1-15.
HOFFMAN, K. and R. KUNZE. *Linear Algebra*. Second edition, Prentice-Hall, 1971.

(Reçu le 16 octobre 1975)

Ladnor Geissinger

Mathematics Department
University of North Carolina
Chapel Hill, North Carolina 27514

Vide-leer-empty