

1. Chevalley's Theorem

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **24 (1978)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **12.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

(1.10) and Lemma 1.2 yield

$$(1.11) \quad \delta_m = \frac{1}{|G|} \sum_{\sigma \in G} (\text{Tr } \sigma)_m = \frac{1}{|G|} \sum_{\sigma \in G} \sum_{|a|=m} \omega^a(\sigma).$$

Multiply both sides of (1.11) by t^m and sum over m from 0 to ∞ . We get

$$\begin{aligned} \sum_{m=0}^{\infty} \delta_m t^m &= \frac{1}{|G|} \sum_{m=0}^{\infty} \sum_{\sigma \in G} \sum_{|a|=m} \omega^a(\sigma) t^m \\ &= \frac{1}{|G|} \sum_{\sigma \in G} \left\{ \sum_{m=0}^{\infty} \omega_1^m(\sigma) t^m \dots \sum_{m=0}^{\infty} \omega_n^m(\sigma) t^m \right\} \\ &= \frac{1}{|G|} \sum_{\sigma \in G} \frac{1}{(1 - \omega_1(\sigma) t) \dots (1 - \omega_n(\sigma) t)} \end{aligned}$$

CHAPTER II

INVARIANT THEORETIC CHARACTERIZATION OF FINITE REFLECTION GROUPS

1. CHEVALLEY'S THEOREM

We showed in chapter I that we can always find a finite number of homogeneous invariants forming a basis for the invariants of G and that this set must contain at least n elements, where $n = \dim V$. We show that this lower bound is attained only for the finite reflection groups. We first define these groups.

DEFINITION 2.1. Let σ be a linear transformation acting on the n -dimensional vector space V . σ is a reflection $\Leftrightarrow \sigma$ fixes an $n - 1$ dimensional hyperplane π and σ is of finite order > 1 . π is called the reflecting hyperplane (r.h.) of σ .

REMARK. Choose $v \notin \pi$. and let $\sigma v = \zeta v + p$, $p \in \pi$. If $\zeta = 1$, then $\sigma^m v = v + mp$, contradicting that σ is of finite order. Hence $\zeta \neq 1$. Let $v' = v + (\zeta - 1)^{-1} p$ and choose p_1, \dots, p_{n-1} as a basis for π . Then $\sigma p_i = p_i$, $1 \leq i \leq n - 1$, $\sigma v' = \zeta v'$. ζ is a root of 1 in k which is distinct from 1, as σ is of finite order > 1 . Thus σ is a reflection iff relative to some basis, the matrix for σ is diagonal, $n - 1$ of the diagonal entries equalling 1 and the remaining one equalling a root of 1 in k distinct from 1.

DEFINITION 2.2. G is a finite reflection group acting on $V \Leftrightarrow G$ is a finite group generated by reflections on V .

As an example of a finite reflection group, let $G = S_n$. It is well known that S_n is generated by transpositions. The transposition of the variables x_i, x_j ($i \neq j$) fixes the hyperplane $x_i - x_j = 0$, so that it is a reflection.

We have the following result

THEOREM 2.1 (Chevalley [4]). *Let G be a finite reflection group acting on the n -dimensional vector space V . The invariants of G have a basis consisting of n homogeneous elements which are algebraically independent over k .*

Let $k[x]$ denote the ring of polynomials in x_1, \dots, x_n with coefficients in k . We prove the following.

LEMMA 2.1. Let I_1, \dots, I_m be invariant polynomials of G , $I_1 \notin (I_2, \dots, I_m)$ = the ideal in $k[x]$ generated by I_2, \dots, I_m . Suppose that $P_1 I_1 + \dots + P_m I_m = 0$, the P_i 's being polynomials with P_1 homogeneous. Then $P_1 \in \mathcal{I}$, where \mathcal{I} is the ideal in $k[x]$ generated by the homogeneous invariants of positive degree.

Proof of Lemma 2.1. The proof proceeds by induction on $\deg P_1$. Suppose $\deg P_1 = 0$, so that $P_1 = c \in k$. If $c \neq 0$, then $I_1 \in (I_2, \dots, I_m)$, contrary to assumption. Hence $c = 0 \Rightarrow P_1 \in \mathcal{I}$. Let $\deg P_1 = n > 0$. Let σ be a reflection in G and $L = 0$ the equation of its r.h. (L is a linear homogeneous polynomial). We have $P_1(x) I_1(x) + \dots + P_m(x) I_m(x) = 0$, $P_1(\sigma x) I_1(x) + \dots + P_m(\sigma x) I_m(x) = 0$. Hence $[P_1(\sigma x) - P_1(x)] I_1(x) + \dots + [P_m(\sigma x) - P_m(x)] I_m(x) = 0$. For $L(x) = 0$, $\sigma(x) = x$, so that $P_i(\sigma x) - P_i(x) = 0$ whenever $L(x) = 0$, $1 \leq i \leq m$. Since $L(x)$ is irreducible it follows that

$$\frac{P_i(\sigma x) - P_i(x)}{L(x)}$$

is a polynomial, $1 \leq i \leq m$. We have

$$\left[\frac{P_1(\sigma x) - P_1(x)}{L(x)} \right] I_1(x) + \dots + \left[\frac{P_m(\sigma x) - P_m(x)}{L(x)} \right] I_m(x) = 0.$$

$$\deg \left[\frac{P_1(\sigma x) - P_1(x)}{L(x)} \right] < \deg P_1(x),$$

so that by the induction hypothesis

$$\frac{P_1(\sigma x) - P_1(x)}{L(x)} \equiv 0 \pmod{\mathcal{I}}.$$

Hence $P_1(\sigma x) \equiv P_1(x) \pmod{\mathcal{I}}$. Since the σ 's generate G , this congruence holds for $\sigma \in G$. We conclude that

$$P_1(x) \equiv \frac{1}{|G|} \sum_{\sigma \in G} P_1(\sigma x) \pmod{\mathcal{I}}.$$

The polynomial $\frac{1}{|G|} \sum_{\sigma \in G} P_1(\sigma x)$ is invariant and homogeneous of degree $n \geq 1$. Hence it $\in \mathcal{I}$, so that $P_1 \in \mathcal{I}$.

Proof of Theorem 2.1. We choose I_1, \dots, I_r to be homogeneous invariants of positive degree forming a minimal basis for \mathcal{I} . Hilbert's proof of Theorem 1.1 shows that I_1, \dots, I_r form a basis for the invariants of G . We show that I_1, \dots, I_r are algebraically independent, so that $r = n$.

Suppose, to the contrary, that I_1, \dots, I_r are algebraically dependent. Choose $H(y_1, \dots, y_r)$ to be a polynomial of minimal positive degree so that $H(I_1(x), \dots, I_r(x)) = 0$. Let x -degree of any monomial $y_1^{a_1} \dots y_r^{a_r}$ be $d_1 a_1 + \dots + d_r a_r$, where $d_i = \deg I_i$. We may assume that all x -degrees of the monomials appearing in H are the same. Let

$$H_i(x) = \frac{\partial H}{\partial y_i}(I_1(x), \dots, I_r(x)), \quad 1 \leq i \leq r.$$

The H_i 's are invariant homogeneous polynomials, as all monomials in H have equal x -degree. Since $H(y_1, \dots, y_r)$ is of positive degree, some $\frac{\partial H}{\partial y_i} \neq 0$. It follows that the corresponding $H_i(x) \neq 0$, as H was chosen

to be of minimal degree; i.e. not all H_i 's = 0. We relabel indices so that $H_1, \dots, H_s, 1 \leq s \leq r$, are ideally independent (i.e. none of the H_i 's is in the ideal generated by the others) and $H_{s+j} \in (H_1, \dots, H_s), 1 \leq j \leq r - s$.

Thus $H_{s+j} = \sum_{i=1}^s V_{ji} H_i, 1 \leq j \leq r - s$, where each V_{ji} is a homogeneous polynomial of degree $d_i - d_{s+j}$ (V_{ji} is interpreted to be 0 if this degree is negative). Differentiating the relation $H(I_1(x), \dots, I_r(x)) = 0$ with respect to x_k , we obtain

$$\begin{aligned} (2.1) \quad \sum_{i=1}^r H_i \frac{\partial I_i}{\partial x_k} &= \sum_{i=1}^s H_i \frac{\partial I_i}{\partial x_k} + \sum_{l=1}^{r-s} H_{s+l} \frac{\partial I_{s+l}}{\partial x_k} \\ &= \sum_{i=1}^s H_i \left[\frac{\partial I_i}{\partial x_k} + \sum_{l=1}^{r-s} V_{li} \frac{\partial I_{s+l}}{\partial x_k} \right] = 0. \end{aligned}$$

Since

$$\frac{\partial I_i}{\partial x_k} + \sum_{l=1}^{r-s} V_{li} \frac{\partial I_{s+l}}{\partial x_k}$$

is homogeneous of degree $d_i - 1$, we conclude from Lemma 2.1 that

$$(2.2) \quad \frac{\partial I_i}{\partial x_k} + \sum_{l=1}^{r-s} V_{li} \frac{\partial I_{s+l}}{\partial x_k} = \sum_{j=1}^r B_j I_j, \quad 1 \leq i \leq s,$$

where the B_j 's are homogeneous and each term in (2.2) is homogeneous of degree $d_i - 1$. This forces $B_i = 0$. Multiply both sides of (2.2) by x_k and sum over k . We conclude, by Euler's identity for homogeneous polynomials,

$$(2.3) \quad d_i I_i + \sum_{l=1}^{r-s} V_{li} d_{s+l} I_{s+l} = \sum_{j=1}^r A_j I_j,$$

the A_j 's being homogeneous with $A_i = 0$.

(2.3) shows that $I_i \in (I_1, \dots, I_{i-1}, I_{i+1}, \dots, I_r)$, contradicting the minimality of the basis I_1, \dots, I_r . Hence I_1, \dots, I_r are algebraically independent and $r = n$.

2. THE THEOREM OF SHEPHARD AND TODD

We obtain in this section a converse to Chevalley's Theorem, thereby obtaining an invariant theoretical characterization of finite reflection groups. We first prove several preliminary results.

LEMMA 2.2. Let H be a finite group of linear transformations acting on the n -dimensional space V and fixing the $n - 1$ dimensional hyperplane π . The elements of H have a common eigenvector $v \in V - \pi$. Let $\sigma(v) = \zeta(\sigma)v$, $\sigma \in H$. $\zeta(\sigma)$ is an isomorphism from H into the multiplicative group of the roots of unity in k . It follows that H is a cyclic group.

REMARK. The above lemma is a consequence of Maschke's Theorem proven in section 2.3. We provide another proof below.

Proof. Let $\sigma_1 \in H$, $\sigma_1 \neq e$ (the identity of H). By the remark following Definition 2.1, there exists $v \in V - \pi$ such that $\sigma_1(v) = \zeta_1 v$, ζ_1 being a root of unity $\neq 1$. For $\sigma \in H$, let $\sigma(v) = \zeta(\sigma)v + p(\sigma)$, $\zeta(\sigma) \in k$ and $p(\sigma) \in \pi$. Let $\sigma^* = \sigma_1^{-1} \sigma^{-1} \sigma_1 \sigma$. Then $\sigma^*(v) = v + (1 - \zeta_1)p(\sigma)$. Since σ^* is of finite order, $(1 - \zeta_1)p(\sigma) = 0 \Rightarrow p(\sigma) = 0$. Hence $\sigma(v) = \zeta(\sigma)v$. $\zeta(\sigma)$ is clearly an isomorphism from H into U , the multiplicative group of