

10. SOME OTHER PROBLEMS

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **24 (1978)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **13.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

<http://www.e-periodica.ch>

which are models for the maps $\Gamma_M \rightarrow \Gamma'_{M,N}$ and $\Gamma'_{M,N} \rightarrow \Gamma'_M$. The first one is obvious and the second one is completely characterized by the map $WU_n \rightarrow WU_{n,p}$.

Now we get the differential on $\Lambda(x^j_\alpha, y^j_\lambda)$ by considering this algebra as the tensor product over $\Lambda(z^j_\alpha)$ of $\Lambda(x^j_\alpha, z^j_\lambda)$ with $\Lambda(y^j_\lambda)$.

One can make a similar construction using for A and B the *DG*-algebras Ω_M and Ω_N of differential forms on M and N . Of course one has to work again in more intrinsic terms and use the C^∞ -topology on Ω_M and Ω_N (compare with [13]). In this way one gets a *DG*-algebra which is also a model for $\Gamma_{M,N}$ (in fact one proves directly that it is a model for the *DG*-algebra constructed above), with a map in $C^*(L_{M,N})$ inducing an isomorphism in cohomology.

Summing up, we get the following result.

THEOREM. *Assume that the inclusion of N in M has a model which is a surjection of finite dimensional *DG*-algebras. One can construct explicitly a model for $C^*(L_{M,N})$ which is finite dimensional in each degree.*

Example. Suppose that M is the disk D^2 and N its boundary $\partial D^2 = S^1$. As the inclusion of $F_{2,1}$ in F_2 is homotopically trivial (equivalently the morphism $WU_2 \rightarrow WU_{2,1}$ is homotopic to zero), the bundle $\Gamma_{M,N} \rightarrow \Gamma'_{M,N}$ is trivial. WU_2 is a model for $S^5 \vee S^5 \vee S^7 \vee S^8 \vee S^8$ and $WU_{2,1}$ for $S^3 \vee S^3 \vee S^3 \vee S^4 \vee S^4$.

Hence $C^*(L_{D^2, \partial D^2})$ is a model for the space which is the product of the space of maps of S^1 in $S^3 \vee S^3 \vee S^3 \vee S^4 \vee S^4$ with the second loop space of $S^5 \vee S^5 \vee S^7 \vee S^8 \vee S^8$.

One can write down quite explicitly the minimal model for that space, but it is harder to compute the cohomology of the first factor. It has an infinite number of multiplicative generators.

10. SOME OTHER PROBLEMS

1. As coefficient for the Gelfand-Fuks cochains, one might consider, instead of the field R with the trivial action of L_M , a topological L_M -algebra A . The problem is to find a model for the *DG*-algebra $C^*(L_M, A)$ of continuous multilinear alternate forms on L_M with values in A . The differential is defined by the usual formula involving the action of L_M on A .

For that case, results similar to the one mentionned in this report have been obtained by Fuks-Segal (unpublished) and by T. Tsujishita [21].

For instance, when A is the algebra of smooth functions on M on which L_M acts by Lie derivative, their result is as follows. As it is described in § 3, the bundle E over M has a fiber F_n which is itself a principal U_n -bundle. Let us fix a fiber $F_n^\circ \approx U_n$ of this bundle; as it is invariant by the structural group $O_n \subset U_n$ of E , we get a subbundle E_o of E with typical fiber F_n° . Then $C^*(L_M, A)$ will be a model for the inverse image of E_o by the evaluation map $M \times \Gamma \rightarrow E$.

2. One of the most interesting problems is to know when, for a given class α in $H^*(L_M)$, there is a space X and a foliation F on $X \times M$ transverse to the fibers such that the image of α in $H^*(X)$ by the characteristic homomorphism (cf. 2) is non zero.

Very recent and remarkable results of Fuchs [23] show that this is the case for all classes coming from WSO_n . (For earlier partial results, see [4].) One might expect that his method will apply in general and show that the answer is affirmative for all classes in $H^*(L_M)$ (and also for the similar problem with $H^*(L_M; G)$).

There is also the problem of the possible continuous variations of characteristic classes for flat bundles which would be interesting to study (cf. [23]).

REFERENCES

- [1] BOTT, R. A topological obstruction for integrability. *Proc. Symp. Pure Math.* 16 (1970), pp. 127-131.
- [2] — On the Gelfand-Fuks cohomology, Differential Geometry. *Proc. of Symp. A.M.S.* 27, Part I, pp. 357-364.
- [3] — Lectures on characteristic classes and foliations (Notes by Lawrence Conlon). *Lecture Notes in Mathematics* 279, pp. 1-94, Springer Verlag, New York, 1972.
- [4] — On the characteristic classes of groups of diffeomorphisms. *Ens. Math.* 23 (1977), pp. 209-220.
- [5] CARTAN, H. Exposés 19 et 20 du Séminaire Henri Cartan 1949-1950 (Espaces fibrés et homotopie).
- [6] GELFAND, I. M. and D. FUKS. The cohomology of the Lie algebra of vector fields on the circle. *Funct. Anal.* 3 (1968), pp. 92-93.
- [7] — The cohomology of the Lie Algebra of Tangent Vector fields on a Smooth Manifold I and II. *Funct. Anal.* 3 (1969), pp. 32-52 and 4 (1970), pp. 23-32.
- [8] — The Cohomology of the Lie Algebra of Formal Vector Fields. *Izvestia Ak. SSSR* 34 (1970), pp. 322-337.
- [9] GODBILLON, G. Cohomologie d'algèbres de Lie de champs de vecteurs formels. *Séminaire Bourbaki* 421 (1972-1973).