

2. Integral representation theorems for linear functionals

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **25 (1979)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **11.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

2. INTEGRAL REPRESENTATION THEOREMS FOR LINEAR FUNCTIONALS

Let A be a commutative Banach algebra over \mathbf{C} and let Δ denote the locally compact space of regular maximal ideals of A . For each $x \in A$ we use \hat{x} to denote the Gelfand-transform; i.e., \hat{x} is the continuous mapping from Δ to \mathbf{C} defined by the relations:

$$\hat{x}(m) = m(x) \quad \text{for } m \in \Delta.$$

By $C_0(\Delta)$ we shall denote the algebra of all complex-valued continuous functions on Δ which vanish at infinity. For any subset $\mathcal{A} \subset A$ we shall use the notation $\hat{\mathcal{A}}$ to denote the set $\{\hat{x} : x \in \mathcal{A}\}$. As usual $\|\hat{x}\|_\infty$ denotes the supremum norm.

THEOREM 1. Let f be a linear form on the complex commutative Banach algebra A and let \mathcal{A} be a linear subspace of A . The following two statements are equivalent:

(1) There exists a constant M such that

$$|f(x)| \leq M \|\hat{x}\|_\infty \quad \text{for every } x \in \mathcal{A}.$$

(2) There exists a bounded complex Radon measure μ on Δ such that

$$f(x) = \int_\Delta \hat{x}(m) d\mu(m) \quad \text{for every } x \in \mathcal{A}.$$

Proof. The implication (2) \Rightarrow (1) is clear with $M = \|\mu\|$. We shall prove (1) \Rightarrow (2). Define a mapping $L : \hat{\mathcal{A}} \rightarrow \mathbf{C}$ by

$$L(\hat{x}) = f(x).$$

It follows from (1) that L is well-defined, and that

$$|L(\hat{x})| \leq M \|\hat{x}\|_\infty \quad \text{for every } \hat{x} \in \hat{\mathcal{A}}$$

and so L is continuous with $\|L\| \leq M$. Using the Hahn-Banach Theorem we can extend L to a bounded linear form L_0 on $C_0(\Delta)$ and by the Riesz Representation Theorem we obtain the existence of a bounded complex Radon measure μ on Δ such that

$$\| \mu \| = \| L \| = \| L_0 \| \quad \text{and}$$

$$L_0(\varphi) = \int_{\Delta} \varphi(m) d\mu(m) \quad \text{for every } \varphi \in C_0(\Delta).$$

In particular

$$f(x) = L(\hat{x}) = L_0(\hat{x}) = \int_{\Delta} \hat{x}(m) d\mu(m) \quad \text{for every } x \in \mathcal{A}.$$

Remark: Suppose that A has an identity and that \hat{A} is closed under complex conjugation, then since \hat{A} contains constants and separates the points of Δ , the Stone-Weierstraß Theorem implies that \hat{A} is dense in $C(\Delta)$, the algebra of all complex-valued continuous functions on the compact Hausdorff space Δ . If we impose these additional conditions on A and if we take $\mathcal{A} = A$ in Theorem 1, we can conclude that in this case the representing measure μ is uniquely determined.

If the algebra A has a continuous involution, one can use Theorem 1 to derive an extended version of a theorem due to Raikov [10]. We proceed to describe the situation.

Let A be a complex commutative Banach algebra with an isometric involution $*$ and a bounded approximate identity $\{u_\lambda\}_{\lambda \in \Lambda}$ i.e., a net satisfying the following conditions:

$$\begin{aligned} \| u_\lambda \| &\leq 1 \quad \text{for each } \lambda \in \Lambda, \\ \| u_\lambda x - x \| &\rightarrow 0 \quad \text{for each } x \in A. \end{aligned}$$

A continuous *positive* functional on A is an element $f \in A'$ such that $f(x^*x) \geq 0$ for every $x \in A$. If f is a continuous positive functional on A then the Cauchy-Schwarz inequality is valid (Dixmier [8, p. 23]) and this implies the following facts:

$$\begin{aligned} f(u_\lambda) &\rightarrow \| f \| \\ | f(x) |^2 &\leq \| f \| f(x^*x) \quad \text{for every } x \in A. \end{aligned}$$

If the involution is *symmetric*, which means $(x^*)^\wedge = \overline{x}$ for every $x \in A$ or, equivalently, that every $m \in \Delta$ is a *positive* linear functional, then by modifying a classical method of Gelfand-Raikov-Silov [10; p. 62] one can prove that

$$| f(x) | \leq \| f \| \| \hat{x} \|_\infty \quad \text{for every } x \in A.$$

As a corollary to Theorem 1 and the above discussion we obtain the following extended theorem of Raikov [10; p. 64], see also Bucy-Maltese [4]):

THEOREM 2. Let A be a commutative Banach algebra with an isometric involution which is symmetric. Suppose that A has a bounded approximate identity and let $f \in A'$ be a continuous positive functional. Then there exists a unique positive Radon measure μ on Δ such that $\|\mu\| = \|f\|$ and

$$f(x) = \int_A \hat{x}(m) d\mu(m) \quad \text{for every } x \in A.$$

Proof. From the above remarks we know that

$$|f(x)| \leq \|f\| \|\hat{x}\|_\infty \quad \text{for every } x \in A.$$

By Theorem 1 there exists a complex Radon measure μ on Δ such that $\|\mu\| \leq \|f\|$ and

$$f(x) = \int_A \hat{x}(m) d\mu(m) \quad \text{for every } x \in A.$$

This formula implies

$$|f(x)| \leq \|\hat{x}\|_\infty \|\mu\| \leq \|x\| \|\mu\| \quad \text{for every } x \in A,$$

so that $\|f\| \leq \|\mu\|$ and hence $\|f\| = \|\mu\|$.

Since \hat{A} is a self-adjoint subalgebra of $C_0(\Delta)$ which separates points and for each $m \in \Delta$ contains a function \hat{x} such that $\hat{x}(m) \neq 0$ (in fact ¹⁾ there exists an element u_β of the approximate identity such that $\hat{u}_\beta(m) \neq 0$), the Stone-Weierstraß Theorem implies the uniqueness of the measure μ . The positivity of μ also follows from the fact that \hat{A} is dense in $C_0(\Delta)$. In fact if p is a non-negative function in $C_0(\Delta)$, then $p = |q|^2$ for some $q \in C_0(\Delta)$. Choose a sequence $\{x_n\}$ in A such that

$$\hat{x}_n \rightarrow q.$$

¹⁾ If $m \in \Delta$, then $\|m\| \neq 0$ and by the assumption of symmetry m is a positive functional. Therefore, as mentioned above, $\|m\| = \lim_{\alpha} m(u_\alpha)$ so that there must exist some u_β of the approximate identity such that $m(u_\beta) \neq 0$.

Since $(x_n^*)^\wedge = \overline{x_n}$ it follows that $(x_n^*)^\wedge \rightarrow \bar{q}$ and hence

$$(x_n x_n^*)^\wedge \rightarrow |q|^2 = p.$$

This implies

$$\begin{aligned} \int_{\Delta} p(m) d\mu(m) &= \lim_n \int_{\Delta} (x_n x_n^*)^\wedge(m) d\mu(m) \\ &= \lim_n f(x_n x_n^*) \geq 0, \end{aligned}$$

so that μ is a positive measure and this completes the proof.

If A has an identity, as is the case in Raikov's original version, the above proof can be somewhat simplified.

THEOREM 3 (Raikov). Let A be a complex commutative Banach algebra with an identity e and with an isometric involution which is symmetric. If f is a continuous positive functional on A , then there exists a unique positive Radon measure μ on Δ such that $\|\mu\| = \|f\|$ and

$$f(x) = \int_{\Delta} \hat{x}(m) d\mu(m) \quad \text{for every } x \in A.$$

Proof. As above we know that

$$|f(x)| \leq \|f\| \|\hat{x}\|_{\infty} \quad \text{for every } x \in A.$$

From Theorem 1 there exists a complex Radon measure μ on Δ such that $\|\mu\| \leq \|f\|$ and

$$f(x) = \int_{\Delta} \hat{x}(m) d\mu(m) \quad \text{for every } x \in A.$$

Hence $\|\mu\| \leq \|f\| = f(e) = \mu(1) \leq \|\mu\|$ so that $\mu(1) = \|\mu\|$ which is enough to imply that μ is positive. The uniqueness of μ follows as in the Remark to Theorem 1.

3. APPLICATIONS OF THE INTEGRAL REPRESENTATION THEOREMS

Application 1 (Bochner's Theorem). Let G be a locally compact abelian group and let \hat{G} denote the (locally compact) character group. Denote